

DOCTORAL THESIS

Hazardous air pollutants from the waste incineration industry: formation mechanisms, distribution characteristics, and potential environmental risks

Wang, Chen

Date of Award:
2020

[Link to publication](#)

General rights

Copyright and intellectual property rights for the publications made accessible in HKBU Scholars are retained by the authors and/or other copyright owners. In addition to the restrictions prescribed by the Copyright Ordinance of Hong Kong, all users and readers must also observe the following terms of use:

- Users may download and print one copy of any publication from HKBU Scholars for the purpose of private study or research
- Users cannot further distribute the material or use it for any profit-making activity or commercial gain
- To share publications in HKBU Scholars with others, users are welcome to freely distribute the permanent URL assigned to the publication

Abstract

In this study, the formation mechanisms, distribution characteristics, and potential environmental risks of hazardous air pollutants from industrial-scale waste incineration processes were investigated.

First, to clarify the dominant formation mechanism of polychlorinated dibenzo-p-dioxins and dibenzofurans (PCDD/Fs) in a hazardous waste (HW) incinerator, three tests were designed by adding different precursors in phenol-containing raw materials. With the addition of p-dichlorobenzene, PCDD/F levels at the quenching tower outlet were ten times higher than levels observed at the inlet. This indicates that the quenching tower failed to suppress the formation of PCDD/Fs and surface-mediated precursor reaction is the dominant formation mechanism in low-temperature stages. Besides, adsorptive memory effect in air pollution control devices (APCDs) also led to high PCDD/F emissions. These findings suggest that to control PCDD/F emissions, strict regulation of chlorine contents in feed materials and frequent cleaning of APCDs are necessary.

Meanwhile, single particles and solid residues were collected from the same HW incinerator. Morphologies and elemental compositions of particles in flue gas and indoor air were characterized by transmission electron microscopy-energy dispersive X-ray spectrometry (TEM-EDS). Eight types of single particles were classified, as organic, soot, K-rich, S-rich, Na-rich, Fe-rich, mineral and fly ash particles. The heavy metal partitioning behavior study suggested that Hg, Cd and Pb were mainly enriched in fly ash through evaporation, condensation, and adsorption; while Cr, Cu, Mn, and Ni were mostly remained in the bottom ash due to their low volatilities.

In addition, the study also investigated environmental behaviors of certain characteristic pollutants. Thirty-two soil samples surrounding a cement plant co-processing HW were collected and analyzed for the presence of 16 polycyclic aromatic hydrocarbons (PAHs) and 12 heavy metals. Ten samples were selected for PCDD/Fs analysis. The highest concentration of PCDD/Fs occurred 1200 meters downwind from the cement plant. Levels of $\sum 16$ PAHs ranged from 130.6 to 1134.3 $\mu\text{g}/\text{kg}$ in soil. Source identification analysis suggested that the cement plant was the most likely source of PAH contamination. The concentrations of Cd and Hg were on average two times and six times higher than background values, respectively. Both incremental lifetime cancer risk model (ILCR) for PAHs and potential ecological risk index (RI) for heavy metals indicate potential risks to the population and the environment surrounding the cement plant.

Last, to identify whether waste incineration is a major source for airborne environmentally persistent free radicals (EPFRs), tree leaf samples were collected from 120 sites surrounding four waste incinerators and one urban area. EPFR concentrations on leaves ranged from 7.5×10^{16} to 4.5×10^{19} spins/g. For the 10 N.D. samples, they were all collected from areas inaccessible by vehicles. Although previous work has linked atmospheric EPFRs to waste incineration, the evidence in this study suggests that vehicle emissions, especially from heavy-duty vehicles, are the main sources. According to our estimation, over 90% of the EPFRs deposited on tree leaves might be attributed to automotive exhaust emissions, as a synergistic effect of primary exhausts and degradation of aromatic compounds in road dust.

Keywords: Waste incineration, PCDD/Fs, PAHs, Heavy metals, EPFRs, Primary particles, Formation Mechanism, Spatial distribution, Environmental risk assessment.

Table of Contents

DECLARATION	i
Abstract.....	ii
Table of Contents.....	vi
List of Figures.....	x
List of Tables	xiii
List of Abbreviations	xv
Chapter 1	1
GENERAL INTRODUCTION.....	1
1.1 Industrial-Scale Waste Incineration and the Related Air Pollution Issues	1
1.1.1 Status of Waste Incineration Industry in Domestic and International Scope.....	1
1.1.2 Hazardous Air Pollutants Emitted from Waste Incineration.....	7
1.1.3 Air Pollution Control Measures in Waste Incinerators and Current Deficiencies	11
1.2 Description of the Selected Hazardous Air Pollutants Emitted from Waste Incineration	16
1.2.1 Polychlorinated Dibenzo-p-dioxins and Dibenzofurans (PCDD/Fs)	16
1.2.2 Polycyclic Aromatic Hydrocarbons (PAHs).....	21
1.2.3 Environmental Persistent Free Radicals (EPFRs)	24
1.2.4 Heavy Metals.....	26
1.2.5 Primary Particles.....	28
1.3 Formation Mechanisms of PCDD/Fs, PAHs, EPFRs and Partitioning Characteristics of Heavy Metals During Waste Incineration.....	30
1.3.1 Formation Mechanisms of PCDD/Fs.....	30
1.3.2 Formation Mechanisms of PAHs	34
1.3.3 Formation Mechanisms of EPFRs	35
1.3.4 Partitioning Characteristics of Heavy Metals	37
1.4 Environmental Impacts of Hazardous Air pollutants from Waste Incineration Process.....	38
1.4.1 Research Progress on Transport Patterns of Hazardous Pollutants	39

1.4.2 Research Progress on Environmental and Health Risk Assessment	42
1.5 Objectives of This Study	43
1.6 Framework of This Study	44
Chapter 2	46
FORMATION MECHANISMS OF PCDD/Fs AND THE RELATED EMISSION REDUCTION STRATEGIES: A CASE STUDY IN A HAZARDOUS WASTE INCINERATOR.....	46
2.1 Introduction	46
2.2 Materials and Methods	50
2.2.1 Basic Information on the HW Incinerator and Characteristics of The Raw Feed	50
2.2.2 Sample Collection.....	54
2.2.3 Sample Preparation and Chemical Analysis	55
2.2.4 Quality Assurance and Quality Control	56
2.3 Results and Discussion	57
2.3.1 PCDD/F Concentrations and Congener Compositions at Different Sampling Points	57
2.3.2 PCDD/F Congener Profile.....	65
2.3.3 Formation Mechanisms of PCDD/Fs with the Addition of Precursors	68
2.3.4 Mass Balance and Emission Factors of PCDD/Fs from the Hazardous Waste.....	73
2.4 Conclusions	77
Chapter 3	79
FORMATION PATHWAYS OF PRIMARY PARTICLES AND PARTITIONING OF HEAVY METALS: A CASE STUDY IN A HAZARDOUS WASTE INCINERATOR.....	79
3.1 Introduction	79
3.2 Materials and Methods	82
3.2.1 Sampling Site Description	82
3.2.2 Morphology and Composition Analysis of Single Particles	85
3.2.3 Heavy Metal Content Measurement	86
3.2.4 Stack Gas Emission Level Monitoring	87

3.3 Results and Discussion	88
3.3.1 Morphology, Element Composition, and Mixing State of Single Particles.....	88
3.3.2 Pollution Emission Characteristics and Heavy Metal Mass Balance	100
3.3.3 Heavy Metal Transfer Behaviors in the Incineration Process.....	105
3.4 Conclusions	110
Chapter 4	112
DISTRIBUTION OF PAHs, PCDD/Fs, AND HEAVY METALS IN SOIL SURROUNDING A CEMENT PLANT CO-PROCESSING HAZARDOUS WASTE AND POTENTIAL ENVIRONMENTAL RISKS	112
4.1 Introduction	112
4.2 Materials and Methods	115
4.2.1 Sampling Methodology.....	115
4.2.2 Sample Pretreatment and Chemical Analysis of PCDD/Fs	116
4.2.3 Sample Pretreatment and Chemical Analysis for PAHs	117
4.2.4 Other Chemical Analysis	118
4.2.5 Quality Assurance and Quality Control	119
4.2.6 Statistical Analysis	121
4.2.7 Incremental Lifetime Cancer Risks (ILCRs) for PAHs	121
4.2.8 Pollution Index and Ecological Risk for Heavy Metals	124
4.3 Results and Discussion	125
4.3.1 Concentrations of PCDD/Fs, PAHs and Heavy Metals in Samples	125
4.3.2 Source Identification of PAHs and Heavy Metals in Soil.....	137
4.3.3 Relationship Between PAHs, Heavy Metals, and Soil Properties (pH, TOC, and granulometric composition)	148
4.3.4 Risk Assessment of PAHs and Heavy Metals for Sampling Area	151
4.4 Conclusions	153
Chapter 5	155
SPATIAL DISTRIBUTION AND SOURCE IDENTIFICATION OF EPFRs SURROUNDING WASTE INCINERATORS.....	155
5.1 Introduction	155

5.2 Methods and Materials.....	159
5.2.1 Sampling Sites	159
5.2.2 Sample Collection.....	165
5.2.3 Sample Pretreatment and Analysis	165
5.2.4 Statistical Analysis and EPFR Distribution Mapping.....	172
5.2.5 Evaluation of the Decay Pattern of PM ₂ from Tree Leaves.....	173
5.3 Results and discussion	174
5.3.1 Levels and Distribution Pattern of EPFRs Collected from Tree Leaves	174
5.3.2 Sources and Estimated Contribution Proportions of Airborne EPFRs on Tree Leaves	187
5.3.3 Degradation Characteristics of Airborne EPFRs	196
5.4 Conclusions	198
Chapter 6	201
GENERAL SUMMARY AND CONCLUSIONS	201
6.1 General Summary	201
6.2 Innovations and Scientific Contributions	208
6.3 Limitation of this study.....	209
6.4 Future work.....	210
REFERENCES.....	213
OUTCOME OF THE THESIS WORK.....	253
CURRICULUM VITAE.....	256