

MASTER'S THESIS

Effects of osmotic stress on molecular responses of gill cells from Japanese eels, *Anguilla Japonica*

Ho, Cheuk Hin

Date of Award:
2021

[Link to publication](#)

General rights

Copyright and intellectual property rights for the publications made accessible in HKBU Scholars are retained by the authors and/or other copyright owners. In addition to the restrictions prescribed by the Copyright Ordinance of Hong Kong, all users and readers must also observe the following terms of use:

- Users may download and print one copy of any publication from HKBU Scholars for the purpose of private study or research
- Users cannot further distribute the material or use it for any profit-making activity or commercial gain
- To share publications in HKBU Scholars with others, users are welcome to freely distribute the permanent URL assigned to the publication

Abstract

Japanese eels (*Anguilla japonicas*) are snakelike fishes living in waters in the Asian region. In contrast to most fishes which are stenohalines that can only live in waters with a narrow range of salinity, Japanese eels are classified as euryhalines that can habitat in a broad range of salinity. As the lifecycle of Japanese eels consists of stages across fresh and seawater districts, a well-developed osmoregulation mechanism is needed to balance the intra- and extra-cellular osmolarity of the fishes throughout the seawater acclimation process. While fish gills are one of the organs that separating the ambient water and the inner body fluid of the fish, the fish gills of the Japanese eels have been studied as one of the most crucial organs for osmoregulation purposes. Yet, the osmoregulation and survival strategies of Japanese eels under hyperosmotic stress has not been fully elucidated.

In chapter 2, this study has performed a transcriptome study on the *ex vivo* gill filament model of the Japanese eel to profile the molecular responses after a hypertonic treatment of 4 hours or 8 hours. The experiment is aimed to mimic the gill cells exposed to seawater in the seawater acclimation process of Japanese eels. A profile of differential expressed genes (DEGs) has been revealed that 577 DEGs were commonly upregulated and 711 DEGs were commonly downregulated in both 4- and 8-hours hypertonic treatment. Functional analysis and annotation have been processed with these DEGs, including Ingenuity Canonical Pathways analysis and gene ontology. These analyses have revealed that the cellular homeostasis of the gill cells has been disrupted and cell death responses has been induced by osmotic stress. The

results have raised a concern that the maintenance of cellular viability and a cell death regulation mechanism are needed for the fishes to survive in the early stage of seawater acclimation.

In chapter 3, this chapter demonstrated that gill cells in Japanese eels are susceptible to apoptosis when they are exposed to hyperosmotic treatments in both in vitro gill cell and the ex vivo gill filament model. To maintain the viability of the gills cells, two inhibitors of apoptosis, XIAP, and survivin, were seen to be expressed in gills cells. The expression of XIAP and survivin were upregulated by dexamethasone, which is an agonist mimicking the effect of cortisol on fishes in seawater acclimation. Meanwhile, the expression levels of the apoptosis executor, caspase 3, were downregulated. These data suggested that with the regulation of cortisol express in the fishes, XIAP and survivin are effective apoptosis regulators in the gill cells of Japanese eels.

The study has demonstrated the molecular responses of the gills of Japanese eels exposed to hyperosmotic stress at the transcriptional level and post-translational level by using transcriptome studies and protein study respectively. The study has paved cell death regulation to be another the key field to study in understanding the ability of salinity tolerance in euryhalines.

Tables of Contents

DECLARATION.....	i
Abstract.....	ii
Acknowledgements	iv
Tables of Contents	v
List of Figures	viii
List of Tables.....	ix
List of Abbreviations.....	x
Chapter 1	1
Literature Review	1
1. Eel	1
1.1.) Anguillid eel	1
1.2) Life cycle of <i>Anguilla japonica</i>	1
2. Fish gills.....	5
2.1) Structure	5
2.2) Gill filament.....	6
2.3) Gill epithelium and gill cell.....	7
3. Osmotic and salinity stress	11
4. Cell Death	12
4.1) Programmed cell death and apoptosis	12
4.2) Caspases.....	14
4.3) Inhibitors of apoptosis (IAPs)	16
5. Objectives	18
Chapter 2	19
Transcriptomic Analysis of <i>Ex Vivo</i> Gill Filaments under Hyperosmotic Challenge ...	19
2.1 Introduction	19
Materials and Methods	21
i) Animal	21
ii) <i>Ex vivo</i> gill filament culture	21
iii) Osmotic stress induction treatment.....	21

iv) RNA isolation and quantification.....	23
v) Transcriptome sequencing	23
vi) DEGs Functional annotation.....	23
vii) Real-time PCR analysis	24
2.3 Results	25
i) Transcriptome revealed DEGs in <i>ex vivo</i> gill filament culture upon hyperosmotic stress	25
ii) Functional analysis.....	26
2.4 Discussion	33
i) Acute osmotic stress induces a broad set of dysregulated genes.....	33
ii) Acute osmotic stress did not induce the expression of ion transporters.....	33
iii) Enhanced <i>tgfb3</i> expression evidenced SW acclimation process in the <i>ex vivo</i> gill model	34
iv) Osmotic stress showed disruptions to cellular homeostasis, and trigger cell death responses in Japanese eel.....	34
v) Enriched molecular function lied clues for the molecular mechanism that interrupted translation process or protein synthesis	35
2.5 Summary.....	37
Chapter 3	38
The Anti-apoptotic Role of Dexamethasone on Gill cells upon Osmotic Challenge ..	38
3.1 Introduction	38
Materials and Methods	40
i) Animal	40
ii) <i>In vitro</i> primary gill cell culture	40
iii) <i>Ex vivo</i> gill filament culture	41
iv) Hormonal treatment and osmotic stress induction treatment	41
vi) Protein extraction and western blot	42
3.3 Result	43

<i>In vitro</i> gill cell model.....	43
i) Hyperosmotic stress reduced the viability of gill epithelial cells	43
ii) Hyperosmolarity triggered the activation of apoptotic executor caspase 3	43
iii) The SW acclimation hormone cortisol reduced the level of cleaved caspase-3 expressions in SW acclimating gill cells.....	44
<i>Ex vivo</i> gill filament model	50
iv) <i>Ex vivo</i> gill filament culture model showed pro-caspase-3 cleavage and the expression of inhibitors of apoptosis induced by osmotic treatment.....	50
v) Dexamethasone reduced caspase-3 cleavage and induced the expression of XIAP and survivin in <i>ex vivo</i> gill filament <i>model</i> under hypertonic treatment	50
3.4 Discussion	53
i) Hypertonicity activates pro-caspase-3 cleavage and apoptosis.....	53
ii) Dexamethasone diminished the threat of apoptosis in gill cells under hyperosmotic treatment.....	54
iii) Expression of inhibitors of apoptosis were induced by osmotic pressure	55
iv) Dexamethasone prompts the expression of both XIAP and survivin in the <i>ex vivo</i> gill filament culture under hypertonic treatment.....	56
3.5 Summary.....	58
Appendix.....	59
Bibliography	230
CURRICULUM VITAE.....	248