

MASTER'S THESIS

Measuring the binding between estrogen receptor alpha and potential endocrine disruptors by fluorescence polarization and total internal reflection fluorescence

Yiu, Kwok Wing

Date of Award:
2013

[Link to publication](#)

General rights

Copyright and intellectual property rights for the publications made accessible in HKBU Scholars are retained by the authors and/or other copyright owners. In addition to the restrictions prescribed by the Copyright Ordinance of Hong Kong, all users and readers must also observe the following terms of use:

- Users may download and print one copy of any publication from HKBU Scholars for the purpose of private study or research
- Users cannot further distribute the material or use it for any profit-making activity or commercial gain
- To share publications in HKBU Scholars with others, users are welcome to freely distribute the permanent URL assigned to the publication

**Measuring the Binding between Estrogen Receptor Alpha and
Potential Endocrine Disruptors by Fluorescence Polarization
and Total Internal Reflection Fluorescence**

YIU Kwok Wing

**A thesis submitted in partial fulfillment of the requirements
for the degree of
Master of Philosophy**

Principal supervisor: Prof CHEUNG Nai Ho

Hong Kong Baptist University

June 2013

Abstract

Endocrine systems are found in all mammals and many other types of living organisms. It consists of glands that secrete hormones and receptors that detect and react to the hormones and regulates all biological processes in the body. Endocrine disruptors are hormone-like chemicals that can be found in the environment around us. Once taken in, they may cause hormonal effects. This thesis reports the measurement of binding strength and binding kinetics between human estrogen receptor and different kinds of endocrine disruptors by means of fluorescence polarization and total internal reflection fluorescence. Our results show that the equilibrium dissociation constants for different endocrine disruptors ranged from $\sim 1\text{nM}$ to $\sim 100\mu\text{M}$ and their off-rates ranged from $\sim 1.5 \times 10^{-3}$ to $\sim 30 \times 10^{-3} \text{ s}^{-1}$. We also found that mixture of endocrine disruptors (PBA, PFOS, PFNA and PFOA) did not show synergistic binding with the endocrine disruptor.

Table of Contents

Declaration	i
Abstract	ii
Acknowledgments	iii
Table of Contents	iv
List of Tables	viii
List of Figures	ix
List of Abbreviations	xiv
Chapter 1 Introduction	1
Chapter 2 Background	4
2.1 Receptor ligand interaction	4
2.1.1 Nuclear Receptor superfamily	4
2.1.2 Estrogen receptor structure and function	5
2.1.3 Breast cancer and drug discovery	10
2.1.4 Endocrine disruptors and contaminant screening	12
2.1.5 Peroxisome proliferator-activated receptors	13
2.1.6 Ginsenosides	14
2.2 <i>In-vitro</i> binding assays	16
2.2.1 Mass difference assays	16
2.2.2 Surface plasmon resonance (SPR)	16
2.2.3 Fluorescence assays	18

2.2.4	Total internal reflection fluorescence (TIRF)	19
2.2.5	Total internal reflection with fluorescence correlation spectroscopy (TIR-FCS)	20
2.2.6	Total internal reflection with fluorescence photobleaching recovery (TIR-FPR)	21
2.2.7	Total internal reflection with reflectance interferometry (TIR-RIf)	22
2.2.8	Fluorescence polarization (FP)	23
2.2.9	Free probes versus tethered probes	24
Chapter 3 Methods		26
3.1	FP-TIRF setup	26
3.2	Setup optimization and calibration	27
3.3	Flow cell fabrication and fluidics	29
3.3.1	Flow cell fabrication	30
3.3.2	Fluidics	31
3.4	Channel surface preparation	32
3.4.1	Coating with PEG	32
3.4.2	Immobilization of primary and secondary antibody	33
3.4.3	Receptor preparation and tethering	34
3.5	Ligand preparation	36
3.6	Program for real-time data capture and display	36
3.7	Data runs	38

3.7.1	Kinetics data capture for surface tethered receptors	38
3.7.2	Kinetics data capture for free receptors and ligands in solution ..	39
3.7.3	Screening ginsenosides for binding with nuclear receptors	40
3.8	Data analysis – From fluorescence anisotropy to bound fraction	41
3.9	Kinetics modeling	43
3.10	Rapid mixing assumption	44
Chapter 4 Results and discussion		46
4.1	Characterization of the FP-TIRF setup	47
4.1.1	Correcting for // and \perp arm imbalance	47
4.1.2	Monitoring [F]	48
4.1.3	Consistency of FP and TIRF	48
4.2	Characterization of each tether link	51
4.2.1	Strength and specificity of tether	52
4.2.2	Tethering preserved receptor function	57
4.2.3	Tethering preserved the random orientation of μ	59
4.3	Binding kinetics of the fluorescent ligand with ER α	62
4.4	Justify rapid mixing assumption	68
4.5	Binding kinetics of nonfluorescent ligands with ER α	72
4.6	Positive and negative controls	78
4.7	Screening ginsenosides for ER α binding	80
4.8	Screening ginsenosides for PPAR γ binding	82
4.9	Binding affinity and chemical structure	85

Chapter 5 Conclusion	91
List of References	94
Curriculum Vitae	100