

DOCTORAL THESIS

Use of food waste feeds for culturing low trophic level fish (grass carp, bighead carp and mud carp): persistent toxic substances

Cheng, Zhang

Date of Award:
2014

[Link to publication](#)

General rights

Copyright and intellectual property rights for the publications made accessible in HKBU Scholars are retained by the authors and/or other copyright owners. In addition to the restrictions prescribed by the Copyright Ordinance of Hong Kong, all users and readers must also observe the following terms of use:

- Users may download and print one copy of any publication from HKBU Scholars for the purpose of private study or research
- Users cannot further distribute the material or use it for any profit-making activity or commercial gain
- To share publications in HKBU Scholars with others, users are welcome to freely distribute the permanent URL assigned to the publication

ABSTRACT

This study aimed at using different types of food wastes as major sources of protein to replace the fish meal used in fish feeds to produce quality fish. The major objectives were to (1) investigate the variations of metalloids/metals, polycyclic aromatic hydrocarbons (PAHs) and organochlorine pesticides (OCPs) in the fish ponds (pond mud and water), and food wastes used as fish feeds; (2) analyze bioaccumulation and biomagnification of pollutants in the food chains; and (3) evaluate the potential health risks of exposure (to these pollutants) via dietary intake of fish fed with food waste feeds.

The traditional fish farming model was used to culture low trophic level fish: a filter feeder (bighead, *Aristichthys nobilis*), a herbivore (grass carp, *Ctenopharyngodon idellus*) and a bottom feeder (mud carp, *Cirrhina molitorella*), which are more environmental friendly as they can utilize more solar energy. Furthermore, low-trophic level fish are less susceptible to the accumulation of toxic chemicals. Two types of food wastes (mainly containing cereal (Food Waste A) and meat waste meal (Food Waste B)) were used as the major source of protein to replace the fish meal in fish feed to culture fish.

The concentrations of metalloids (arsenic (As)), metals (mercury (Hg), cadmium (Cd), chromium (Cr), lead (Pb), zinc (Zn), copper (Cu) and nickel (Ni)) in water, suspended particulate matter and sediment of the 3 experimental fish ponds located in Sha Tau Kok Organic Farm were monitored (bi-monthly during the first half year and tri-monthly during the second half year) and the results were similar to or lower than those in the commercial fish ponds around the Pearl River Delta (PRD) region. Results of the health risk assessments indicated that human consumption of grass carp (a herbivore) which fed

food waste feed pellets would be safer than other fish species (mud carp, bighead carp and largemouth bass). There were no or lower magnifications, and low concentrations of metalloids/metals contained in the ponds indicated that the practice of traditional pond management by draining pond water regularly can provide a better fish pond habitat for birds and other wildlife. Furthermore, the use of food waste instead of fish meal (mainly consisted of contaminated trash fish) further reduced Hg accumulation in the cultured fish.

During October 2011 - December 2012, the concentrations of PAHs and OCPs in three experimental fish ponds were monitored (bi-monthly during the first half year and tri-monthly during the second half year). The results were similar to or lower than those obtained in commercial fish ponds around the PRD region. The mean concentrations of Σ PAHs and Σ OCPs in sediment and fish collected from the experimental fish ponds during the 2nd half year (May 2012 to December 2012) were significantly higher ($p < 0.05$) than those in the 1st half year (October 2011 to April, 2012). Σ PAHs and Σ DDTs in the two species of fish (grass carp and bighead carp) were significantly increased ($p < 0.05$) with time, and PAHs and DDTs in grass carp and bighead carp fed with commercial fish feed pellets (control group) were significantly higher ($p < 0.05$) than the fish fed with food waste pellets (Food Waste A and Food Waste B). Fruit, vegetables, bone meal and meat products were the major sources of PAHs and OCPs contamination for producing Food Waste A and Food Waste B. No significant increases in PAHs and DDTs concentrations with trophic levels were observed in the experimental ponds, showing that PAHs were not biomagnified in the omnivorous food chains (plankton, grass carp, bighead carp and mud carp). DDTs were lower magnifications than those predatory food chains (plankton, trash fish, and largemouth bass) in farmed ponds. There was a very low cancer risk for

PAHs and DDTs exerted on humans via consumption of bighead carp, grass carp and mud carp (fed with food waste and commercial pellets). Furthermore, the use of food waste instead of fish meal (mainly consisted of contaminated trash fish) further reduced accumulation of PAHs and DDTs in the cultured fish.

The present results revealed that recycling of food waste for cultivating low trophic level fish (mainly bighead carp and grass carp) is feasible, which will also ease the disposal pressure of the large volume of food waste, a common problem encountered in densely populated cities such as Hong Kong.

ACKNOWLEDGEMENTS

I want to express my sincere appreciation and gratitude to my principal supervisor, Prof. M.H. Wong, for providing an opportunity for me to pursue a doctorate degree and for his professional guidance, encouragement, constructive suggestions and continuous support during my study period. Being able to study at Hong Kong Baptist University under guidance of Prof. M.H. Wong is a proud memory of my life journey. I would also like to acknowledge my co-supervisor, Prof. Chris K.C. Wong, for his enthusiastic guidance and support.

Special thanks are given to Prof. X. P. Nie at the Jinan University for his enthusiastic guidance and support during my time in Hong Kong Baptist University. I also thank Prof. K.B. Li and Prof. K.C. Chen at Pearl River Fisheries Research Institute for their assistance in field work.

I am grateful to Ms. S. Fung for her help in improving my manuscripts, and Dr. S.C. Wu for providing me valuable suggestions on my experiments. I would like to thank my co-workers Mr. W.Y. Mo, Mr. W.M. Choi and Mr C.L. Lam, and also wish to thank Dr. J.L. Hu, Dr. Y Kang, Dr. H.S. Wang, Dr. P. Liang, Dr. X.W. Chen, Dr. Y.G. Zhao, Dr. Y.L. Zhang, Dr. M.J. Huang, Dr. J.L. Liu and Dr. Y.B. Man for their assistance and useful comments. Special thanks are also given to Ms. H.Y. Ng, Dr. F.Y Wu, Dr. W. Wang, Ms. J.L. Miao and Mr. L.Y. Zhang for the joy they have brought me during my time in Hong Kong.

Lastly, I express my sincere gratitude to my parents for their love and moral support throughout the period of my study.

TABLE OF CONTENT

DECLARATION	i
ABSTRACT.....	ii
ACKNOWLEDGEMENTS	v
TABLE OF CONTENT	vi
LIST OF TABLES.....	xii
LIST OF FIGURES	xvi
ABBREVIATIONS	xx
CHPAPTER 1 GENERAL INTRODUCTION	1
1.1 Food Wastes	1
1.2 Food Safety Issues and Pollutants in Fish Products in Hong Kong	6
1.3 Hong Kong Inland Fisheries	9
1.4 Fish Culture	9
1.5 Integrated Fish Farming System	11
1.6 Fish Pond Habitat Conservation.....	12
1.7 Objectives of the Present Study	13
1.7 Framework	14
1.8 Significance of the Present Study.....	16
CHAPTER 2 TROPHIC RELATIONSHIPS AND HEALTH RISK ASSESSMENTS OF HEAVY METALS IN THE AQUACULTURE POND ECOSYSTEM OF PEARL RIVER DELTA, CHINA.....	17
2.1 Introduction.....	17

2.2 Materials and Methods	19
2.2.1 Sampling.....	19
2.2.2 Analytical Methods.....	21
2.2.3 Quality Control.....	23
2.2.4 Assessment of Sediment Quality	24
2.2.5 Biota-Sediment Accumulation Factor (BSAF).....	26
2.2.6 Stable Isotope Analysis	26
2.2.7 Trophic Level and Biomagnification Calculations.....	26
2.2.8 Risk Assessment.....	27
2.2.9 Data Analyses	29
2.3 Results and Discussion.....	29
2.3.1 Sediment	29
2.3.2 Fish	36
2.4 Conclusions	46
 CHAPTER 3 VARIATIONS OF METALLOID/METALS IN FRESHWATER FISH PONDS, USING FOOD WASTES AS FISH FEEDS	
3.1 Introduction	48
3.2 Materials and Methods	51
3.2.1 Experimental Design	51
3.2.2. Sampling.....	53
3.2.3. Analytical Methods.....	56
3.2.4. Quality Control	57
3.2.5 Statistical Analyses.....	58

3.3 Results and Discussion.....	58
3.3.1 Fish Feeds.....	58
3.3.2 Environmental Samples.....	60
3.4 Conclusions.....	67
 CHAPTER 4 BIOACCUMULATION AND HEALTH RISK ASSESSMENT OF METALLOID/METALS IN FRESHWATER FISH, USING FOOD WASTES AS FISH FEEDS	
4.1 Introduction.....	68
4.2 Materials and Methods.....	70
4.2.1 Experimental Design.....	70
4.2.2 Sampling.....	70
4.2.3 Analytical Methods.....	71
4.2.4 Quality Control.....	71
4.2.5 Bioaccessibility of Metalloid/Metals.....	72
4.2.6 Bioaccumulation Factor.....	73
4.2.7 Stable Isotope Analysis.....	73
4.2.8 Trophic Level and Biomagnification Calculations.....	73
4.2.9 Health risk Assessment.....	74
4.2.10 Statistical Analyses.....	75
4.3 Results and Discussion.....	75
4.3.1 Metalloid/Metal Concentrations.....	75
4.3.2 Bioaccessibility and Health Risk assessment.....	78
4.3.3 Bioaccumulation and Biomagnification.....	85

4.4 Conclusions	90
CHAPTER 5 VARIATIONS OF PAHs AND OCPs IN FRESHWATER FISH PONDS, USING FOOD WASTES AS FISH FEEDS	91
5.1 Introduction	91
5.2 Materials and Methods	94
5.2.1 Experiment Design	94
5.2.2 Sampling of Water, SPM and Sediment	94
5.2.3 Sample Extraction.....	94
5.2.4 Instrumental Analyses	95
5.2.5 QA/QC.....	96
5.2.6 Statistical Analyses.....	96
5.3 Results and Discussion.....	99
5.3.1 Fish Feeds.....	99
5.3.2 PAHs in Environmental Samples	100
5.3.3 OCPs in Environmental Samples	108
5.4 Conclusions	113
CHAPTER 6 HEALTH RISK ASSESSMENTS OF PAHs AND OCPs IN FRESHWATER FISH, USING FOOD WASTE AS FISH FEEDS	115
6.1 Introduction	115
6.2.2 Sampling.....	118
6.2.3 Chemical Analyses	118
6.2.4 Digestible Fraction of PAHs and OCPs	118
6.2.5 Calculation.....	119

6.2.6. Bioaccumulation and Biomagnification	121
6.2.7 Statistical Analyses.....	121
6.3 Results and Discussion.....	121
6.3.1 OCPs and PAHs Concentrations	121
6.3.2 Bioaccessibilities and Health Risk Assessment.....	130
6.3.3 Bioaccumulation and Biomagnification	136
6.4 Conclusions	142
CHAPTER 7 GENERAL DISCUSSION AND CONCLUSIONS.....	143
7.1 Introduction.....	143
7.2 Growth Performance, Feed Nutrient Utilization, and Levels of Environmental Pollutants in Fish Feeds.....	145
7.3 The Levels of Environmental Pollutants in Water, SPM, and Sediments Collected from Experimental Ponds and Other Fish Ponds (Control Sites).....	146
7.4 The Levels of Environmental Pollutants in Fish Collected from Experimental Ponds and Other Fish Ponds (Control Sites).....	147
7.5 Bioaccessibilities and Health Risk Assessments of Consumption The Fish Fed with Food Waste Feeds	149
7.7 Bioaccumulation and Biomagnification of Environmental Pollutants in Fresh Water Fish, Using Food Waste as Fish Feed	154
7.8 General Conclusions	157
7.9 Limitations of the Present Study	160
7.10 Future Work	161
References.....	163

Publications.....	188
Curriculum Vitae	192