

## DOCTORAL THESIS

### Characterization of in-situ $\text{Ca}^{2+}$ -sensing mechanisms and stanniocalcin-1 target cells in gills of Japanese eels

Gu, Jie

*Date of Award:*  
2014

[Link to publication](#)

#### General rights

Copyright and intellectual property rights for the publications made accessible in HKBU Scholars are retained by the authors and/or other copyright owners. In addition to the restrictions prescribed by the Copyright Ordinance of Hong Kong, all users and readers must also observe the following terms of use:

- Users may download and print one copy of any publication from HKBU Scholars for the purpose of private study or research
- Users cannot further distribute the material or use it for any profit-making activity or commercial gain
- To share publications in HKBU Scholars with others, users are welcome to freely distribute the permanent URL assigned to the publication

## Abstract

Calcium ion has diverse beneficial roles in living organisms. Failure in  $\text{Ca}^{2+}$  homeostasis affects a variety of molecular and cellular processes, ultimately leading to many pathological consequences. In mammals, body  $\text{Ca}^{2+}$  homeostasis is maintained by the coordinated calcium (re)absorption that occurs in the small intestines, kidneys and bones, and is under tight hormonal control. In fish, two special organs, Corpuscles of Stannius (CS) glands and gills form a regulatory circuit to detect and regulate blood  $\text{Ca}^{2+}$  homeostasis. However, the underlying molecular mechanism in the regulation of gill  $\text{Ca}^{2+}$  uptake has not been fully examined. Moreover, some putative biological active substances in CS glands have not been identified. To address these research questions, a euryhaline fish, Japanese eel (*Anguilla japonica*) was used as an animal model for the study.

Fish gill is equipped with epithelial calcium channel (ECaCl) as gatekeeper of  $\text{Ca}^{2+}$  entry, and membrane  $\text{Ca}^{2+}$ -ATPase (PMCA) for  $\text{Ca}^{2+}$  efflux. To test if branchial ECaCl and PMCA responded to change in water  $\text{Ca}^{2+}$  level, we investigated the changes in fish adapted in artificial freshwater (AFW),  $\text{Ca}^{2+}$ -deficient AFW (D-AFW) or high  $\text{Ca}^{2+}$ -AFW (H-AFW). Our data illustrated both short-term and long-term effects on modulations of the transporters. The changes correlated with expression levels of stanniocalcin-1 (STC-1) in CS glands. This part of study supports the regulatory circuit between gills and the glands. In primary cell culture of Japanese eel gill cells,  $\text{Ca}^{2+}$  sensing was shown to be mediated by  $\text{Ca}^{2+}$  sensing receptor (CaSR) coupled to phospholipase C (PLC)-extracellular signal-regulated kinase (ERK) and PLC-inositol triphosphate ( $\text{IP}_3$ )- $\text{Ca}^{2+}$ /calmodulin-dependent protein kinase-II (CaMK-II) pathways. And CaSR-STC-1/cyclo-oxygenase-2 (COX-2) mediated protective pathway in gill cells that exerts a possible protective mechanism against an increase in intracellular  $\text{Ca}^{2+}$  levels associated with transepithelial  $\text{Ca}^{2+}$  transport. Apparently, the protective effects against  $\text{Ca}^{2+}$ -mediated cytotoxicity of gill cell were mediated by STC-1 binding on gill cells that led to elevations of cytosolic cAMP. In a follow-up experiment of using  $\text{Ca}^{2+}$ -imaging system in a model of thapsigargin (TG)-induced elevation of cytosolic  $\text{Ca}^{2+}$ , a hypocalcemic action of STC-1 was demonstrated and was found to be mediated by cAMP and COX-2 pathway. To further determine the gene expressed in CS gland responsive to changes in water salinity, the first transcriptome database of CS glands from fish adapted in freshwater or seawater condition. A *de novo* assembly of RNA sequencing data generated 11747 unigenes and revealed 475 genes that were differentially expressed. Three functional clusters: (1)  $\text{Ca}^{2+}$ -metabolism, (2) blood pressure and (3) ion-osmoregulation were revealed. Gene targets, in addition to STC-1 in related to the regulation of calcium metabolism and blood pressure, like calcitonin, atrial natriuretic peptide-converting enzyme and endothelin-converting enzyme 1 were identified. Taken together this thesis described a comprehensive study on the functional circuit between gills and CS glands to decipher the regulation and functions of transporters and hormones in calcium metabolism in fish.

## **Acknowledgments**

I would like to express my sincere gratitude to my principal supervisor, Prof. Chris KC Wong, for his guidance, support, and many helpful comments throughout the study. I am also grateful to Prof. NK MAK, my co-supervisor, for his advice and support. Also, special thanks to Prof. Ricky NS Wong, Prof. J Zhang for their precious advice and generous support.

Besides, I would also like to thank Dr. Bonnie Yeung, Ms. Eman. Lee, Ms. Louise Ng, Ms. Olivia Chau, Mr. KW Chan, Mr. WS Chung, Mr. KK Ma, Mr. WK Ip, and other technicians in the Biology Department (Hong Kong Baptist University), for giving their help and their precious suggestion. And I also appreciate Dr. GG Zhao, Dr. X Wei, Dr. Jamie Wan, Dr. Alice Law, Dr. William Tse, Mr. Oscar Chow and other lab colleagues, for their technical assistances and advices.

Last but not least, I would like to thank my family, colleagues and friends for their encouragement and understanding.

## Table of Contents

	Page no.
<b>Declaration</b>	<b>i</b>
<b>Abstract</b>	<b>ii</b>
<b>Acknowledgments</b>	<b>iii</b>
<b>Table of Contents</b>	<b>iv</b>
<b>List of Figures</b>	<b>viii</b>
<b>List of Tables</b>	<b>x</b>
<b>List of Abbreviations</b>	<b>xi</b>
<b>Chapter 1 Literature Review</b>	<b>1</b>
<b>1. General Information of Japanese Eel</b>	<b>1</b>
<b>(I) Classification</b>	<b>1</b>
<b>(II) Life Cycle of Japanese Eel</b>	<b>1</b>
<b>2. Anatomical Structure Fish Gills and the Major Types of Epithelial Gill Cells</b>	<b>4</b>
<b>(I) Fish Gill</b>	<b>4</b>
<b>(II) Epithelial Gill Cells</b>	<b>5</b>
<b>(i) Pavement Cells (PVCs)</b>	<b>6</b>
<b>(ii) Chloride Cells (CCs)</b>	<b>7</b>
<b>(iii) Accessory Cells (ACs)</b>	<b>7</b>
<b>3. Corpuscles of Staninus (CS)</b>	<b>8</b>
<b>4. Ca<sup>2+</sup> Homeostasis</b>	<b>9</b>
<b>(I) Ca<sup>2+</sup> Homeostasis in Mammals</b>	<b>9</b>
<b>(II) Ca<sup>2+</sup> Homeostasis in Fish</b>	<b>10</b>
<b>(i) Stanniocalcin-1 (STC-1)</b>	<b>11</b>
<b>(ii) Calcitonin</b>	<b>12</b>
<b>(III) Ca<sup>2+</sup> Transporters</b>	<b>13</b>

(i) ECaCl	13
(ii) PMCA and NCX	14
(IV) Calcium Sensing Receptor (CaSR) Mediated Signaling Pathways	15
5. Objectives	18
Chapter 2 Molecular Responses of Gill Cells of Japanese Eels to Change in Ambient Ca <sup>2+</sup> Levels	19
2.1 Introduction	20
2.2 Materials and Methods	22
2.3 Results	26
2.3.1 Cloning and Expression of Eel <i>Ecacl</i> and <i>Pmca</i>	26
2.3.2 Change of Serum Osmolarity, and Ions Concentrations in the Fish Acclimated in H-AFW and D-AFW	32
2.3.3 Gene Expressions of Ca <sup>2+</sup> Transporters ( <i>ecacl</i> , <i>pmca</i> ) and <i>stc-1</i> in Gill Scrapings, Gill PVCs and CCs in the Fish Acclimated in H-AFW and D-AFW	34
2.3.4 Gene Expressions of Ca <sup>2+</sup> -Related Proteins in CS gland in the Fish Acclimated in H-AFW and D-AFW	36
2.3.5 Cell Density of CCs in Gills	38
2.4 Discussion	40
2.4.1 In the Acclimation at H-AFW or D-AFW, the Changes of Serum Osmolarity and Ion Concentrations were Accompanied with Modulations in mRNA Expression Levels of <i>ecacl</i> in Gill Cells	40
2.4.2 Changes in <i>stc-1</i> Expression in CS glands of Fish Acclimated in H-AFW or D-AFW	41

<b>Chapter 3</b>	<b>Activation of Gill Ca<sup>2+</sup>-Sensing Receptor as a Protective Pathway to Reduce Ca<sup>2+</sup>-Induced Cytotoxicity</b>	<b>43</b>
3.1	Introduction	44
3.2	Materials and Methods	46
3.3	Results	50
3.2.1	Detection of Gill CaSR and the Effects of Ca <sup>2+</sup> and/or Calcimimetic Treatment on the Activation of PC-PLC-ERK Signaling Pathway in Isolated Gill Cells	50
3.2.2	Effects of Ca <sup>2+</sup> and/or Calcimimetic Treatment on the Activation of CaMK-II in Isolated gill cells	54
3.2.3	Effects of Ca <sup>2+</sup> and/or Calcimimetic Treatment on stc-1 and cox-2 mRNA and Protein Expression in Isolated Gill Cells	56
3.2.4	Effects of CaSR Activation, stc-1 Overexpression, and cox-2 Inhibition (NS-398) on Thapsigargin-induced Cell Death	58
3.4	Discussion	63
<b>Chapter 4</b>	<b>Characterization of STC-1-binding, signaling and function in gill cells</b>	<b>67</b>
4.1	Introduction	68
4.2	Materials and Methods	70
4.3	Results	76
4.3.1	Effects of CS Extracts on the Production of cAMP in Gill Cells	76
4.3.2	The Binding of the STC-1 Recombinant Protein (STC-1-V5) on Paraffin-embedded Gill Tissues of FW and High-Ca <sup>2+</sup> FW Adapted Fish	78

4.3.3	<b>The Binding of the STC-1 Recombinant Protein (STC-1-AP) on Subcellular Fractions of Gill Preparations</b>	80
4.3.4	<b>Effects of STC-1-V5 on Intracellular Ca<sup>2+</sup> Level, Expression of Epithelial Ca<sup>2+</sup> Channel (ECACL) and Cyclo-oxygenase-2 (COX-2)</b>	82
4.4	<b>Discussion</b>	87
<b>Chapter5</b>	<b>Transcriptome Analysis of CS Gland of Japanese eel (<i>Anguilla japonica</i>) in Response to Changed Water Environment</b>	<b>90</b>
5.1	<b>Introduction</b>	91
5.2	<b>Materials and Methods</b>	93
5.3	<b>Results</b>	96
5.3.1	<b>Illumina Draft Reads and Sequence Assembly</b>	96
5.3.2	<b>Annotation of Unigenes</b>	100
5.3.3	<b>GO Enrichment Analysis and Cluster Classification</b>	102
5.3.4	<b>Differentially Expressed Genes</b>	106
5.4	<b>Discussion</b>	110
	<b>Appendix</b>	114
	<b>References</b>	146
	<b>Curriculum Vitae</b>	166