

MASTER'S THESIS

Differentiation inducing effect of isoflavonoids on neuroblastoma cells

Tai, Lai Shan

Date of Award:
1999

[Link to publication](#)

General rights

Copyright and intellectual property rights for the publications made accessible in HKBU Scholars are retained by the authors and/or other copyright owners. In addition to the restrictions prescribed by the Copyright Ordinance of Hong Kong, all users and readers must also observe the following terms of use:

- Users may download and print one copy of any publication from HKBU Scholars for the purpose of private study or research
- Users cannot further distribute the material or use it for any profit-making activity or commercial gain
- To share publications in HKBU Scholars with others, users are welcome to freely distribute the permanent URL assigned to the publication

**Differentiation Inducing Effect of Isoflavonoids on
Neuroblastoma Cells**

TAI Lai Shan

A thesis submitted in fulfillment of the requirements
for the degree of
Master of Philosophy

October 1999

Hong Kong Baptist University

Abstract

Neuroblastoma, an embryonal tumor of the nervous system, is one of the major solid tumors that usually occur in children. It is a group of primitive and clinically aggressive neoplasm. In my present study, murine neuroblastoma Neuro-2A was used as a cell model to study the anti-tumor activities of two selected isoflavonoids, namely biochanin A and genistein. These two isoflavonoids have been previously reported to mediate anti-tumor activities in mammalian cancer cells.

Biochanin A and genistein not only can inhibit the growth of Neuro-2A, but also induce morphological differentiation of this tumor cell. Morphological differentiation was observed as early as 1 day after isoflavonoids treatment. Neuro-2A is a heterogenous population of cells. Two subclones, namely BU-1 and BU-2, were isolated for the subsequent studies. The cytotoxic and growth inhibitory effects of both biochanin A and genistein on Neuro-2A, BU-1 and BU-2 cells were similar. The degree of differentiation in isoflavonoids treated BU-1 cell was higher than Neuro-2A and BU-2 cells. Morphological features of biochanin A- and genistein-treated BU-1 cells are different. Biochanin A treated BU-1 cells are generally elongated in shape, whereas genistein treated BU-1 cells produce extensive neurite network. Both biochanin A and genistein induce morphological differentiation in a dose- and time dependent manner. Results from limiting dilution analysis reveal that isoflavonoids treated BU-1 cells undergo terminal differentiation at 6 days after treatment.

Axonal (NF-H and tau) and dendritic (MAP-2) markers were employed to characterize the nature of induced neuritic outgrowth in differentiated BU-1 cells. Laser scan confocal microscopy was used to study the distribution of these neuronal markers. A mixed phenotype of axon- and dendrite-like processes was observed 3 days after biochanin A and genistein treatment. In terminally differentiated BU-1 cells (after 6 days of incubation), axonal property was observed in the neurites.

Neurotrophic cytokines are important in controlling neuronal growth and differentiation in the nervous system. In this study, the roles of some selected neurotrophic cytokines (IL-6, CNTF and LIF) upon isoflavonoids induced differentiation of neuroblastoma BU-1 cells was examined. The mRNA expression of neurotrophic cytokine genes was examined using Reverse-transcription Polymerase chain reaction (RT-PCR). The level of expression was semi-quantified by cycle titration and dot blot hybridization. The expression of CNTF was not significantly altered in isoflavonoids-treated BU-1 cells. However, the expression profile of IL-6 and LIF was different in biochanin A- and genistein-treated BU-1 cells. Our results suggest that IL-6 and LIF possibly interact with other neurotrophic cytokines in controlling neuronal differentiation of BU-1 cells.

In summary, biochanin A and genistein may act as anti-cancer compounds upon neuroblastoma cells. The induced neuronal differentiation and also the nature of isoflavonoids driven neuritogenesis may be under the control of neurotrophic cytokines. The role of cytokines on isoflavonoids induced neuronal maturation remains to be investigated. The biological effects of isoflavonoids on the development of neurons should also be studied. Hopefully, the underlying mechanism of cellular differentiation of neuroblastoma can be elucidated in future.

Contents

Declaration	i
Abstract	ii
Acknowledgements	iv
Contents	v
List of Tables	xiii
List of Figures	xiv
Abbreviations	xvi
Chapter One Introduction and Literature Review	1
1.1 Neuroblastoma	1
1.1.1 Pathology and clinical features of neuroblastoma	1
1.1.1.1 Origin of neuroblastoma	1
1.1.1.2 Histology of neuroblastoma	2
1.1.1.3 Clinical features of neuroblastoma	3
1.1.2 Diagnosis and investigations of neuroblastoma	4
1.1.2.1 Tumor markers	4
1.1.2.2 Serum markers	5
1.1.2.3 Tumor cell DNA content	5
1.1.2.4 N-myc oncogene activation	6
1.1.2.5 Gene expression	6
1.1.3 Clinical staging of neuroblastoma	7
Table 1. Staging system developed by Evans and his colleagues	8
Table 2. International Neuroblastoma Staging System	9
1.1.4 Treatment of neuroblastoma	10
Table 3. Guidelines for treatment of neuroblastoma	11
1.1.5 Murine neuroblastoma, Neuro-2A	11

1.2 Flavonoids	13
1.2.1 Classification and occurrence of flavonoids	13
Figure 1. Basic structure of flavonoid	15
Table 4. Classification and terminology of major flavonoids	16
1.2.2 Biological and pharmacological activities of flavonoids	17
Table 5. Functions of major flavonoids subclasses	17
1.2.3 Isoflavonoids	19
1.2.3.1 Biochanin A	20
1.2.3.2 Genistein	21
Figure 2A. Basic structure of isoflavone	22
Figure 2B. Structure of biochanin A	22
Figure 2C. Structure of genistein	22
1.3 Architecture and Cytoskeletons of neurons	23
1.3.1 Structural proteins of cytoskeleton in neurons	24
1.3.1.1 Microtubules : An important determinant of cell architecture	24
Table 6. Major microtubule proteins in mammalian brain	28
1.3.1.2 Microfilaments : Involved in intracellular transport and cell movement	29
Table 7. Major proteins of the microfilament cytoskeleton in brain	31
1.3.1.3 Intermediate filaments : Prominent constituents of nervous system	32
Table 8. Subclasses of intermediate filament proteins and their distribution	33
1.4 Cytokines	34
1.4.1 General features	34
Table 9. Utilization of Jak/STAT members by some soluble mediators	36
1.4.2 Cytokines in neuronal cell growth and differentiation	37
Table 10.	38
1.4.2.1 CNTF	39
1.4.2.2 LIF	40
1.4.2.3 IL-6	41
1.5 Aims of study	42

Chapter Two Materials and Methods	44
2.1 Materials	44
2.1.1 Cell line	44
2.2 Methods	
2.2.1 Biological effects of biochanin A and genistein on murine Neuro-2A and its subclone, BU-1 and BU-2	45
2.2.2.1 Isolation of BU-1 and BU-2 subclones	45
2.2.2.2 MTT cytotoxicity assay	45
2.2.2.3 ³ H-Thymidine incorporation assay	46
2.2.2.4 Differentiation inducing effects of isoflavonoids	47
2.2.2.5 Frequency analysis of proliferative neuroblastoma cells	47
2.2.2 Neuropoietic cytokine expression in biochanin A- and genistein-treated BU-1 cells	48
2.2.2.1 Materials	48
2.2.2.2 Isolation of total RNA by guanidinium thiocyanate / cesium chloride isopycnic gradient method	49
2.2.2.3 Reverse-transcription Polymerase chain reaction (RT-PCR)	51
Table 11. List of Specific primers for different genes	52
2.2.2.4 Cycle titration and dot blot hybridization	53
2.2.2.5 DIG 3' end labeling of probes	53
2.2.2.6 Hybridization	54
2.2.2.7 Chemiluminescent detection	55
2.2.2.8 Quantitation by densitometry	55
Figure 3. Assembly of the dot blot apparatus	56
2.2.3 Immunohistochemical characterization of neurites in differentiating neuroblastoma cells	57
2.2.3.1 Materials	57
2.2.3.2 Induction of neuronal maturation of BU-1 by isoflavonoids	57
2.2.3.3 Immunocytochemistry	58
2.2.3.4 Laser-scan confocal microscopy	58

Chapter Three	Biological Effects of Biochanin A and Genistein on Murine Neuro-2A Cell and its Subclones, BU-1 and BU-2	59
3.1	Introduction	
3.1.1	Anti-proliferative effects of flavonoids on mammalian cancer cells	59
3.1.2	Biological effects of biochanin A and genistein on Neuro-2A and its subcloned cells	59
3.1.2.1	Cytotoxicity	60
3.1.2.2	Growth inhibitory effects	61
3.1.2.3	Differentiation inducing effects	61
3.2	Results	63
3.2.1	Cytotoxicity of biochanin A and genistein on neuroblastoma cells	63
3.2.2	Growth inhibitory effects of biochanin A and genistein	63
3.2.3	Differentiation inducing effects of biochanin A and genistein	64
3.2.3.1	Morphological differentiation of neuroblastoma cells	64
3.2.3.2	Phenotypes of differentiated neuroblastoma cells	66
3.2.3.3	Frequency analysis on the number of terminally differentiated neuroblastoma BU-1 cells	67
	Table 12. Frequency of proliferative BU-1 cells after isoflavonoids treatment	68
	Figure 4. Cytotoxic effects of isoflavonoids on Neuro-2A cells	69
	Figure 5. Cytotoxic effects of isoflavonoids on BU-1 cells	70
	Figure 6. Cytotoxic effects of isoflavonoids on BU-2 cells	71
	Figure 7. Growth inhibitory effects of isoflavonoids on Neuro-2A cells	72
	Figure 8. Growth inhibitory effects of isoflavonoids on BU-1 cells	73
	Figure 9. Growth inhibitory effects of isoflavonoids on BU-2 cells	74
	Figure 10. Differentiation inducing effects of isoflavonoids on Neuro-2A cells	75
	Figure 11. Differentiation inducing effects of isoflavonoids on BU-1 cells	76
	Figure 12. Differentiation inducing effects of isoflavonoids on BU-2 cells	77

Figure 13. Photomicrographs of biochanin A and genistein induced morphological changes in parent Neuro-2A cells	78
Figure 14. Photomicrographs of biochanin A and genistein induced morphological changes in subclone BU-1 cells	80
Figure 15. Photomicrographs of biochanin A and genistein induced morphological changes in subclone BU-2 cells	82
Figure 16. Frequency analysis of biochanin A treated BU-1 cells (50 μ M, 3 days)	84
Figure 17. Frequency analysis of biochanin A treated BU-1 cells (50 μ M, 6 days)	85
Figure 18. Frequency analysis of biochanin A treated BU-1 cells (100 μ M, 3 days)	86
Figure 19. Frequency analysis of biochanin A treated BU-1 cells (100 μ M, 6 days)	87
Figure 20. Frequency analysis of genistien treated BU-1 cells (25 μ M, 3 days)	88
Figure 21. Frequency analysis of genistien treated BU-1 cells (25 μ M, 6 days)	89
Figure 22. Frequency analysis of genistien treated BU-1 cells (50 μ M, 3 days)	90
Figure 23. Frequency analysis of genistien treated BU-1 cells (50 μ M, 6 days)	91
3.3 Discussion	
3.3.1 Cytotoxicity of biochanin A and genistein	92
3.3.2 Growth inhibitory effects of biochanin A and genistein	94
3.3.3 Differentiation inducing effects of biochanin A and genistein	95
Chapter Four Molecular Analysis of Neuroipoietic Cytokine Expression in Biochanin A- and Genistein-treated BU-1 cells	98
4.1 Introduction	98

4.1.1 Cytokines in neuronal cell growth and differentiation	99
4.2 Results	101
4.2.1 Analysis of total RNA	101
Table 13. Concnetration of various RNA samples	101
4.2.2 mRNA phenotyping	102
4.2.2.1 Expression Profile of IL-6	102
4.2.2.2 Expression profile of CNTF	103
4.2.2.3 Expression profile of LIF	103
4.2.2.4 Expression profile of LIFR-L and LIFR-S	104
Figure 24A. Normalization of samples by GADPH gene expression	105
Figure 24B. Normalization of samples by GADPH gene expression	105
Figure 25. Cycle titration of IL-6 mRNA prepared from biochanin A treated BU-1 cells	106
Figure 26. Cycle titration of IL-6 mRNA prepared from genistein treated BU-1 cells	107
Figure 27. Cycle titration of CNTF mRNA prepared from biochanin A treated BU-1 cells	108
Figure 28. Cycle titration of CNTF mRNA prepared from genistein treated BU-1 cells	109
Figure 29. Cycle titration of LIF mRNA prepared from biochanin A treated BU-1 cells	110
Figure 30. Cycle titration of LIF mRNA prepared from genistein treated BU-1 cells	111
Figure 31. Cycle titration of LIR-S mRNA prepared from biochanin A treated BU-1 cells	112
Figure 32. Cycle titration of LIFR-S prepared from genistein treated BU-1 cells	113
Table 14. Summary of expression profiles of various neuropoietic cytokines and receptors in BI-1 cells	114
4.3 Discussion	115
4.3.1 mRNA phenotyping	115

Chapter Five Immunohistochemical Characterization of Neuritic Outgrowth in Differentiating BU-1 and BU-2 cells	119
5.1 Introduction	119
5.1.1 Architecture and cytoskeleton of neurons	119
5.1.2 Microtubules: Microtubule-associated proteins (MAPs)	120
5.1.3 Intermediate filaments: Neurofilament (NFs)	121
5.2 Results	123
5.2.1 Immunoreactivity of MAP-2 in biochanin A- and genistein-treated BU-1 cells	123
5.2.2 Immunoreactivity of tau in biochanin A- and genistein-treated BU-1 cells	124
5.2.3 Immunoreactivity of NF-H in biochanin A- and genistein-treated BU-1 cells	124
Figure 33. Photomicrographs of immunoreactivity of MAP-2 after 3 days isoflavonoids treatment	126
Figure 34. Photomicrographs of immunoreactivity of MAP-2 after 6 days isoflavonoids treatment	128
Figure 35. Photomicrographs of immunoreactivity of tau after 3 days isoflavonoids treatment	130
Figure 36. Photomicrographs of immunoreactivity of tau after 6 days isoflavonoids treatment	132
Figure 37. Photomicrographs of immunoreactivity of NF-H after 3 days isoflavonoids treatment	134
Figure 38. Photomicrographs of immunoreactivity of NF-H after 6 days isoflavonoids treatment	136
Table 15. Relative intensity of immunoreactivity of MAP-2 in isoflavonids treated BU-1 cells	138
Table 16. Relative intensity of immunoreactivity of tau in isoflavonids treated BU-1 cells	138

Table 17. Relative intensity of immunoreactivity of NF-H in isoflavonoids treated BU-1 cells	139
Table 18. Percentage of NF-H positive neurons in isoflavonoids treated BU-1 cells	139
5.3 Discussion	140
5.3.1 Immunoreactivity of MAP-2	140
5.3.2 Immunoreactivity of tau	142
5.3.3 Immunoreactivity of NF-H	143
5.3.4 Characterization of neuritic outgrowth in differentiating BU-1 cells	144
Chapter Six General Discussion	147
6.1 Anti-tumor effects of biochanin A and genistein	147
6.2 Differentiation inducing effects of biochanin A and genistein	150
6.3 Neuropoietic control of isoflavonoids-induced differentiation	152
6.4 Further studies of control of differentiation on neuroblastoma cells	153
6.5 Conclusion	155
Appendix	157
References	159
Curriculum Vitae	178