

## MASTER'S THESIS

### Effects of agronomic practices on aluminium and fluoride concentrations in soil and tea plants

Fung, Ka Fai

*Date of Award:*  
1999

[Link to publication](#)

#### General rights

Copyright and intellectual property rights for the publications made accessible in HKBU Scholars are retained by the authors and/or other copyright owners. In addition to the restrictions prescribed by the Copyright Ordinance of Hong Kong, all users and readers must also observe the following terms of use:

- Users may download and print one copy of any publication from HKBU Scholars for the purpose of private study or research
- Users cannot further distribute the material or use it for any profit-making activity or commercial gain
- To share publications in HKBU Scholars with others, users are welcome to freely distribute the permanent URL assigned to the publication

Effects of Agronomic Practices on Aluminium and Fluoride  
Concentrations in Soil and Tea Plants

FUNG Ka Fai

A thesis submitted in partial fulfillment of the requirements

for the degree of

Master of Philosophy

December 1999

Hong Kong Baptist University

## Abstract

Tea is one of the most popular beverages in the world and it is well-known accumulator of Al and F. The present study aims to investigate the effects of different agronomic practices on Al and F uptake by tea plants, and to find ways to reduce the concentration of these elements within the plants. A reduction is desirable as Al and F absorbed will leach into the tea liquor through infusion thus posing a serious threat to the health of the consumers of tea.

The soil pH of the tea plantation (4-4.52) was lower than that in the normal soil obtained near the tea plantation on Lantau Island, Hong Kong. The soil extractable Al and F concentration ranged from 168-193 Al mg/kg and 1.41- 2.61 F mg/kg, and were found to be inversely proportional to the soil pH ( $p < 0.001$ ). Extractable of both species decreased ( $p < 0.05$ ) gently along the soil profile. The ecological survey indicated that the extractable amount of Al and F and their concentrations within the plants were directly proportional to the age of the plantation. Al accumulated in old leaves and roots were quantified to be 15,268 mg/kg and 10,101 mg/kg in average, and F accumulated in old tea leaves was 2072 mg/kg. The average amount of Al and F in young leaves, which makes up of the bulk of commercial tea products, were, 7017 mg/kg and 310 mg/kg respectively. Among the various species tested, "small leafed variety" had the highest ( $p < 0.05$ ) Al and F concentration, followed by "large leafed variety" and Assam variety. Aluminium and F contents in black tea were higher than those in green tea, and both elements' concentrations in tea liquor under repeated infusion was slightly higher than in continuous infusion.

Acidifying materials and lime are commonly applied on tea plantations. All acidifying materials included in this studies ( $\text{Al}_2(\text{SO}_4)_3$ ,  $\text{H}_2\text{SO}_4$ ,  $(\text{NH}_4)_2\text{SO}_4$  and  $\text{NH}_4\text{NO}_3$ ) increased extractable Al and F concentration but decrease available nutrient contents (such as P and K) in soil while calcium-containing materials ( $\text{CaCO}_3$ ,  $\text{CaCl}_2$  and  $\text{CaSO}_4$ ) reduced extractable Al and F but maintained the normal nutrient level. Therefore, Ca-containing materials were chosen for the purpose of Al and F uptake reduction by tea seedlings.

Extractable Al, F, Fe, Mn and Zn concentrations decreased whereas extractable Ca, Cu, K, Mg, Na and P concentration increased when the soil pH was raised from 3 to 6. The three growth measurement, relative dry weight gain (RDW), relative leaf number gain (RLN) and relative leaf area gain (RLA), decreased when the soil pH increased which resulted in a decrease in soil extractable Al and F. Tea seedlings grown in soils of pH 3.0 and 3.5 were the tallest and the healthiest in both varieties while those in pH 6.0 died after 3 months of cultivation. "Large leafed variety" was noted to display higher growth rates than "small leafed variety". The highest ( $p < 0.05$ ) amount of Al and F, 4225 mg/kg and 430 mg/kg, were found in seedlings under treatment of pH 3.5, and they were observed to be directly proportional to the plant growth rate. When Al and F contents increased in the seedlings, nutrient contents such as Ca, Mg, Mn, Na and Zn gently decreased ( $p < 0.05$ ) whereas P and Fe concentrations increased. Potassium and Cu were not

influenced by varying the amount Al and F. It has been suggested high pH values (> 4) can reduce the Al and F concentrations in tea plants.

All true lime material ( $\text{CaCO}_3$ ,  $\text{Ca(OH)}_2$  and  $\text{CaO}$ ) increase the soil pH and decrease soil extractable Al and F concentrations. There are no ( $p < 0.05$ ) change of pH when  $\text{CaCl}_2$  was added, and a slight decrease in Al and F concentrations was observed in the treatments with  $\text{CaSO}_4$ . Tea seedlings under the addition of  $\text{CaSO}_4$  were the tallest and the healthiest, followed by  $\text{CaCl}_2$ ,  $\text{CaCO}_3$ ,  $\text{Ca(OH)}_2$  and  $\text{CaO}$ . The treatment with 4,000 mg  $\text{CaSO}_4$  /kg yield the tallest seedlings and the height decreased subsequently ( $p < 0.05$ ). Seedlings under large doses of  $\text{Ca(OH)}_2$  and  $\text{CaO}$ , there were dead seedlings. RDW, RLN and RLA decreased significantly with increasing calcium compounds added except in the treatment with  $\text{CaSO}_4$ . The highest Al concentration was also found in the treatment with 4,000 mg  $\text{CaSO}_4$  /kg, it then decreased gently from 3364 to 933 mg Al/kg in leaves. For all treatments in both varieties, F concentrations in leaves were lower than the control (374 mg/kg). Therefore,  $\text{CaSO}_4$  is a material capable of accelerating growth of tea seedlings, without reducing the uptake of Al and F. The growth rate, as well as Al and F concentrations, were greatly reduced in the true lime treatments. Calcium chloride was a suitable agent reduction of Al and F uptake by Al and F while maintaining a high growth rate of the seedlings.

The present experiment indicated that there was a decrease in Al and F concentrations in soil and tea plants with the addition of organic amendments applied to soil. the amount of organic Al concentration in soil was directly proportional to ( $p < 0.001$ ) the values of CEC and TOC in the soil. Tea plants exhibited poorer growth when the application rates of the organic amendments were increased, and death was observed at higher doses. The highest growth rate (RDW, RLN and RLA values) was found in the treatments of 1% peat moss (361-398% in average). Peanut cake was not a suitable amendment for tea plantation as it reduced Al and F concentrations in both soil and plants, and resulted in death of the plants. Although the leaves of seedlings did not grow so healthily, pig manure compost could still be one of the organic amendments for reducing Al and F availability to tea plants. Peat moss, when adjusted to a suitable pH (e.g. pH 4.5), could be another suitable organic amendment, for it was effective in reducing Al and F availability while promoting healthy plant growth.

# TABLE of CONTENTS

	Page
<b>Declaration</b>	i
<b>Abstract</b>	ii
<b>Acknowledgements</b>	iv
<b>Table of Contents</b>	v
<b>Lists of Tables</b>	ix
<b>List of Figures</b>	xi
<b>Chapter I. Introduction</b>	
1.1 Botanical classification of tea	1
1.2 Condition of tea cultivation	2
1.3 Types of tea leaves and their production	2
1.3.1 Black tea	3
1.3.2 Green tea	3
1.3.3 Oolong tea	4
1.3.4 Brick tea	4
1.4 Aluminium in soil, in tea soil and in tea	4
1.5 Potential health hazard of Al in tea	6
1.6 Fluoride in soil, in tea soil, and in tea	6
1.7 Potential health hazard of F in tea	7
1.8 Aim and objectives of the research	9
<b>Chapter II. Aluminium and Fluoride Concentrations of Three Tea Varieties Growing on Lantau Island of Hong Kong</b>	
2.1 Introduction	11
2.2 Materials and methods	13
2.2.1 Site description	13
2.2.2 Soil and tea bushes samplings	14
2.2.3 Soil analysis	14
2.2.4 Plant analysis	15
2.2.5 Infusion methods	15
2.2.6 Statistical analysis	16
2.3 Results	16
2.3.1 General soil properties, Al and F contents of surface soils of tea plantation	16
2.3.2 General soil properties, Al and F contents of soil profiles	20
2.3.3 Distribution of Al and F in tea plants	25
2.3.4 Metal contents and phosphorus contents	27
2.3.5 Total Al and F contents in tea products and effects of infusion methods on Al and F contents of tea liquor	32

2.3.5.1 <i>Repeated infusion</i>	33
2.3.5.2 <i>Continuous infusion</i>	33
2.4 Discussion	35
2.4.1 Soil properties in tea soil	35
2.4.2 Distribution of Al and F different tea varieties	36
2.4.3 Al and F in tea infusions	37
2.4.4 Al and F contamination in production process	38
2.5 Conclusions	39

**Chapter III. Effects of Various Soil Ameliorants on Soil Extractable Aluminium and Fluoride Concentrations**

3.1 Introduction	40
3.1.1 Common ameliorants applied to tea soil	40
3.1.2 Aluminium solubility and the interaction with other elements in soil	41
3.2 Materials and methods	44
3.3 Results	
3.3.1 pH, extractable aluminium and fluoride concentrations	45
3.4 Discussion	52
3.5 Conclusion	54

**Chapter IV. The effects of Soil pH on the Uptake of Al and F by Tea Plants**

4.1 Introduction	55
4.2 Materials and methods	58
4.2.1 Experiment Setup	58
4.2.2 Soil and plant analyses	60
4.2.3 Plant growth rate analysis	62
4.2.4 Statistics	63
4.3 Results	63
4.3.1 Comparison of extractable aluminium, fluoride, phosphorus and other metals contents in the pH dependent experiment	63
4.3.2 the growth study of tea seedlings after six months	65
4.3.3 Distribution of Al and F contents in tea seedlings	65
4.3.4 Distribution of other nutrient contents in tea seedlings	68
4.3.5 Relationships of Al and F with other elements in soil and in tea seedlings	77
4.3.6 Relationships of Al and F with other elements in roots, in leaves, and between roots and leaves	80
4.3.7 Relationships of Al and F, and the relative growth rates of tea seedlings	80
4.4 Discussion	80
4.4.1 Extractable aluminium, fluoride, phosphorus and other metals contents	80

4.4.2 The growth study of tea seedlings after the six -months treatments	81
4.4.3 Distribution of Al and F contents and other nutrients in tea seedlings.	81
4.5 Conclusion	85

**Chapter V: Application of Different Forms of Calcium to Tea Soil  
(Liming) in Order to Prevent Al and F Accumulation in  
Tea Plants**

5.1 Introduction	86
5.2 Materials and methods	90
5.2.1 Experimental setup	90
5.2.2 Soil and plant analysis	92
5.2.3 Plant growth rate analysis	93
5.2.4 Statistics	93
5.3 Results	93
5.3.1 Comparison of soil properties after application of calcium compounds	93
5.3.2 Growth performance of tea seedlings after six -months	97
5.3.3 Distribution of Al and F contents in tea seedlings	102
5.3.4 Distribution of other nutrient contents in tea seedlings	104
5.3.5 Relationships of Al and F with other elements in the soil and in tea seedlings	113
5.3.6 Relationships of Al and F with other elements in roots, in leaves, and between roots and leaves	113
5.3.7 Relationships of Al and F, and the relative growth rates of tea seedlings	116
5.4 Discussion	116
5.4.1 Comparison of extractable aluminium, fluoride, phosphorus and other metals contents in calcium application experiment	116
5.4.2 growth performance of tea seedlings after six- months	118
5.4.3 Distribution of Al and F contents in tea seedlings	119
5.5 Conclusion	120

**Chapter VI. The Effects of Organic Amendments for Reducing the  
Uptake of Al and F of Tea Plant**

6.1 Introduction	121
6.2 Materials and methods	124
6.2.1 Experimental setup	124
6.2.2 Soil and plant analyses	125
6.2.3 Plant growth rate analysis	127
6.2.4 Statistics	127
6.3 Results	127

6.3.1 Comparison of soil properties after treatment of organic amendments	127
6.3.2 The growth of tea seedlings applied with different soil amendments after six-months	130
6.3.3 Distribution of Al and F contents in tea seedlings	134
6.3.4 Distribution of other nutrient contents in tea seedlings	137
6.3.5 Relationships of Al and F with other elements in the soil and in tea seedlings	137
6.4 Discussion	147
6.4.1 Comparison of extractable aluminium, fluoride, phosphorus and other metals contents in organic amendments application experiment	147
6.4.2 The growth of tea seedlings after six-months' treatments	150
6.4.3 Distribution of Al and F contents in tea seedlings	151
6.5 Conclusion	152
<b>Chapter VII. General Discussion and Conclusion</b>	
7.1 The role of aluminium and fluoride in tea plants and their effects on growth rates	153
7.1.1 Aluminium	153
7.1.2 Fluoride	156
7.2 Nutrient contents in tea plants	157
7.3 Factors affecting bioaccumulation of Al and F by tea plants	158
7.4 General conclusion	160
7.5 Comments on future studies	161
<b>Appendices</b>	162
<b>References</b>	186
<b>List of papers published, accepted and submitted to journals or presented in meetings</b>	198
<b>Curriculum Vitae</b>	199