

DOCTORAL THESIS

Functional studies of gill epithelial cells isolated from Japanese eels (*anguilla japonica*)

Tse, Ka Fai William

Date of Award:
2008

[Link to publication](#)

General rights

Copyright and intellectual property rights for the publications made accessible in HKBU Scholars are retained by the authors and/or other copyright owners. In addition to the restrictions prescribed by the Copyright Ordinance of Hong Kong, all users and readers must also observe the following terms of use:

- Users may download and print one copy of any publication from HKBU Scholars for the purpose of private study or research
- Users cannot further distribute the material or use it for any profit-making activity or commercial gain
- To share publications in HKBU Scholars with others, users are welcome to freely distribute the permanent URL assigned to the publication

**Functional Studies of Gill Epithelial Cells Isolated from Japanese
Eels (*Anguilla japonica*)**

TSE Ka Fai William

A thesis submitted in partial fulfillment of the requirements

for the degree of

Doctor of Philosophy

Principal Supervisor: Dr. WONG Kong Chu

Hong Kong Baptist University

January 2008

Abstract

Eel (*Anguilla japonica*) is a euryhaline fish that resides both in freshwater and seawater. The fish, however, encounters totally opposite osmotic challenges in the two media. In freshwater, the fish deals with osmotic water uptake and loss of salts and vice versa in seawater. To tolerate the wide range of salinities during migration, they develop special structures and mechanisms for osmoregulation, of which are the chloride cells (CCs) and pavement cells (PVCs) in the gill epithelium. Over the years, studies on the ion regulation in gills were mainly based on the electrophysiological, biochemical, and immunocytochemical staining methods on whole gill preparation. Although a great deal of information has been gathered, the direct interaction of the hormones and osmotic effects with individual type of gill cells in mediating cellular mechanisms in ion transport is still not fully understood. The progress of the study is basically hampered by the anatomical complexity of the gill tissues and the limited nucleotide databases as well as the commercially available antibodies for fish protein.

In the present study, we developed a model that is able to extract purified isolated chloride cells (CCs) and pavement cells (PVCs) from the gill of acclimating fish as well as to establish the primary gill cell culture to study the effect of osmotic stress on the expression profiles of ion transporters and ion channels. To increase the availability of molecular tools in studying fish osmoregulation, in chapter 2 we have cloned a number of ion transporters/channels and osmotic-stress induced transcription factor from the Japanese eel model. Partially cDNAs of sodium-hydrogen exchanger 3 (NHE3), sodium-bicarbonate co-transporters (NBC1), vacuolar H⁺-ATPase and osmotic stress transcription factor 1 (Ostf1) were cloned. Real-time PCR primers were designed for mRNA quantification. For the NHE3, NBC1 and vacuolar H⁺-ATPase, their expression levels were in general higher in freshwater gill cells. Our data indicated that the expression level of Ostf1 was significant higher in freshwater CCs than that of the PVCs. Upon seawater transfer, significant inductions of Ostf1 transcripts were observed in both CCs and PVCs. Interestingly in seawater to freshwater transfer experiments, there was no significant change in Ostf1 expression levels. Our data also indicated that the activation of Ostf1 expression was sensitive to water Na⁺ and Cl⁻ levels.

In chapters 3 & 4, we aimed to determine the expression profiles of ion transporters and ion channels in CCs and PVCs of freshwater- and seawater-acclimating eels. Upon freshwater to seawater transfer (Chapter 3), significant increase of the mRNA and/or protein expressions of Na⁺/K⁺-ATPase α and β subunits, cystic fibrosis transmembrane conductance regulator (CFTR), sodium potassium chloride co-transporter (NKCC), and inwardly rectifying potassium channel (eKir) were found in gill CCs. However, reduction of aquaporin 1 and 3 (AQP1, and 3) transcript levels

were detected. While in seawater to freshwater transfer (Chapter 4), increased phosphorylation of some signaling proteins like mitogen-activated protein kinases (MAPK), myristoylated alanine-rich C-kinase substrate (MARCKs), and cAMP response element-binding (p-CREB) were observed. The expression levels of the transporters like anion exchanger 1 (AE1), NHE3, NBC1, and V-H⁺-ATPase were significantly increased in freshwater CCs. The results suggested that the rapid response of signaling molecules in freshwater acclimation modulated the expression levels of different ion transporters for osmoregulation. In these two chapters, we provide the first evidence to demonstrate the expression profiles of different ion transporters and ion channels in individual cell types in fish gills. Our data demonstrated that the hyper- or hypo-osmotic adaptation remodeled the tissue distribution of CCs as well as the differential gene expression in the cells. The study support the notion that gill CCs are important in both freshwater and seawater adaptation.

To elucidate the importance of freshwater CCs in Na⁺ and Cl⁻ uptake, in chapter 5 we attempted to establish an experimental model to study the reactivation mechanism in branchial ion uptake. Deionized water adapted eels were transferred to artificial freshwater (ion-supplement deionized water, ISDI) with or without Na⁺ and/or Cl⁻. Upon the transfer, our data demonstrated that the fish acclimating in ISDI without Na⁺, showed significant inductions in the expression of the ion transporters (i.e. NHE1, NHE3, NBC1, and V-H⁺-ATPase) in gill CCs. In the acclimation at ISDI without Cl⁻, significant inductions of AE and NKCC in gill CCs were detected. Comparatively, the changes in the transporter expression levels were trifling in gill PVCs. Our study demonstrated that the gill CCs were the major targets in the reactivation of ion absorption. The data further suggest the importance of freshwater CCs in ion uptake.

In the last chapter (Chapter 6), we have developed a primary gill cell culture model and studied the direct effect of osmotic stress on cell volume disturbance and the expression of ion transporters in the cells. Our data demonstrated that the hypertonic (500 mOsm) treatment of PVCs induced cell shrinkage, followed by regulatory volume increase (RVI). The cell volume regulation was found to be modulated by the transporters – Na⁺/K⁺-ATPase, NKCC, and NHE1. The stimulators, dexamethasone (DEX) and dibutyryl cAMP treatments significantly induced the expression levels of the three ion transporters. Collectively, this thesis represents the first to combine *in-vivo* and *in-vitro* studies to obtain a multi-facet understanding on the gill cell functions.

Table of Contents

Declaration	i
Abstract	ii
Acknowledgement	iv
Table of Contents	v
List of Figures	ix
List of Abbreviation	xv
Chapter 1 Literature Review	1
1.1 Eel	1
1.2 Gill	4
1.3 Ion Transporters and Channels	10
i Anion Exchanger (AEs)	13
ii Aquaporins (AQPs)	14
iii Cystic Fibrosis Transmembrane Conductance Regulator (CFTR)	16
iv Sodium Bicarbonate Co-transporter (NBC)	18
v Na ⁺ /K ⁺ -ATPase	19
vi Sodium-hydrogen Exchanger (NHE)	20
vii Na ⁺ /K ⁺ /2Cl ⁻ Co-transporter (NKCC)	21
viii Inwardly Rectifying Potassium Channel (eKir)	22
ix Vacuolar Proton ATPase (V-H ⁺ -ATPase)	23
1.4 Hormonal Effects on FW and SW Adaptation	24
i Glucocorticoid	26
ii Growth Hormone (GH) and Insulin-like Growth Factor (IGF)	28
iii Thyroid Hormones	29
iv Prolactin (PRL)	30
Working Hypothesis	31
Chapter 2 The Cloning of Eel Osmotic Stress Transcription Factor 1 (Ostf1); Sodium-hydrogen Exchanger 3 (NHE3); Sodium-bicarbonate Co-transporters (NBC1); and Vacuolar H⁺-ATPase (V-H⁺-ATPase) and the Determination of their mRNA Expression Levels in Isolated Pavement Cells (PVCs) and Chloride Cells (CCs) at Different Osmotic Conditions	32

	Abstract	32
2.1	Introduction	33
2.2	Materials and Methods	35
2.3	Results	39
	i Cloning and Characterization of Eel Ostf1 in Eel's Gills	39
	ii Cloning and Characterization of Eel NBC1, NHE3, and V-H ⁺ -ATPase in Eel's Gills	42
	iii The Differential Expression of Ostf1 mRNA in PVCs and CCs at Freshwater to Seawater and Seawater to Freshwater Transfer Experiments	47
	iv Ostf1 mRNA Expression in PVCs and CCs in the DI to ISDI Transfer Experiments	49
	v mRNA Expressions of NBC1, V-H ⁺ -ATPase, and NHE3 in Freshwater and Seawater PVCs and CCs	50
2.4	Discussion	52
Chapter 3 Characterization of Ion Channel and Transporter mRNA Expressions in Isolated Gill Chloride and Pavement Cells of Seawater Acclimating Eels		56
	Abstract	56
3.1	Introduction	57
3.2	Materials and Methods	59
3.3	Results	63
	i Expression Profile of Ion Channels and Transporters	63
	ii Western Blot and Immunocytochemical (ICC) Studies of the Ion Channel and Transporters	70
3.4	Discussion	75
Chapter 4 Characterization of Ion Channel and Transporter mRNA Expressions in Isolated Gill Chloride Cells of Freshwater Acclimating Eels		79
	Abstract	79
4.1	Introduction	80
4.2	Materials and Methods	82
4.3	Results	86
	i Plasma Osmolarity and Na ⁺ and Cl ⁻ Concentrations	86
	ii Protein Expression Profile of Signaling Molecules in Gills	87

iii	Gene Expression Profile of Ion Transporters in Isolated Gill Chloride Cells	88
iv	Immunocytochemistry Staining of Ion Transporters in Gills	91
4.4	Discussion	94
Chapter 5	Selective Activation of V-H⁺-ATPase, Na⁺/HCO₃⁻ Cotransporter-1, Na⁺/H⁺ Exchanger-1 and 3 Expressions in Gill Chloride Cells of Na⁺-deficient Freshwater Adapted Japanese Eel	98
	Abstract	98
5.1	Introduction	99
5.2	Materials and Methods	101
5.3	Results	104
i	Plasma pH, Osmolarity, Na ⁺ and Cl ⁻ Concentrations	104
ii	Gene Expression Profile of Ion Transporters in Isolated Gill PVCs and CCs from Deionized Water Adapted Eels	106
iii	Gene Expression Profiles of Ion Transporters in Isolated Gill PVCs from ISDI Acclimating Fish	107
iv	Gene Expression Profile of Ion Transporters in Isolated Gill CCs and Immunocytochemistry Staining of Gill Filaments after 2 days Transfer from DI to ISDI	108
v	Gene Expression Profile of Ion Transporters in Isolated Gill CCs and Immunocytochemistry Staining of Gill Filaments after 2 days Transfer from DI to Na ⁺ -deficient ISDI	110
vi	Gene Expression Profile of Ion Transporters in Isolated Gill CCs and Immunocytochemistry Staining of Gill Filaments after 2 Days Transfer from DI to Cl ⁻ -deficient ISDI	112
5.4	Discussion	114
Chapter 6	Effect of Osmotic Shrinkage and Hormones on the Expression of Na⁺/H⁺ exchanger-1, Na⁺-K⁺-2Cl Cotransporter and Na⁺/K⁺-ATPase in Gill Pavement Cells of Freshwater Adapted Japanese Eel, <i>Anguilla japonica</i>	117
	Abstract	117
6.1	Introduction	118
6.2	Materials and Methods	121
6.3	Results	125

	i Effect of Hypertonic Treatment on the Cell Size and the expressions of NHE1, NKCC and Na ⁺ /K ⁺ -ATPase in PVCs	125
	ii Effect of the Hormones (DEX, prolactin, T ₃ and IGF-1), dbcAMP and dbcGMP on the Expression of Ion Transporters in the Primary PVC Culture	129
6.4	Discussion	133
	Conclusion Remarks	138
	List of References	140
	Curriculum Vitae	165