

MASTER'S THESIS

Impacts of *Pomacea canaliculata* on freshwater macrophytes and water quality, as well as its control with fish

Wong, Pak Ki

Date of Award:
2010

[Link to publication](#)

General rights

Copyright and intellectual property rights for the publications made accessible in HKBU Scholars are retained by the authors and/or other copyright owners. In addition to the restrictions prescribed by the Copyright Ordinance of Hong Kong, all users and readers must also observe the following terms of use:

- Users may download and print one copy of any publication from HKBU Scholars for the purpose of private study or research
- Users cannot further distribute the material or use it for any profit-making activity or commercial gain
- To share publications in HKBU Scholars with others, users are welcome to freely distribute the permanent URL assigned to the publication

**Impacts of *Pomacea canaliculata* on Freshwater Macrophytes and Water Quality, as well as
its Control with Fish**

WONG Pak Ki

**A thesis submitted in partial fulfillment of the requirements
for the degree of
Master of Philosophy**

Principle Supervisor: Dr. QIU Jianwen

Hong Kong Baptist University

February 2010

Abstract

The golden apple snail (*Pomacea canaliculata* Lamarck, 1819) is native to freshwater wetlands of South America, but has invaded many Asian countries and grazed heavily in agricultural and natural habitats. This species has become the focus of research in recent years, with the main targets of assessing its impact and reducing its grazing damage. However, several aspects of its trophic relationship remain understudied. Previous studies of the grazing of *P. canaliculata* have focused on its direct effects on macrophytes, but have neglected the indirect effects on water quality. Feeding preference of this invader on local wild macrophytes and plant traits that determine such preference are unknown. Besides, although common carp (*Cyprinus carpio*) has been proposed as a biological control agent against *Pomacea canaliculata*, its impact on non-target aquatic plants and animals has not been assessed.

This study aimed to provide information on the direct and indirect effects of the grazing by *P. canaliculata* on local wetlands, understand the plant traits that determine its feeding preference, and assess the effectiveness as well as the potential side effects of its biological control using common carp. To quantify the direct and indirect snail grazing effects, my colleagues and I confined a known amount of *Myriophyllum aquaticum* or *Eichhornia crassipes* with 0, 2, 4 or 8 apple snails in 1 x 1 x 1 m enclosures for approximately 1 month in a freshwater pond. Apple snails grazed heavily on both species of macrophytes, with higher overall weight losses at higher snail densities. The damage patterns differed between the two macrophytes. In *M. aquaticum*, both leaves and stems suffered from substantial herbivory, whereas in *E. crassipes*, only the roots suffered significant weight reduction. In addition to grazing on macrophytes, apple snails appeared to have controlled the growth of filamentous algae, as these did not develop in the snail treatments. The ability of *P. canaliculata* to control filamentous algae was supported by a laboratory experiment where the consumption was as high as 0.25 g g⁻¹ snail DW d⁻¹. Nitrogen and phosphorous concentrations remained low throughout both experiments, and were not correlated with apple snail

density. The treatment effects on chlorophyll *a* (Chl *a*) and phytoplankton composition varied in the two experiments. In the *M. aquaticum* experiment, with increasing snail density, Chl *a* increased and the phytoplankton community became dominated by Cryptophyceae. In the *E. crassipes* experiment, Chl *a* level was independent of snail density, but with increasing snail density, the phytoplankton community became co-dominated by Cryptophyceae, Chlorophyceae and Bacillariophyceae.

To study the palatability of local macrophytes to *P. canaliculata*, I offered 21 species of freshwater macrophytes to the snail in no-choice feeding assays. I also examined the plant traits that determine the food preference in apple snails. Snail daily feeding rate varied greatly with plant species, ranging from 1.1% to 22% of its body mass. Total nitrogen content was positively correlated, and C:N ratio and dry matter content (DMC) were negatively correlated with snail feeding rate. There was no significant correlation between snail feeding rates and plant phenolic content, but the feeding rate on *Myriophyllum aquaticum* (a plant with the highest phenolic content) was very low. Two macrophytes which bear the highest DMC, viz. *Phragmites australis* and *Vallisneria natans* were eaten much more as reconstituted plant than as fresh leaves, indicating structure (i.e. DMC) to be important in their defence against snail herbivory. The extracts from *M. aquaticum* and *Alternanthera philoxeroides* greatly reduced snail feeding rate, indicating chemical defense to be present in these two species. These results indicate that the apple snail's feeding decision is affected by multiple plants traits. The snail favours plants with high nitrogen and phosphorous contents. High DMC and secondary metabolite (i.e. phenolics and other plant defence chemicals) deter the feeding of the snail.

To examine the effects of common carp on *P. canaliculata* and local freshwater snails and macrophytes, I conducted a 2-month enclosure experiment with three species of aquatic macrophytes and nine species of snails, including the apple snail, in a shallow pond within a plot of freshwater wetland. The results showed that the apple snail or carp alone significantly reduced the

plant biomass, although the apple snail had a stronger herbivorous effect than the carp. The carp completely removed juvenile apple snails, had a weaker predatory effect on adult apple snails, but no effect on the adults' oviposition frequency. Furthermore, the carp significantly reduced the populations of most species of other gastropods that occurred naturally in the pond.

In summary, this study has identified some key plant traits that are useful to predict the direct impact on macrophytes. Given the multiple effects of *P. canaliculata* on wetland biodiversity and function, and the potential significant side effects of using common carp as a biological control agent for this invasive snail, management strategies should be developed to prevent its further spread. In invaded wetlands, strategies should be developed to eradicate the apple snail and reintroduce native snails that can control the development of filamentous algae.

Table of Content

Declaration	i
Abstract	ii
Acknowledgments	v
Table of Content	viii
List of Tables	xi
List of Figures	xii
Chapter 1 General introduction	1
1.1 Research background.....	1
1.1.1 Wetland habitat and biological invasion.....	1
1.1.1 <i>Pomacea canaliculata</i> : one of the 100 worst exotic species around the world.....	2
1.1.2 Control measures for <i>Pomacea</i> spp.	4
1.2 Research objectives and thesis organization	6
1.2.1 Impact of invasive apple snails on wetland macrophytes, nutrients, phytoplankton and filamentous algae in Hong Kong (Chapter 2).....	7
1.2.2 Palatability of macrophytes to the invasive freshwater snail <i>Pomacea canaliculata</i> : differential effects of multiple plant traits (Chapter 3).....	8
1.2.3 Complex interactions among fish, gastropods and macrophytes: implications for biological control of an invasive snail (Chapter 4).....	9
Chapter 2 Impact of invasive apple snails in Hong Kong on wetland macrophytes, nutrients, phytoplankton and filamentous algae	11
2.1 Abstract	11
2.2 Introduction	12

2.3 Methods	15
2.3.1 Characteristics of the study site	15
2.3.2 Field experiments.....	15
2.3.3 Laboratory experiment.....	19
2.4 Results	19
2.4.1 The <i>M. aquaticum</i> experiment.....	19
2.4.2 The <i>E. crassipes</i> experiment.....	22
2.4.3 Grazing rate on filamentous algae	25
2.5 Discussion	25

Chapter 3 Palatability of macrophytes to the invasive freshwater snail *Pomacea canaliculata*:

differential effects of multiple plant traits	42
3.1 Abstract.....	42
3.2 Introduction	44
3.3 Methods	46
3.3.1 <i>Pomacea canaliculata</i> consumption of fresh plant leaves.....	46
3.3.2 <i>Pomacea canaliculata</i> consumption of reconstituted plants	47
3.3.3 Macrophyte chemical defense	48
3.3.4 Plant traits	50
3.3.5 Statistical analyses	50
3.4 Results	52
3.4.1 Palatability of fresh macrophyte leaves	52
3.4.2 Palatability of reconstituted plants.....	53
3.5 Discussion	55
3.5.1 Determinants of plant palatability	55
3.5.2 Implications for wetland management.....	58

Chapter 4 Complex interactions among fish, snails and macrophytes: implications for biological control of an invasive

snail.....63

4.1 Abstract..... 63

4.2 Introduction 64

4.3 Methods 66

4.3.1 Study site 66

4.3.2 Experimental design 67

4.3.3 Sampling and analysis 70

4.4 Results 71

4.4.1 Fish survival and effects of fish on apple snails..... 71

4.4.2 Effects of fish on non-Pomacea snails..... 72

4.4.3 Effects of fish and apple snails on macrophytes..... 73

4.5 Discussion 74

4.5.1 Effects of apple snails on plant biomass..... 74

4.5.2 Effects of apple snails on other snails..... 76

4.5.3 Effects of common carp on macrophytes and snails 78

4.5.4 Management implications..... 80

Chapter 5 General summary and discussion..... 86

5.1 Contribution to current knowledge of *P. canaliculata* 86

5.2 Implications for management 88

6.3 Further studies 89

Chapter6 List of References..... 92

Curriculum Vitae 110

Publication..... 110