

MASTER'S THESIS

Coordination of division timing for intestine precursor cells during *C.elegans* embryogenesis

Wong, Ming Kin

Date of Award:
2016

[Link to publication](#)

General rights

Copyright and intellectual property rights for the publications made accessible in HKBU Scholars are retained by the authors and/or other copyright owners. In addition to the restrictions prescribed by the Copyright Ordinance of Hong Kong, all users and readers must also observe the following terms of use:

- Users may download and print one copy of any publication from HKBU Scholars for the purpose of private study or research
- Users cannot further distribute the material or use it for any profit-making activity or commercial gain
- To share publications in HKBU Scholars with others, users are welcome to freely distribute the permanent URL assigned to the publication

Abstract

Metazoan development is a complex and tightly controlled process that not only requires precise cell fate differentiation, but also demands accurate timing of cell division and precise cell migration. Genetic regulation of cell cycle length throughout metazoans embryogenesis is largely unknown, mainly due to the technical hurdle in quantifying cell division timing during development. *Caenorhabditis elegans* embryogenesis provides an excellent opportunity to study the genetic regulation of division timing because of its invariant cell lineage and widespread division asynchronies between sister cells. A combination of in toto imaging and automated cell lineaging coupled with high throughput RNAi allows genetic screening of genes involved in regulation of Asynchrony of Division between Sister cells (ADS) or cousin cells. One of the most pronounced asynchronies between cousin cells during *C. elegans* embryogenesis is a significant elongation of division timing in two endoderm progenitor cells, Ea and Ep (E2), versus their cousins MSa and MSp (MS2) that mainly develop into mesoderm organs. Out of a total of 822 essential and conserved genes that were perturbed by RNAi in our previous genetic screening, 53 genes are found to produce significantly reduction specifically in the E2 cell cycle length ($p < 0.01$). Surprisingly, nearly 70% of the 53 genes are involved in mRNA production or its regulation, indicating a differential requirement of transcription for division timing between E2 and MS2. Reduction in E2 cell cycle length is frequently associated with cell migration defect and gastrulation failure. Furthermore, our systematic data on cell division timings upon perturbation of a large cohort of essential genes provide a valuable source for inferring the function of uncharacterized gene. For example, phenotypical clustering based on cell division timings suggested that an essential gene, *gad-1*, is likely to be involved in general transcription, which is in agreement with its further functional assays. In summary, a combining of the published data with our own demonstrates that E2 specific cell cycle elongation requires robust and earlier zygotic genome activation (ZGA) during *C. elegans* embryogenesis. The cell-specific elongation might be important for coordinate fate differentiation, division timing and cell migration of E2 to ensure proper intestine development.

Table of contents

DECLARATION	i
Abstract.....	ii
Acknowledgement.....	iii
Table of contents	iv
List of Tables	vii
List of Figures.....	viii
List of Abbreviations	x
Chapter 1 Introduction	1
1.1. <i>Caenorhabditis elegans</i> embryogenesis.....	1
1.2. Regulation of division asymmetry and asynchrony	3
1.3. Maternal-to-zygotic transition in gene regulation.....	7
1.4. Roles of zygotic genome activation in controlling intestine fate specification.....	9
1.5. Roles of zygotic genome activation in controlling division timing	12
1.6. Cell migration and gastrulation.....	16
1.7. Study objective and approach	18
Chapter 2 Results.....	20
2.1. Screening for genes required for division asynchrony during <i>C. elegans</i> embryogenesis using RNAi followed by automated lineaging	20
2.2. Division timing of intestine progenitor during <i>C. elegans</i> embryogenesis	24
2.2.1. <i>Perturbation of multiple genes in C. elegans embryos specifically reduced cell cycle length in Ea and Ep.....</i>	<i>24</i>
2.2.2. <i>mRNA production pathway may play a major role in E2 cell cycle length elongation</i>	
33	
2.2.3. <i>Robust zygotic transcription activation may be initiated at around E2 stage.....</i>	<i>35</i>

2.2.4.	<i>Cell cycle regulation of E2 may or may not be coupled with intestine fate specification</i>	40
2.2.5.	<i>wee-1.1 functioned downstream of general transcription machinery regulating E2 cell cycle length</i>	43
2.2.6.	<i>E2 division timing elongation might facilitate gastrulation</i>	45
2.3.	<i>gad-1 gene characterization</i>	52
2.3.1.	<i>gad-1 encoded a unknown protein that regulates E2 division timing</i>	52
2.3.2.	<i>Deletion knockout of gad-1 led to larval arrest</i>	53
2.3.3.	<i>Rescue of gad-1 deletion by GAD-1::GFP fusion</i>	56
2.3.4.	<i>Nuclear expression of GAD-1 enriched on transcriptionally active chromosomes</i> .	57
2.3.5.	<i>Zygotic expression of GAD-1 started around 28-cell stage</i>	64
Chapter 3 Conclusion and Discussion		66
3.1.	<i>Developmental regulation of intestinal progenitor cells in C. elegans</i>	66
3.1.1.	<i>Robust zygotic genome activation is required for proper E2 cell cycle timing</i>	66
3.1.2.	<i>Proper cell division timing of E2 facilitate gastrulation</i>	72
3.2.	<i>The possible role of gad-1 in transcription</i>	73
Chapter 4 Materials and Methods		78
4.1.	<i>Worm strains and maintenance</i>	78
4.1.1.	<i>Maintance of worms</i>	78
4.1.2.	<i>Worm strains</i>	78
4.1.3.	<i>Generation of lineaging strain with wee-1.1 homozygous deletion</i>	79
4.1.4.	<i>Generation of deletion rescue strain of gad-1 by GAD-1::GFP</i>	80
4.1.5.	<i>Generation of lineaging strain with gad-1 homozygous deletion rescued by GAD-1::GFP</i>	81
4.1.6.	<i>Generation of gad-1 homozygous deletion strain</i>	82

4.2.	RNAi and microinjection.....	83
4.3.	Microscopy, automated lineaging and gene expression profiling	84
4.3.1.	<i>Time-lapse imaging for automated cell lineaging.....</i>	84
4.3.2.	<i>Single-shot imaging</i>	85
4.3.3.	<i>Z-series imaging and maximum 3D projection of fluorescent protein co-localization</i> <i>86</i>	
4.3.4.	<i>Automated cell lineaging and gene expression profiling</i>	87
4.4.	Statistical analysis of temporal and spatial data.....	87
4.4.1.	<i>Quantification of cell cycle length</i>	87
4.4.2.	<i>Hierarchical clustering of E2 cell cycle length.....</i>	87
4.4.3.	<i>GO analysis.....</i>	88
4.4.4.	<i>Data analysis of C. elegans blastomere-specific RNA-seq data</i>	88
4.4.5.	<i>Data analysis on cell migration.....</i>	89
4.5.	Genetic analysis	89
4.5.1.	<i>Analysis of genetic interaction of wee-1.1</i>	89
4.5.2.	<i>Embryonic zygotic expression detection of gad-1</i>	90
4.6.	Phenotyping and genotyping	90
4.6.1.	<i>Brood size, hatching ratio and survival ratio.....</i>	90
4.6.2.	<i>Single-worm PCR.....</i>	91
4.6.3.	<i>Genotyping of homozygous deletion</i>	91
	Appendices.....	93
	Appendix 1. Data analysis on cell migration (Supplementary).....	93
	References.....	96
	CURRICULUM VITAE.....	116