

DOCTORAL THESIS

Effects of radial oxygen loss (ROL) on arsenic tolerance, uptake and distribution by rice (*Oryza sativa* L.)

Wu, Chuan

Date of Award:
2011

[Link to publication](#)

General rights

Copyright and intellectual property rights for the publications made accessible in HKBU Scholars are retained by the authors and/or other copyright owners. In addition to the restrictions prescribed by the Copyright Ordinance of Hong Kong, all users and readers must also observe the following terms of use:

- Users may download and print one copy of any publication from HKBU Scholars for the purpose of private study or research
- Users cannot further distribute the material or use it for any profit-making activity or commercial gain
- To share publications in HKBU Scholars with others, users are welcome to freely distribute the permanent URL assigned to the publication

**Effects of Radial Oxygen Loss (ROL) on Arsenic
Tolerance, Uptake and Distribution by Rice
(*Oryza sativa* L.)**

WU Chuan

**A thesis submitted in partial fulfillment of the requirements
for the degree of
Doctor of Philosophy**

Principal Supervisor: Prof. WONG Ming Hung

Hong Kong Baptist University

Aug 2011

Abstract

Arsenic (As) contamination is currently an environmental problem that has been receiving increasing concern. Substantial arsenic contamination of paddy soil has now been found in many areas around the world, and has adversely affected the health of millions of people that relies on rice as a staple food source. Therefore, there is an urgency to understand the mechanism of As tolerance and uptake by rice. The results of present study may be very valuable in revealing and explaining the physiological mechanisms of As uptake by rice, and can potentially provide valuable information in screening and breeding rice with low concentrations of As in their grains.

Root aeration, arsenic accumulation and speciation in rice of 20 different genotypes treated with 0.4 mg l^{-1} of As were investigated. Different genotypes had different root anatomy demonstrated by entire root porosity (ranging from 12.43 to 33.21%), which was significantly correlated with radial oxygen loss (ROL) ($R = 0.64$, $P < 0.01$). As accumulation differed among genotypes, but there were no significant differences between Indica and Japonica subspecies, as well as paddy and upland rice. Generally, total ROL from entire roots was correlated with metal tolerance and negatively correlated with total As ($R = -0.67$, $P < 0.01$) and inorganic As ($R = -0.47$, $P < 0.05$) in rice grains. There were also significant genotypic effects in percentage of inorganic As ($F = 15.8$, $P < 0.001$) and percentage of DMA ($F = 22.1$, $P < 0.001$) respectively. Root aeration of different genotypes and variation of genotypes on As accumulation and speciation would be useful for selecting genotypes to grow in areas contaminated by As.

Another greenhouse experiment was conducted to examine As dynamics, iron

plaque formation and radial oxygen loss and their relationships with As uptake and speciation in rice. Three genotypes were grown in low As soil amended with different concentrations of As. As concentrations in soil solution were built up during the flooded period, with Arsenic (III) the predominant form. There was a slight increase of iron plaque formation, a decrease in the rates of ROL, and induced a shift in the spatial ROL pattern from the “tight” to the “partial” barrier type. Arsenic speciation results showed that the percentage of DMA increased from 19-28% to 53-58 %, while the percentage of inorganic As decreased from 53-58% to 36-42% with the increasing soil As concentrations; the percentage of organic As was greatly elevated in rice grains (53-70%) compared with small percentages in rice roots (1-2%) and straws (2-6%).

The experiment was conducted to further study whether rice paddy soils would indicate enhanced mobility and bioavailability of this metalloid in soils with impeded drainage. The fate of As was investigated in two types of soils originated from the United Kingdom that were, derived from different sources of industrial waste. Arsenic mobility of soil was measured by soil pore-water sampling under flooded and drained watering regimes. Results showed that the flooding of one test soil greatly enhanced iron (Fe) and manganese (Mn) content of pore-water, followed by increases in As content and poor growth of rice. It is argued that testing As-contaminated soils by flooding combined with chemical analyses and plant assays, provides a realistic indication of future risk associated with waterlogged conditions.

In addition, to elucidate the mechanisms for genotypic differences of rice in As

uptake and tolerance, and that cause rice genotypes with higher ROL to have lower As accumulation, hydroponic experiments were conducted to investigate the effect of internal and external aeration on iron plaque formation, As accumulation and speciation in rice. The present study shows that there were significant correlations between ROL and Fe concentrations in iron plaque ($R = 0.61$, $P < 0.001$) of different genotypes of rice. Iron plaque sequestered more arsenate than arsenite, leading to decreased As accumulation in rice roots. In addition, there were significant correlations between Fe concentrations and As accumulation in iron plaque in both arsenite ($R = 0.87$, $P < 0.001$) and arsenate ($R = 0.81$, $P < 0.001$) treatments. There were significant differences in the amounts of iron plaque formed in different genotypes ($P < 0.01$), in different positions of rice roots ($P < 0.001$) and under different aerated conditions (aerated, normal and stagnant treatments) ($P < 0.001$). Compared with aerated treatment, rice tended to have a higher ROL in stagnant solution, with the highest in the root tip ($P < 0.001$), in accordance with the trend of spatial ROL. The majority As species detected in rice roots and shoots was arsenite ranging from 34-78% of total As in different treatments and genotypes.

Moreover, the effects of root oxidization and P fertilization were conducted to investigate As mobility in soils, As uptake, translocation and speciation in rice. Results showed that root oxidization could significantly influence As mobility in rhizosphere. Genotype TD71 with higher ROL could induce more Fe plaque formation and sequester more As and P in iron plaque and rhizosphere soil,

finally reduce As accumulation in rice plants. Additionally, P mobilized As in soil solution, and increased As accumulation in rice plants. Arsenic speciation results showed that the majority of As species in husks detected was inorganic As, accounting for 82-93% of the total As, while in rice husks and rice grains the majority of As was inorganic As and DMA, accounting for 33-64% of the total As. It was also noted that the percentage of inorganic As decreased, while percentage of DMA increased along the concentration gradient of As and P. The study elucidated the mechanisms involved in ROL on As tolerance and accumulation in rice.

The present study provides valuable information regarding As uptake, accumulation and speciation in rice related with ROL, iron plaque formation and soil chemistry. It provides potential strategies to mitigate the health risk posed by As contamination in rice. However, the mechanism for As tolerance and accumulation, as well as field application for mitigating As accumulation in rice grains need further investigation.

Table of Contents

Declaration	
Abstract	
Acknowledgement	
Table of Contents	
List of Figures	
List of Tables	
Chapter 1 General Introduction	1
1.1 Background of Research	1
1.1.1 Arsenic contamination in paddy rice and paddy field	2
1.2 Literature Review	7
1.2.1 Aerenchyma, porosity, radial oxygen loss (ROL) and iron plaque..	7
1.2.2 Arsenic (As), iron (Fe) and phosphate (P) in soil and in rhizosphere of rice	11
1.2.3 Arsenic in soil	13
1.2.4 Metals and wetland plants.....	13
1.3 Objective of research	16
1.4 Contributions and Significance of the Present Research	19
1.5 Frame work of the PhD thesis.....	19
Chapter 2 Arsenic accumulation and speciation in rice related to root aeration and variation of genotypes.....	21
2.1 Introduction.....	23
2.2 Materials and Method	25
2.2.1 Hydroponic experiment	25
2.2.2 Pot experiment	30
2.2.3 Data analyses	34
2.3 Results.....	35
2.3.1 Porosity and ROL of different genotypes	35
2.3.2 Plant growth during the pot experiment	36
2.3.3 Effects of ROL on arsenic accumulation in different genotypes of rice	40
2.3.4 As speciation in different genotypes	43
2.4 Discussion	49
2.5 Conclusion	53
Chapter 3 Arsenic accumulation and speciation in rice plants related to ROL and iron plaque formation.....	55
3.1 Introduction.....	55
3.2 Materials and Methods.....	59
3.2.1 Plant culture	59
3.2.2 As determination and speciation in soil solution	59
3.2.3 Iron plaque extraction and scanning electron microscopy with	

X-ray analysis (SEM X-ray analysis)	59
3.2.4 Measurement of ROL	60
3.2.5 Plant analysis	62
3.2.6 Statistical analyses	64
3.3 Results.....	65
3.3.1 As in soil solution	65
3.3.2 Effects of As on growth, iron plaque formation, ROL of entire roots and ROL spatial patterns in three genotypes of rice	66
3.3.3. Arsenic uptake and speciation in rice plants	73
3.4 Discussion	76
3.5 Conclusion	80
Chapter 4 Effects of soil chemistry on arsenic uptake of rice genotypes with different ROL associated with macrophyte duckweed	81
4.1 Introduction.....	81
4.2 Materials and Methods.....	83
4.2.1 Study sites and sources of arsenic.....	83
4.2.2 Collection and preparation of soil samples	84
4.2.3 Plant investigation.....	85
4.2.4 Pore-water sampling	86
4.2.5 Analytical methods.....	88
4.2.6 Harvesting and digestion of rice and Lemna tissues.....	89
4.2.7 Statistical analysis	90
4.3 Results.....	91
4.3.1 Soils and As fractionation	91
4.3.2 Trace elements in soil pore-water	91
4.3.3 Rice yield and As uptake.....	92
4.3.4 Arsenic uptake in Lemna minor	94
4.4 Discussion	99
4.5 Conclusion	105
Chapter 5 Do internal and external aeration affect iron plaque formation and As accumulation and speciation in rice?	106
5.1 Introduction.....	106
5.2 Materials and Methods.....	109
5.2.1 ROL of entire roots and iron plaque formation.....	109
5.2.3 Spatial pattern of ROL and spatial distribution of iron plaque	112
5.2.4 Arsenic accumulation and speciation in rice plants	114
5.2.4 Statistical analyses	116
5.3 Results and Discussion	116
5.3.1 Plant growth and iron plaque formation	116
5.3.2 Spatial pattern of ROL and spatial distribution of iron plaque	117
5.3.3 As accumulation and speciation in rice plants	121
5.4 Discussion	128
5.5 Conclusions.....	131
Chapter 6 Effects of root oxidation ability and P on As mobility and	

bioavailability in rice	132
6.1 Introduction.....	132
6.2 Materials and Methods.....	136
6.2.1 Plant culture	136
6.2.2 As determination in soil solutions.....	137
6.2.3 DCB-extraction of iron plaque.....	137
6.2.4 Plant analysis for total As	138
6.2.5 Plant analysis for As speciation	138
6.2.6 Sequential extraction of arsenic fractionations in soils.....	139
6.2.7 Statistical analyses	140
6.3 Results.....	140
6.3.1 As in soil solution and soil As sequestration.....	140
6.3.2 Plant growth.....	144
6.3.3 P and Fe sequestration in iron plaque	144
6.3.4 P accumulation in rice plants	147
6.3.5 As accumulation and speciation in rice plants	147
6.4 Discussion.....	150
6.5 Conclusion	154
Chapter 7 General Discussion and Conclusions	155
7.1 Risk assessment posed by As contamination in rice plants	155
7.2 Effects of radial oxygen loss on As uptake, tolerance and distribution in rice	157
7.3 Effects of genotype variation on As uptake and speciation in rice	158
7.4 Effects of genetic and environmental factors on As Speciation in Rice	159
7.5 Mechanisms evolved in As tolerance and accumulation in rice	161
7.6 Strategies to mitigate As contamination in paddy soil and rice	162
7.7 Future Work	165
7.7.1 Mechanisms of As tolerance, accumulation and speciation in rice	165
7.7.2 Rice breeding and field trials: Rice genotypes with high ROL grown on As-contaminated soils.....	167
References.....	169
Publications.....	188
Curriculum Vitae.....	189