

MASTER'S THESIS

Anti-tumour and anti-angiogenic effects of euxanthone

Tsang, Hing Yan

Date of Award:
2001

[Link to publication](#)

General rights

Copyright and intellectual property rights for the publications made accessible in HKBU Scholars are retained by the authors and/or other copyright owners. In addition to the restrictions prescribed by the Copyright Ordinance of Hong Kong, all users and readers must also observe the following terms of use:

- Users may download and print one copy of any publication from HKBU Scholars for the purpose of private study or research
- Users cannot further distribute the material or use it for any profit-making activity or commercial gain
- To share publications in HKBU Scholars with others, users are welcome to freely distribute the permanent URL assigned to the publication

Anti-tumour and Anti-angiogenic Effects of Euxanthone

TSANG Hing Yan

A thesis submitted in partial fulfillment of the requirements

for the degree of

Master of Philosophy

November, 2001

Hong Kong Baptist University

Abstract

Plants of *Polygala sp.* have long been used as folk medicines to improve the learning and memory process. The roots of *Polygala* have also been used as sedative agent and anti-tumor drug in some Chinese medicines. Both naturally occurring and synthetic xanthone derivatives have been reported to mediate anti-tumor activities on human cells *in vitro*. In the current study, the potential anti-tumor and anti-angiogenic effects of euxanthone isolated from the medicinal herb *Polygala caudata* were investigated. Murine neuroblastoma (Neuro 2A, subclone BU-1) was used as a model to study the anti-tumor activities, and the human umbilical vein endothelial cells (HUVECs) was employed as a model to evaluate the anti-angiogenic activities of euxanthone.

In the studies of anti-tumor activity, euxanthone (50 - 100 μ M) was found not only to inhibit the growth and arrest BU-1 at G2/M phase, but also to induce the morphological differentiation of the cells. The morphological differentiation of BU-1 cell was associated with the outgrowth of neurite and the enlargement of cell bodies. Previous studies suggested that protein kinase C (PKC) signaling pathway may be involved in the neuronal differentiation of Neuro-2A cells. Thus, the effects of PKC inhibitor on the growth and neuronal differentiation of euxanthone-treated BU-1 cells were also examined. Significant reduction of euxanthone-induced neuritogenic effect was observed when the conventional PKC isoform specific inhibitor Gö6976 was included in the culture. The results suggest that the conventional PKC isoforms may be involved in euxanthone-induced neuritogenesis in the BU-1 cells. Axonal (NF-H and tau) and dendritic (MAP-2) markers were employed to characterize the nature of

induced neuritic outgrowth in euxanthone-treated BU-1. Laser scanning confocal microscopy was used to study the distribution of these neuronal markers. A mixed phenotype of axon- and dendrite-like processes was observed 5 days after euxanthone treatment. These data suggested that euxanthone might be one of the anti-tumor compounds in the medicinal plant *Polygala caudata*.

In the second part of the study, the anti-angiogenic effect of euxanthone on HUVECs was examined. Euxanthone was found to inhibit the *in vitro* migration and proliferation of HUVECs. The growth inhibition of HUVECs was also supported by the results from the flow cytometric analysis. The proportion of HUVECs in S-phase was reduced after euxanthone treatment. The induction of morphogenesis of HUVECs, namely changing from the typical cobblestone appearance into elongated spindle shape, was observed after euxanthone treatment. Different *in vitro* and *in vivo* assays (*in vitro* tube formation assay and *in vivo* chick chorioallantoic membrane study) were evaluated on the effect of euxanthone on the vessel formation. The *in vitro* 3-D tube formation assay was finally selected in this study. The anti-angiogenic effect was also observed in the *in vitro* tube formation on Matrigel. It has been previously reported that cell adhesion molecules including platelet endothelial cell adhesion molecules-1 (PECAM-1) and cadherin-5 are critical molecules expressed on HUVECs for tube formation. In this study, PECAM-1 and cadherin-5 were found to express mainly at the cell junction of HUVECs. Euxanthone did not alter the expression level of these two adhesion molecules as revealed by the flow cytometric analysis. Inhibition of HUVEC migration might be responsible for the anti-angiogenic effect of euxanthone. The data generated in the study allow us to further

investigate the potential anti-angiogenic effects of euxanthone in other *in vivo* angiogenic models.

Table of Contents

Declaration	i
Abstract	ii
Acknowledgments	v
Table of Contents	vi
List of Abbreviations	xi
List of Tables	xiv
List of Figures	xv
Chapter I Introduction	
1.1. The Nervous System	1
1.1.1. Architecture and cytoskeletons of neurons	1
1.1.1.1. Structural proteins of cytoskeleton in neurons	2
1.1.1.1.1. Microtubules - an important determinant of cell architecture	2
1.1.1.1.2. Involvement of microfilaments in intracellular transport and cell movement	6
1.1.1.1.3. Intermediate filaments - the prominent constituents of nervous system	9
1.1.2. Neuroblastoma cells	11
1.1.3. Pathology and clinical features of neuroblastoma	11
1.1.3.1. Origin of neuroblastoma	11
1.1.3.2. Histology of neuroblastoma	12
1.1.3.3. Clinical features of neuroblastoma	13
1.1.4. Diagnosis and investigations of neuroblastoma	14
1.1.4.1. Tumor markers	14
1.1.4.2. Serum markers	15
1.1.4.3. Tumor cell DNA content	15
1.1.4.4. N-myc oncogene activation	16
1.1.4.5. Gene expression	16
1.1.5. Clinical staging of neuroblastoma	17
1.1.6. Treatment of neuroblastoma	20
1.1.7. Murine neuroblastoma, Neuro-2A	21
1.1.8. Protein kinase C (PKC) signaling pathway	22
1.2. The Chinese medicines	23

1.2.1. Medicinal herbs as drugs and diet supplements	24
1.2.2. Xanthones	25
1.2.3. Nature of euxanthone	25
1.3. The angiogenesis	26
1.3.1. Angiogenesis process	27
1.3.2. Regulation of angiogenesis and angiogenic diseases	27
1.3.3. Clinical applications of angiogenesis research	30
1.3.3.1. Anti-angiogenic therapy: a novel weapon to fight cancer	30
1.3.3.1.1. Role of angiogenesis in tumor development	30
1.3.3.1.2. Therapeutic strategies of anti-angiogenesis	31
1.3.3.1.3. Challenges to anti-angiogenic therapy	35
1.3.3.1.4. Combination of anti-angiogenic and other kinds of therapies	36
1.3.3.2. Therapeutic angiogenesis	37
1.3.4. Endothelial cell adhesion molecules	38
1.3.4.1. Platelet endothelial cell adhesion molecule-1 (PECAM-1/CD31)	39
1.3.4.2. Cadherin	39
1.3.5. Review on the assays for angiogenesis used in the study	40
1.3.5.1. <i>In vitro</i> assays for angiogenesis	40
1.3.5.1.1. Endothelial cell migration assay	41
1.3.5.1.2. Endothelial cell proliferation assay	42
1.3.5.1.3. Three-dimensional differentiation: <i>in vitro</i> capillary tube formation	42
1.3.5.2. <i>In vivo</i> models for angiogenesis	44
1.3.5.2.1. Chick embryo chorioallantoic membrane (CAM) assay	44
1.4. Aims of Study	45
Chapter II Materials and Methods	
2.1. Materials	47
2.1.1. Cell culture	47
2.1.1.1. Culture media for neuroblastoma BU-1 cells	47
2.1.1.2. Culture media for human umbilical vein endothelial cells (HUVECs)	47
2.1.2. Effects of euxanthone on neuroblastoma	47
2.1.2.1. Reagents to measure of morphological differentiation	48
2.1.2.2. Reagents for MTT cytotoxicity assay	48
2.1.2.3. Reagents for cell cycle analysis	48
2.1.2.4. Reagents for [³ H]-thymidine incorporation assay	48
2.1.2.5. Immunocytochemical staining of axonal and dendritic markers	48

2.1.3. Anti-angiogenic effect of euxanthone	49
2.1.3.1. Cell migration assay	49
2.1.3.2. Trypan blue exclusion method	49
2.1.3.3. Chick embryo chorioallantoic membrane (CAM) assay	49
2.1.3.4. <i>In vitro</i> tube formation assay	49
2.1.3.5. Immunostaining and flow cytometric analysis of PECAM-1 and cadherin-5	49
2.2. Methods	50
2.2.1. Cell culture	50
2.2.1.1. Neuroblastoma BU-1 cells	50
2.2.1.2. Human umbilical vein endothelial cells (HUVECs)	50
2.2.2. Effects of euxanthone on neuroblastoma	51
2.2.2.1. MTT cytotoxicity assay	51
2.2.2.2. Measurement of morphological differentiation	52
2.2.2.3. [³ H]-thymidine incorporation assay	53
2.2.2.4. Cell cycle analysis	53
2.2.2.5. Immunocytochemical staining of axonal and dendritic markers	54
2.2.2.5.1. Immunostaining of BU-1 cells	54
2.2.2.5.2. Laser scanning confocal microscopy	55
2.2.3. Anti-angiogenic effect of euxanthone	55
2.2.3.1. Cell migration assay	55
2.2.3.2. Measurement of cell proliferation	56
2.2.3.2.1. Trypan blue exclusion method	56
2.2.3.2.2. [³ H]-thymidine incorporation assay	58
2.2.3.3. Cell cycle analysis	58
2.2.3.4. Chick embryo chorioallantoic membrane (CAM) assay	58
2.2.3.5. Morphological change of euxanthone-treated HUVECs	59
2.2.3.6. <i>In vitro</i> tube formation assay	59
2.2.3.7. Immunofluorescent staining of PECAM-1 and cadherin-5	60
2.2.3.8. Determination of PECAM-1 and cadherin-5 expression by FACS	61

Chapter III Effect of Euxanthone on Neuroblastoma (BU-1 cells)

3.1. Introduction	63
3.2. Results	64
3.2.1. Cytotoxicity of euxanthone	64
3.2.2. Induction of morphological differentiation	64
3.2.2.1. Morphological differentiation of neuroblastoma cells	64
3.2.2.2. Phenotypes of differentiated neuroblastoma cells	65
3.2.3. Anti-proliferative activity of euxanthone on BU-1 cells	66

Chapter V General Discussion

5.1. Induction of neuronal differentiation and the anti-tumor activities	121
5.2. Involvement of PKC signaling in neuronal differentiation	122
5.3. Characterization of neuritic outgrowth	124
5.4. Controlling of tumor growth using anti-angiogenic drugs	125
5.5. Conclusion	127
5.6. Further studies	128
References	130
Lists of Publications	158
Cirriculum Vitae	159