

## DOCTORAL THESIS

### Interactions of arbuscular mycorrhizal fungi with an arsenic hyperaccumulator plant (*pteris vittata*) on the uptake of arsenic

Leung, Ho Man Homan

*Date of Award:*  
2008

[Link to publication](#)

#### General rights

Copyright and intellectual property rights for the publications made accessible in HKBU Scholars are retained by the authors and/or other copyright owners. In addition to the restrictions prescribed by the Copyright Ordinance of Hong Kong, all users and readers must also observe the following terms of use:

- Users may download and print one copy of any publication from HKBU Scholars for the purpose of private study or research
- Users cannot further distribute the material or use it for any profit-making activity or commercial gain
- To share publications in HKBU Scholars with others, users are welcome to freely distribute the permanent URL assigned to the publication

**Interactions of Arbuscular Mycorrhizal Fungi with an Arsenic  
Hyperaccumulator Plant (*Pteris vittata*) on the Uptake of Arsenic**

**LEUNG Ho Man, Homan**

**A thesis submitted in partial fulfillment of the requirements  
for the degree of  
Doctor of Philosophy**

**Principal Supervisor: Prof. WONG Ming Hung**

**Hong Kong Baptist University**

**July 2008**

## ABSTRACT

The major objective of the project is to study the roles and effects of arbuscular mycorrhiza (AM) fungi on the growth, metal uptake and distribution in an arsenic (As) hyperaccumulator (*Pteris vittata*) and a non-hyperaccumulator (*Cynodon dactylon*) when grown in metal contaminated soil.

A field survey of metal concentrations and AM components of plants growing on five mining sites was conducted in Chenzhou City, Hunan Province, Southern China and a control site in Hong Kong. Significant differences were observed on the average total concentrations of four heavy metals [lead (Pb), zinc (Zn), copper (Cu), cadmium (Cd)] and one metalloid [arsenic (As)] in contaminated soils compared with the control site. Three components of mycorrhizal colonization (arbuscules, vesicles and coiled hyphae) were found in the roots of *C. dactylon* and *P. vittata* growing at Do Shun Long (DSL) mine site. Concentrations of As in fronds were 24-fold higher than in roots of *P. vittata* with the highest mycorrhizal colonization rate (73%) among all sampling sites. Extensive mycorrhizal colonization (85%) was also recorded in the roots of *C. dactylon* with As accumulation 57 times higher than in shoots.

A greenhouse trial was conducted to investigate the role of indigenous mycorrhiza (IM) isolated from As mine in aiding As uptake and tolerance by *P. vittata* (As hyperaccumulator) and *C. dactylon* (a multi-metal root accumulator). The infectious percentage of mycorrhiza (0mg/kg As: 26.4%, 50 mg/kg As: 30.3%, 100mg/kg As: 40.6%)

and the average biomass of shoots in infected *P. vittata* increased (0mg/kg As: 2.45g/pot, 50mg/kg As: 2.48g/pot, 100mg/kg As: 10.9g/pot) according to the increase of As levels when compared to control. IM also enhanced As accumulation (0mg/kg As: 3.70mg/kg, 50mg/kg As: 58.3mg/kg; 100mg/kg As: 88.1mg/kg) in the As mine populations of *P. vittata* and also sustained its growth by aiding P absorption. For *C. dactylon*, As was mainly accumulated in mycorrhizal roots and translocation to shoots was inhibited.

The effects of three types of single inoculum [IM, *Glomus mosseae* (GM) and *Glomus intraradices* (GI)] and two types of mixed inoculum (mixed with IM and either GM or GI) on the growth response of *P. vittata* and *C. dactylon* were studied. Mixed inoculum IM/GM promoted substantially higher mycorrhizal colonization and arsenate reductase (AR) activity in *P. vittata* than *C. dactylon* among all As levels. This may be attributed to IM/GM inoculum which could improve soil conditions and eventually favour AR formation and synthesis in plant root. The portion of *Paris* AM structure (observed in colonized roots) together with the highest As translocation factor of 10.2 in *P. vittata* inoculated with IM/GM was also noted.

The contributions of AM fungi and phosphate (P) amendment on the As uptake and accumulation of *P. vittata* and *C. dactylon* in As contaminated soil was investigated. It was found that the growth of mycorrhizal plants treated with P amendment was significantly improved when compared with non-amended plants. In addition, plant biomass, root surface area and chlorophyll concentrations in *P. vittata* inoculated with IM/GM were significantly higher when treated with P rock in 300 mg/kg As contaminated soil than the

other treatments. Henceforth, such combination seemed to play a potential role in enhancing phytoextraction efficiency of hyperaccumulating plants growing in As contaminated soil.

The effects of AM fungi and P rock on the phytoremediation efficiency of *P. vittata* and *C. dactylon* growing on As contaminated field soil were studied. Significant As reduction in soil, increased plant biomass and As accumulation were found for *P. vittata* treated with IM/GM and P rock at 100% water holding capacity (WHC) level. The overall results indicated that the synergistic effect of mycorrhiza and P rock affected As subcellular distribution of the hyperaccumulator and thereby altered its As removal efficiency under well-watered conditions.

The present study has proved that IM/GM combined with P rock would enhance the host plant in As accumulation when grown in As contaminated substrates. Therefore, mycorrhizal inoculation could be combined with other practices such as appropriate soil amendments to maximize re-vegetation success at field.

## TABLE OF CONTENTS

<b>DECLARATION</b> .....	i	
<b>ABSTRACT</b> .....	ii	
<b>ACKNOWLEDEMENTS</b> .....	v	
<b>TABLE OF CONTENTS</b> .....	vii	
<b>LIST OF FIGURES</b> .....	xi	
<b>LIST OF TABLES</b> .....	xiii	
<b>LIST OF PHOTOGRAPHS</b> .....	xvi	
<b>LIST OF APPENDICES</b> .....	xvii	
<b>LIST OF ABBREVIATIONS</b> .....	xviii	
<b>CHAPTER 1 GENERAL INTRODUCTION</b>		
<b>1.1 Background of Research</b>		
1.1.1 Current Issues of Metal Contamination in Soil.....	1	
1.1.2 Technologies for Treatment of Metal Contaminated Soil.....	5	
1.1.3 Current Progress of Phytoremediation.....	8	
1.1.4 Advantages of Using Higher Plants to Remediate Polluted Soil.....	11	
1.1.5 Mechanisms of Metal Uptake and Accumulation in Plant.....	11	
<b>1.2 Mycorrhiza: A Helping Agent for Contaminated Land Remediation</b>		
1.2.1 Mycorrhiza: Overview.....	17	
1.2.2 Roles of Mycorrhiza on Metal Contaminated Land.....	18	
1.2.3 Contradictory Role of AM for Phytoremediation.....	24	
<b>1.3 Objectives of Research</b> .....		31
<b>1.4 Contributions and Significance of the Present Research</b> .....		33
<b>1.5 Framework of Research</b> .....		33
<b>CHAPTER 2 ACCUMULATION OF METALS and METALLOIDS BY PLANTS THRIVING ON METAL MINES WITH HEAVY METALS CONTAMINATION</b>		
<b>2.1 Introduction</b> .....		35
<b>2.2 Materials and methods</b>		
2.2.1 Study Sites.....	37	
2.2.2 Sampling area.....	38	
2.2.3 Preparation of Soil and Plant Samples for Chemical Analysis.....	38	
2.2.4 Mycorrhizal Colonization in Plants.....	41	
2.2.5 Statistical Analysis.....	42	
<b>2.3 Results</b>		
2.3.1 Physico-chemical Characteristics of Sampling Sites.....	42	
2.3.2 Vegetation on Mine Tailings, Similarity Index of Plants in Six Sites .....	44	
2.3.3 Mycorrhizal Structures of Plants Commonly Grown in Mining Sites.....	44	
2.3.4 Plant Metal Concentrations and Metal TFs.....	49	
<b>2.4 Discussion</b>		
2.4.1 Common Soil Characteristics at the Five Mine Sites.....	53	
2.4.2 Incidence of AM Fungi.....	54	

2.4.3 Survival Strategies of Different Plant Species at Mine Sites and Their Potential for Phytoremediation	
2.4.3.1 <i>C. moritolum</i> and <i>M. florodulus</i> (Metal Indicators).....	55
2.4.3.2 <i>C. dactylon</i> (Metal Excluder).....	56
2.4.3.3 <i>P. vittata</i> (Metal Accumulator).....	57
2.5 Conclusions.....	58

### **CHAPTER 3 INTERACTIONS OF MYCORRHIZAL FUNGI WITH *PTERIS VITTATA* (AS HYPERACCUMULATOR) IN AS-CONTAMINATED SOILS**

<b>3.1 Introduction</b> .....	60
<b>3.2 Materials and Methods</b>	
3.2.1 Field Sampling and Treatment.....	63
3.2.2 Mycorrhizal Infection.....	64
3.2.3 Trap Culture Establishment.....	64
3.2.4 Cultivation System.....	65
3.2.5 Experimental Set Up.....	65
3.2.6 Chemical Analyses.....	66
3.2.7 Statistical Analysis.....	68
<b>3.3 Results</b> .....	68
<b>3.4 Discussion</b>	
3.4.1 The Performance of <i>P. vittata</i> With and Without Mycorrhizal Inoculations.	73
3.4.2 The Role of P Absorbed by Mycorrhiza in Improving As Uptake by <i>P. vittata</i> .....	77
3.4.3 Uniqueness of Indigenous Mycorrhiza Found at As Contaminated Site.....	78
<b>3.5 Conclusion</b> .....	79

### **CHAPTER 4 THE ROLE OF MIXED ARBUSCULAR MYCORRHIZAL (AM) INOCULUM ON THE GROWTH OF *PTERIS VITTATA* (AS HYPERACCUMULATOR) AND *CYNODON DACTYLON* IN AS CONTAMINATED SOILS**

<b>4.1 Introduction</b> .....	80
<b>4.2 Materials and Methods</b>	
4.2.1 Field Sampling and Sample Treatment.....	83
4.2.2 Trap Culture for AM Fungi Inoculum.....	83
4.2.3 Experimental Set Up.....	83
4.2.4 Chemical Analyses.....	85
4.2.5 Determination of Arsenate Reductase (AR) Activities in Plants.....	86
4.2.6 Statistical Analysis.....	87
<b>4.3 Results</b> .....	87
<b>4.4 Discussion</b>	
4.4.1 Contribution of IM/GM Inoculum to the Uptake of As by Plants.....	97

4.4.2 The Importance of Hyphae and <i>Paris</i> -form AM Formed in Plants and Their Significance to Phytoremediation.....	98
4.4.3 Uniqueness of Arsenate Reductase (AR) Activity in <i>P. vittata</i> .....	99
4.4.4 Potential Application of Mixed AM Inoculum for Soil Remediation at Mining Site.....	101
<b>4.5 Conclusion</b> .....	102

**CHAPTER 5 THE ROLES OF ARBUSCULAR MYCORRHIZAL FUNGI AND PHOSPHATE AMENDMENT ON AS UPTAKE AND ACCUMULATION OF *PTERIS VITTATA* IN AS CONTAMINATED SOIL**

<b>5.1 Introduction</b> .....	103
<b>5.2 Materials and Methods</b>	
5.2.1 Field Sampling and Sample Treatment.....	106
5.2.2 Trap Culture for AM Fungi Inoculum.....	106
5.2.3 Experimental Set-up.....	106
5.2.4 Plant Analysis.....	108
5.2.5 Statistical Analysis.....	110
<b>5.3 Results</b> .....	111
<b>5.4 Discussion</b>	
5.4.1 Importance of Adding Soil Amendments for Phytoremediation.....	119
5.4.2 Comparison Between Natural and Synthetic P Amendments in Treating As-contaminated Soil.....	122
5.4.3 Contributions of AM Fungi to the Host for As Uptake and Accumulation in Amended Soil.....	124
<b>5.5 Conclusion</b> .....	128

**CHAPTER 6 FEASIBLE BIOREMEDIATION THROUGH ARBUSCULAR MYCORRHIZAL FUNGI AND PHOSPHATE ROCK FACILITATING THE HEAVY METAL UPTAKE AND ACCUMULATION OF HYPERACCUMULATORS AND NON-HYPERACCUMULATORS**

<b>6.1 Introduction</b> .....	129
<b>6.2 Materials and Methods</b>	
6.2.1 Plant Selection.....	131
6.2.2 Suitability of <i>Glomus mosseae</i> as a Mixed AM Inoculum.....	131
6.2.3 Soil Moisture Effects on the Uptake of As by <i>P. vittata</i> and <i>C. dactylon</i> ...	132
<b>6.3 Results</b> .....	138
<b>6.4 Discussion</b>	
6.4.1 Potential of Using Mycorrhiza to Aid the Host in Phytoremediation.....	152
6.4.2 Significance of As Species in Contaminated Soil for the Clean-up of As By Mycorrhizal Hyperaccumulators.....	154
6.4.3 Strategies of Mycorrhizal Plants for As Uptake and Accumulation .....	156



<b>6.5 Conclusion.....</b>	<b>159</b>
<b>CHAPTER 7 GENERAL DISCUSSION AND CONCLUSION</b>	
<b>7.1 Introduction.....</b>	<b>160</b>
<b>7.2 Potential of Mycorrhizal Fungi in Phytoremediation of Arsenic.....</b>	<b>161</b>
<b>7.3 Contribution of Mixed AM Inoculum to its Host for As Phytoremediation..</b>	<b>162</b>
<b>7.4 Effect of Soil Amendment on Arsenic Uptake and Accumulation of the Host Plants.....</b>	<b>163</b>
<b>7.5 Possible Ways for the Interaction of Mycorrhizal Fungi and their Host Plants with Arsenic.....</b>	<b>163</b>
<b>7.6 General Conclusion.....</b>	<b>164</b>
<b>7.7 Further Study.....</b>	<b>165</b>
<b>REFERENCES.....</b>	<b>167</b>
<b>APPENDICES.....</b>	<b>219</b>
<b>PUBLICATIONS.....</b>	<b>223</b>
<b>CURRICULUM VITAE.....</b>	<b>224</b>