

MASTER'S THESIS

The study of novel dioxin antagonist-euxanthone and its derivatives

Zhang, Qi

Date of Award:
2003

[Link to publication](#)

General rights

Copyright and intellectual property rights for the publications made accessible in HKBU Scholars are retained by the authors and/or other copyright owners. In addition to the restrictions prescribed by the Copyright Ordinance of Hong Kong, all users and readers must also observe the following terms of use:

- Users may download and print one copy of any publication from HKBU Scholars for the purpose of private study or research
- Users cannot further distribute the material or use it for any profit-making activity or commercial gain
- To share publications in HKBU Scholars with others, users are welcome to freely distribute the permanent URL assigned to the publication

**The Study of Novel Dioxin
Antagonist-Euxanthone and its Derivatives**

ZHANG Qi

**A thesis submitted in partial fulfillment of the requirements
for the degree of
Master of Philosophy**

**Principal Supervisor: Dr. Ricky N.S. WONG
Hong Kong Baptist University
December 2003**

Abstract

2,3,7,8-Tetrachlorodibenzo-*p*-dioxin (TCDD), a widespread environmental pollutant, has been shown to act as an agonist of the aryl hydrocarbon receptor (Ah receptor) subsequently leading to adverse effects. While much work has been done on the toxicity mechanism of TCDD, some attention begins to turn to discerning the mechanism by which phytochemicals are utilized to inhibit TCDD toxicity, such as carcinogenesis and hepatotoxicity.

The present study disclosed a novel antagonist, named PW-1, which was extracted from Chinese medicine herb *Polygala caudata*. The antagonist showed significant suppression of TCDD-induced EROD activity at a dose-dependent manner as well as *CYP1A1* gene expression in HepG2 cells. Furthermore, six synthetic derivatives of PW-1, namely CYL-1, -2, -3, -4, -5, and -6, also showed to possess inhibitory effects on TCDD-induced EROD activity in a dose-response manner. The PW-1 and all its derivatives *per se* did not significantly result in induction of EROD activity as compared to TCDD. However, CYL-4, PW-1, CYL-1, CYL-2, and CYL-3 at the concentration of 1.65 μM , 1.95 μM , 2.5 μM , 5.0 μM and 8.2 μM , respectively, can give rise to a 50% inhibition (IC_{50}) of EROD activity generated by induction of 0.5 nM TCDD. Almost complete inhibition was observed by CYL-4, PW-1 and CYL-1 at the concentration of 12.5 μM . Real-time PCR as well as agarose gel electrophoresis were employed to evaluate the validity of the *CYP1A1* gene expression changes induced by TCDD and the antagonistic effects of PW-1 on TCDD.

On the other hand, cDNA microarray technology was employed to uncover the possible antagonistic mechanism of PW-1 against TCDD. For this, approximately two thousands of liver-specific cDNAs were firstly prepared from a human liver lambda cDNA library with the titer of 2.97×10^{11} pfu/ml. These cDNAs were then printed on polylysine-coated slides by an arraying robot. The microarrays thus made were hybridized simultaneously with three probes which were labeled separately with a type of fluorescent dye, namely Alexa Fluor 546 for TCDD treatment, Alexa Fluor 488 for TCDD plus PW-1, and Alexa Fluor 647 for solvent control. Finally, the hybridization signals were scanned by the laser confocal scanner, and the digitized fluorescence signals were normalized by an approach of total measured fluorescence intensity. As a result, fifty candidate genes were chosen for sequencing. By blasting the obtained sequences in the GenBank database, 21 genes were identified as unique to the GeneBank. Among them, 18 genes were more than 2.0 fold, 2 genes were without change, and only one gene was less than 1-fold change. Among these 18 genes, seven were firstly disclosed to be induced by TCDD. Obviously, the expressions of most genes identified were altered by exposure to TCDD, and most of genes up-regulated by TCDD were down-regulated by PW-1, such as the gene encoding Alpha-1-microglobulin/bikunin precursor (AMBIP); the gene encoding Retinoid X receptor, alpha (RXRA); the gene encoding complement component 3 (C3). An antagonistic mechanism of PW-1 against TCDD was deduced in the present study. Especially, an interleukin-6 (IL-6)-mediated regulation pathway of TCDD function was also proposed.

Table of Contents

| | Page |
|---|-------------|
| Declaration | i |
| Abstract | ii |
| Acknowledgements | iv |
| Table of Contents | v |
| List of Tables | ix |
| List of Figures | x |
| List of Abbreviations | xii |
| | |
| Chapter 1 Introduction | |
| | |
| 1.1 General Introduction | 1 |
| | |
| 1.2 Dioxin | 2 |
| 1.2.1 Nature of Dioxin | 2 |
| 1.2.2 Action Mechanism of Dioxin | 3 |
| | |
| 1.3 Biological Effects of Dioxin | 4 |
| 1.3.1 <i>In Vivo</i> Biological Effects | 7 |
| 1.3.1.1 In Human | 7 |
| 1.3.1.2 In Laboratory Animals | 8 |
| 1.3.1.3 Apoptosis | 9 |
| 1.3.1.4 Oxidative Stress | 10 |
| 1.3.2 <i>In Vitro</i> Biological Effects | 11 |
| | |
| 1.4 Bio-analytical Techniques for Analyzing Dioxin | 12 |
| 1.4.1 Biomarker | 12 |
| 1.4.1.1 EROD | 13 |
| 1.4.1.2 AHH | 14 |
| 1.4.1.3 Chemical Marker-Vitamin A | 14 |
| 1.4.1.4 CYP1A1 mRNA | 15 |
| 1.4.2 DNA Micoarray | 16 |
| 1.4.3 Bioassay (Reporter Gene Expression Assay) | 18 |
| 1.4.3.1 CALUX | 18 |
| 1.4.3.2 GRAB | 19 |
| 1.4.3.3 EGFP | 21 |
| | |
| 1.5 Antagonism of Dioxin | 22 |
| 1.5.1 Antagonist | 22 |

| | | |
|--------------|--|----|
| 1.5.2 | Action Mechanism of Antagonism | 23 |
| 1.6 | Euxanthone (PW-1) and its Synthetic Derivatives | 25 |

Chapter 2 Identification of Euxanthone (PW-1) as a Novel TCDD Antagonist

| | | |
|--------------|---|----|
| 2.1 | Introduction | 28 |
| 2.2 | Materials and Methods | 30 |
| 2.2.1 | Materials | 30 |
| 2.2.1.1 | Culture Media for HepG2 Cells | 30 |
| 2.2.1.2 | Reagents for MTT Cytotoxicity Assay | 30 |
| 2.2.1.3 | Reagents for Ethoxyresorufin- <i>O</i> -deethylase (EROD) Assay | 31 |
| 2.2.1.4 | CYP1A1 Gene Expression Assay | 31 |
| 2.2.2 | Methods | 31 |
| 2.2.2.1 | Cell Culture and Treatment | 31 |
| 2.2.2.2 | Ethoxyresorufin- <i>O</i> -dealkylase (EROD) Assay | 32 |
| 2.2.2.3 | MTT Cytotoxicity Assay | 32 |
| 2.2.2.4 | Total RNA Isolation | 33 |
| 2.2.2.5 | Real-time RT-PCR | 33 |
| 2.2.2.5.1 | Preparation of cDNAs | 33 |
| 2.2.2.5.2 | Real-time PCR | 34 |
| 2.3 | Results | 35 |
| 2.3.1 | Effect of TCDD on EROD Activity in HepG2 Cells | 35 |
| 2.3.2 | Effects of PW-1 and Its Synthetic Derivatives on EROD Activity in HepG2 Cells | 36 |
| 2.3.3 | Inhibitory Effects of Euxanthone (PW-1) and Its Synthetic Derivatives on EROD Activity Induced by TCDD in Intact HepG2 Cells | 36 |
| 2.3.4 | Real-Time PCR Quantification of the Effect of Euxanthone (PW-1) on CYP1A1 Expression | 37 |

| | | |
|-------|---|----|
| 2.3.5 | Effect of Alpha-naphthoflavone (ANF) on EROD Activity | 38 |
| 2.3.6 | Effect of 7-ketocholesterol (7-KC) on EROD Activity | 39 |
| 2.3.7 | Cytotoxicity of Euxanthone (PW-1) | 39 |
| 2.4 | Discussion | 40 |

Chapter 3 Determination of Antagonism Mechanism of Euxanthone (PW-1) via Three-color cDNA Microarray Technology

| | | |
|-------------|---|----|
| 3.1 | Introduction | 54 |
| 3.2 | Materials and Methods | 56 |
| 3.2.1 | Materials | 56 |
| 3.2.1.1 | Materials for Preparation of cDNA Array | 56 |
| 3.2.1.2 | Materials for Preparation of cDNA Probe | 57 |
| 3.2.1.3 | Materials for Hybridization | 57 |
| 3.2.2 | Methods | 58 |
| 3.2.2.1 | Preparation of cDNA Microarray | 58 |
| 3.2.2.1.1 | Recovery of cDNA-containing Phagemids From Uni-ZAP XR Vector | 58 |
| 3.2.2.1.1.1 | Preparation of the <i>E. coli</i> Host Strains (Strains XL1-Blue MRF' and SOLR) | 58 |
| 3.2.2.1.1.2 | Titer of the ExAssist™ Interference-Resistant Helper Phage | 58 |
| 3.2.2.1.1.3 | Titer of the Human Liver Lambda cDNA Library Using Blue-White Selection | 59 |
| 3.2.2.1.1.4 | <i>In Vivo</i> Mass Excision of the pBluescript Phagemid From the Uni ZAP XR Vector | 59 |
| 3.2.2.1.1.5 | Phagemid Transformation and Blue-White Color Selection | 60 |
| 3.2.2.1.1.6 | Lager-scale Isolation of pBluescript Phagemid Using Montage™ Plasmid Miniprep ₉₆ Kit | 60 |
| 3.2.2.1.2 | PCR Amplification of cDNA | 61 |
| 3.2.2.1.3 | Agarose Gel Electrophoresis Analysis | 62 |

| | | |
|--------------|---|------------|
| 3.2.2.1.4 | PCR Product Purification | 62 |
| 3.2.2.1.5 | Preparation of the Human Liver Cell cDNA Microarray | 63 |
| 3.2.2.1.5.1 | Rehydration | 63 |
| 3.2.2.1.5.2 | Ultraviolet Cross-linking | 64 |
| 3.2.2.1.5.3 | Blocking | 64 |
| 3.2.2.1.5.4 | Denaturation | 64 |
| 3.2.2.2 | Isolation and Purification of Total RNA | 65 |
| 3.2.2.3 | cDNA Labeling and Hybridization | 65 |
| 3.2.2.4 | cDNA Microarray Assay | 66 |
| 3.2.2.5 | Sequencing of cDNA | 67 |
| 3.2.2.6 | Analysis of cDNA Sequence | 67 |
| 3.3 | Results | 68 |
| 3.3.1 | Amplification of Human Liver Cell cDNA | 68 |
| 3.3.2 | Microarray Image Scanning | 68 |
| 3.3.3 | cDNA Microarray Studies | 72 |
| 3.4 | Discussion | 73 |
| 3.5 | Conclusion | 86 |
| 3.6 | Prospective work | 87 |
| | Appendix-1 | 107 |
| | Appendix-2 | 124 |
| | References | 125 |
| | Curriculum Vitae | 145 |