

DOCTORAL THESIS

Oligosaccharides approach for the qualitative analysis of dendrobium officinale polysaccharides

Wong, Tin Long

Date of Award:
2019

[Link to publication](#)

General rights

Copyright and intellectual property rights for the publications made accessible in HKBU Scholars are retained by the authors and/or other copyright owners. In addition to the restrictions prescribed by the Copyright Ordinance of Hong Kong, all users and readers must also observe the following terms of use:

- Users may download and print one copy of any publication from HKBU Scholars for the purpose of private study or research
- Users cannot further distribute the material or use it for any profit-making activity or commercial gain
- To share publications in HKBU Scholars with others, users are welcome to freely distribute the permanent URL assigned to the publication

ABSTRACT

Numerous of studies have reported that polysaccharides have many bioactivities including immune system modulation, anti-oxidative and anti-tumor activities. Because herbal materials are rich in polysaccharides, some of the herbs like *Dendrobium officinale*, its polysaccharide marker (DOP) has been applied in the quality analysis of polysaccharides in *D. officinale* by high performance gel permeation chromatography (HPGPC). However, such polysaccharide marker is not presented in every herb, and DOP also has limitation in monitoring herb formula. In addition, even though the same herbal materials and the same extraction methods were used, different studies gave out different conclusion regarding the structure of polysaccharides. Therefore, in the current study, *D. officinale* was used as a case study to demonstrate the application of ABEE-labeled oligosaccharides approach on structure elucidation and quantification of polysaccharides in herbs and herb formulae.

First, the elucidation of polysaccharide structure remains challenging due to the lack of accurate analytical methods to determine the sequence and nature of glycosidic linkages. Oligosaccharide fragments from hydrolysis of polysaccharides are believed to provide accurate structure information, however, they are hard-to-separate and hard-to-detect. In the proposed method, the oligosaccharides generated from partial acid hydrolysis of DOP were labeled with p-aminobenzoic ethyl ester (ABEE), which made them separable and detectable by ultra-high performance liquid chromatography-quadrupole time-of-flight mass spectrometry (UHPLC-Q-TOF-MS-DAD). Subsequently, nine ABEE-labeled oligosaccharide fragments (dimer to decamer) were isolated and identified by MS sequencing and 2D-NMR, and were confirmed by methylation analysis. The results indicated that the backbone should be β -D-1,4-linked Man_n chain instead of mixed mannose and glucose chain.

Second, two ABEE-labeled oligosaccharides namely, Te-Man-ABEE and Pen-Man-ABEE, were selected as chemical markers in the quantification of DOP in *D. officinale* and *D. officinale* (DO) products due to their high specificity in herb formula. The linear relationship between the content of these two markers and the content of DOP was successfully established. The linear

relationship was further transformed to that between peak area of chemical markers and DOP content so that chemical markers were not necessary to be isolated for analysis. This linear relationship was systemically validated in terms of repeatability, precision and accuracy. The results showed that these two oligosaccharide markers presented a good linear relationship with DOP ($R^2 \geq 0.997$) in the range of 0.68-16.02 μg and also demonstrated satisfactory repeatability ($\text{RSD} < 7.0\%$), and recovery (91.41% - 118.30%) in real sample determination. There was no significant difference between the results given by the two chemical markers as the RSD values were not more than 7.0%. While concerning the results given by the oligosaccharide-markers and the previously-published polysaccharide marker, the RSD value was not more than 6.4%.

In conclusion, this approach provided an efficient and reliable method to obtain accurate structure information of polysaccharides and quantify specific polysaccharide in herb formula. it is believed that ABEE-labeled oligosaccharides approach can be also applied in the analysis of other saccharide-dominant herbal materials which in turns helps to find out the bioactivity mechanism of polysaccharides.

TABLE OF CONTENTS

DECLARATION	i
ABSTRACT	ii
ACKNOWLEDGEMENTS	iv
TABLE OF CONTENTS	v
LIST OF TABLES	ix
LIST OF FIGURES	x
LIST OF ABBREVIATIONS	xiii
CHARTER 1 RECENT DEVELOPMENT ON QUALITY CONTROL OF <i>DENDROBIUM OFFICINALE</i>	1
1.1. General introduction of <i>Dendrobium officinale</i>	1
1.2. Authentication strategies on <i>Dendrobium officinale</i> based on small molecules and DNA barcoding.....	3
1.3. Quality control on polysaccharides	5
1.3.1. Introduction of polysaccharides	5
1.3.2. Extraction, purification, and fractionation of <i>Dendrobium</i> polysaccharides ...	7
1.3.3. Qualitative and quantitative analysis of polysaccharides.....	15
1.4. Current limitations on the quality control of <i>Dendrobium officinale</i> polysaccharides.....	24
1.5. Derivatization in carbohydrate analysis	25
1.6. Objectives of this study	29
CHARTER 2 Structural elucidation of polysaccharide backbone using a modified ABEE-labeled oligosaccharide approach: <i>Dendrobium officinale</i> , a case study.....	31
2.1. Introduction	31
2.2. Materials and methods	35
2.2.1. Materials	35

2.1.1.	Extraction of DOP	36
2.1.2.	Partial acid hydrolysis and ABEE derivatization	36
2.1.3.	UHPLC-Q-TOF-MS analysis for acid hydrolysates	37
2.1.4.	Isolation of the oligosaccharide markers	38
2.1.5.	ABEE-based sequence determination of isolated oligosaccharides by UHPLC-Q-TOF-MS	39
2.1.6.	Glycosidic linkages as determined by methylation reaction and GC-MS analysis	40
2.1.7.	Nuclear magnetic resonance (NMR) spectroscopy	42
2.3.	Results	42
2.3.1.	Protocol optimization for preparation of DOP hydrolysates (HDOP)	42
2.3.2.	Isolation and purity check of ABEE-labeled oligosaccharides	47
2.3.3.	Structure characterization of HDOP by monosaccharide composition and methylation	50
2.3.4.	Structure elucidation of ABEE-labeled oligosaccharides	51
2.3.5.	Structure elucidation of the backbone of DOP	61
2.3.6.	Limitation of structural characterization using methylation analysis.....	65
2.3.7.	Limitation of the proposed method on structure elucidation.....	70
2.4.	Discussion	73
2.5.	Conclusions	76
CHARTER 3 Oligosaccharide-marker Approach for Qualitative and Quantitative Analysis of Specific Polysaccharide in Herb Formula by UHPLC-Q-TOF-MS: <i>Dendrobium officinale</i> , a case study		78
3.1.	Introduction	78
3.2.	Material and methods	81
3.2.1.	Chemicals and materials	81
3.2.2.	Preparation of <i>Dendrobium officinale</i> polysaccharide-marker (DOP).....	82

3.2.3.	HPGPC-CAD analysis.....	83
3.2.4.	Monosaccharide composition analysis	84
3.2.5.	Preparation of water extract of Dendrobii samples, real product samples and Dendrobii blank product samples	85
3.2.6.	Partial acid hydrolysis and ABEE derivatization	86
3.2.7.	UHPLC-Q-TOF-MS analysis of acid hydrolysates.....	87
3.2.8.	Verification of the specificity of oligosaccharide markers in product samples	88
3.2.9.	Isolation and structure characterization of ABEE-labeled oligosaccharide markers	88
3.2.10.	Methylation reaction and GC-MS analysis	89
3.2.11.	Nuclear magnetic resonance (NMR) spectroscopy	91
3.2.12.	Linearity between the selected oligosaccharide markers and the polysaccharide marker DOP	91
3.2.13.	DOP determination in real samples	94
3.3.	Results and discussion.....	95
3.3.1.	Specificity of the oligosaccharide markers in herbal product samples	95
3.3.2.	Linearity between the daughter oligosaccharide markers and parent polysaccharide.....	101
3.3.3.	Method validation of quantitative analysis.....	104
3.4.	Quantitative analysis of real sample.....	108
3.5.	Conclusion.....	111
CHAPTER 4 SUMMARY AND FUTURE PROSPECTS		113
4.1.	Summary	113
4.2.	Future prospect	117
REFERENCES.....		123
PUBLICATIONS		139

CURRICULUM VITAE 141