

MASTER'S THESIS

Optimizing food waste composting process in fed-batch composter

Chan, Man Ting

Date of Award:
2015

[Link to publication](#)

General rights

Copyright and intellectual property rights for the publications made accessible in HKBU Scholars are retained by the authors and/or other copyright owners. In addition to the restrictions prescribed by the Copyright Ordinance of Hong Kong, all users and readers must also observe the following terms of use:

- Users may download and print one copy of any publication from HKBU Scholars for the purpose of private study or research
- Users cannot further distribute the material or use it for any profit-making activity or commercial gain
- To share publications in HKBU Scholars with others, users are welcome to freely distribute the permanent URL assigned to the publication

ABSTRACT

Composting is considered as an effective and sustainable food waste treatment technology from the perspectives of volume reduction, stabilization and releasing the pressure on landfills. Community composter is a decentralized composting facility in fed-batch operational mode which is usually being installed in the backyard of institutes, hospitals, housing estate etc. to handle the food wastes generated daily. Albeit numerous operational issues including high initial acidity and oil content, poor decomposition and odor generation are commonly encountered in these facilities, which make it difficult to be accepted by the public. Therefore, the aim of the present study is to develop a composting mix formulation that can provide a solution to all these issues in a fed-batch food waste composting process.

The first phase of this study aims at finding out an optimized formulation in a batch-scale food waste composting process through the use of alkaline amendments and microbial inoculum. For the first two experiments, artificial food wastes were prepared by mixing 1.3kg bread, 1kg boiled rice, 1kg cabbage, 0.5kg fully boiled pork and mixed with sawdust to obtain a C/N of 30 and adjusted moisture of the mixtures to 55%. The effect of different concentrations of zeolite compared to lime was studied in the first experiment. Zeolite was amended with food wastes and sawdust mixtures at 2% (ZI-2), 5% (ZI-5), 10% (ZI-10) to compare with lime in 2.25% (L-2.25) w/w (dry weight basis) and composted for 56 days. Results demonstrated that 10% of zeolite was optimal amendment rate compared to lower dosage of zeolite (2% & 5%) with stronger pH buffering capacity and greater decomposition efficiency. Addition of 2.25% of lime buffered the pH efficiently but increased the ammonia loss significantly which eventually reduced total nitrogen (TN) content of final product and posed odor emission problem. Amendment of 10% zeolite provided a higher adsorption affinity on ammonia resulting in 2.05% of TN value of final product which was higher than 1.72% of lime treatment. Furthermore, significantly higher seed germination 150% was achieved of ZI-10 compost compared to 135% of L-2.25 due to low ammonium content of product.

The first experiment showed that application of less than 10% zeolite was not sufficient to buffer the acidity; as a result, organic matter decomposition was inhibited. However, the cost and reduction in treatment percentage of food waste in 10% application rate of zeolite is an issue of concern. To tackle this dilemma, food waste was amended with struvite salts at 1:2 molar ratio of MgO and K₂HPO₄ (Mg:P) with or without zeolite amended at either 5% or 10% amendment (Mg:P, Z5 + Mg:P & Z10 + Mg:P) and a control treatment with food waste only was also included. Results showed that treatment of Z10 + Mg:P was synergistically achieved of pH and EC buffering, and N conservation but not for the case of 5 % zeolite. Treatment of Z10 + Mg:P further reduced the N loss to 18% compared to 25% and 27% of Mg:P and Z5 + Mg:P respectively. However, there was insignificant difference in the final nitrogen content and decomposition rate among all treatments with struvite salts amendment. Comparing to the treatment of Z-10 of the first experiment to Z10 + Mg:P of the second experiment, Z-10 showed superior performance since better decomposition efficiency, shorter time to require to pass the GI (28 Days) and lower cost because of salts exclusion.

To develop a multipurpose formulation for the fed-batch operational food waste composter, high lipids problem in food waste cannot be neglected because it is a critical factor to hinder the decomposition efficiency. Inoculation of oil degradative microorganisms was reported as an effective approach to facilitate the lipids. Therefore, the third experiment was to investigate the overall composting performance supplemented with 10% zeolite and microbial consortium. 10% zeolite with bacterial consortium significantly reduced the lipid contents from 7% to 1%

compared to control treatments. Furthermore, treatments amended with 10% zeolite was proved to reduce ammonia emission and total volatile fatty acids level in the composting mass, therefore the total odor emission level can be reduced. Zeolite at 10% was found to be a suitable optimum additive for both synthetic and real-food wastes. Therefore, treatment of 10% zeolite with bacterial consortium is selected as an optimized formulation for further study of its application in a fed-batch composter.

Following the food waste zeolite composting formulation obtained in Phase I, the aim of Phase II was to develop an ideal composting mix formulation for on-site commercial composters. Although the results have been demonstrated 10% zeolite with bacterial consortium facilitated the composting efficiency in batch composter, those amendments may be over-estimated if applied in a fed batch composter by using real food wastes. With this constraint, the applicability of these additives in commercial fed-batch composter needs to be assessed using locally generated food wastes. Treatments included food waste and sawdust mixtures at 4:1 mixing ratio (wet weight basis) were mixed with 2.25% of lime (L2.25), 10% of zeolite (Z10) and 10% zeolite with bacterial inoculum (Z10+O) and a control of food waste with sawdust mixture only was also included. 35 kg compost mixture was fed into each composter respectively daily for a period of 42 days. Only Z10+O was the most suitable composting mix for fed-batch food waste composting process with continuous sustained high temperature (55-60°C), optimal moisture (55%-60%), alkaline pH and low EC during the experimental period. Bacterial inoculum significantly improved the lipids decomposition from 22.16% (C) to 3.10% (Z10+O) after the composting period. In contrast, lime and zeolite alone treatments could not maintain the optimal pH that led to reduce degradation and longer stabilization period. Only compost taken from Z10+O treatment could be classified as mature compost.

The aim of the third study phase was to examine an optimal application rate of food waste compost produced from decentralized food waste composter for plant. A plant growth experiment was conducted in this phase to evaluate the change in soil properties and plant growth of *Brassica chinensis* and *Lycopersicon esculentum*. The experiment was conducted in a loamy soil amended with 0%, 2.5%, 5% and 10% food waste compost amendment rate compared to the control soil with chemical fertilizer amendment only. Results indicated that 5% was the optimal application rate of food waste compost for both crops among all treatments which can be evidenced by the highest biomass production and nutrients value of the plant tissues. Plant available nutrients such as NH_4^+ , NO_3^- , PO_4^{3-} were proportionally increased with increase in compost application rate. However, 2.5% of the food waste compost did not provide sufficient nutrients for plant growth and 10% showed negative effects due to increased salts content. Plants amended with chemical fertilizer had relatively low biomass production compared to compost amended treatments due to soil compaction and fast leaching of nutrients.

It can be concluded that application of 10% zeolite with microbial consortium is an ideal composting mix formulation for on-site commercial composters and 5% is an optimal application rate of food waste compost of *Brassica chinensis* and *Lycopersicon esculentum*.

TABLE OF CONTENTS

DECLARATION	1
ABSTRACT	2
ACKNOWLEDGEMENTS	4
LIST OF TABLES	9
LIST OF FIGURES	10
CHAPTER 1 AIMS AND OBJECTIVES	13
1.1 Background	13
1.2 Research Objectives and Outlines of the Study	15
CHAPTER 2 - LITERATURE REVIEW: OPTIMIZING FOOD WASTE COMPOSTING PROCESS IN FED BATCH COMPOSTER.....	18
2.1 Introduction.....	18
2.2 Definition of Composting	19
2.3 Composting Process	19
2.4 Food Waste Composting	21
2.4.1 Acidity	22
2.4.2 Odor	23
2.4.3 Lipids content	24
2.4.4 Loss of nutrients	25
2.5 Factors affecting food waste composting process	26
2.5.1 Properties of raw materials.....	27
2.5.1.1 Carbon to nitrogen ratio	27
2.5.1.2 Moisture	28
2.5.1.3 Particle size.....	29
2.5.1.4 Temperature.....	30
2.5.1.5 pH.....	31
2.5.1.6 Oxygen level	31
2.6 Enhancing food waste composting	32
2.6.1 Zeolite.....	32
2.6.2 Lime.....	33
2.6.3 Struvite	34
2.6.4 Microbial inoculums.....	36
2.7 Compost maturity evaluation	37
2.7.1 Physical tests	37
2.7.2 Chemical tests	39
2.7.2.1 pH.....	39
2.7.2.2 Electrical conductivity	40
2.7.2.3 Ammonium, nitrite and nitrate	41
2.7.2.4 Soluble organic carbon.....	42
2.7.2.5 Carbon to nitrogen (C/N) ratio	43
2.7.2.6 Organic matter reduction.....	44
2.7.2.7 Humic acids	44
2.7.3 Biological tests	45
2.7.3.1 Oxygen uptake	46
2.7.3.2 Carbon dioxide evolution.....	46
2.7.3.3. Enzymatic activity	47
2.7.3.4 Phytotoxicity.....	48

2.8 Compost application	48
2.8.1 Effect of compost on soil properties	49
2.8.2 Effect of compost on plants	49
2.9 Introduction of community composters	50
2.9.1 Poor decomposition	51
2.9.2 Odor emission	52
2.9.3 Production of immature compost	53
CHAPTER 3 - EFFECT OF ZEOLITE ON FOOD WASTE COMPOSTING IN LABORATORY SCALE BATCH COMPOSTERS	54
3.1 Introduction.....	54
3.2 Materials and Methods.....	56
3.2.1 Feedstock Preparation	56
3.2.2 Experimental Design and compost reactor operation.....	57
3.3 Statistical analysis	61
3.4 Results and discussion	61
3.4.1 Changes of temperature	61
3.4.2 Carbon dioxide evolution.....	62
3.4.3 Changes of pH	63
3.4.4 Changes of EC.....	65
3.4.5 Ammonia emission.....	66
3.4.6 Changes of extractable ammonium	68
3.4.7 Loss of total organic matter, total organic carbon and total organic nitrogen during food waste composting.....	70
3.4.8 Compost Quality.....	71
3.4.8.1 Compost Maturity	71
3.4.8.2 Seed Germination Index	72
3.4.8.3 Nutrient content.....	73
3.5 Conclusions.....	74
CHAPTER 4 - FEASIBILITY OF CO-COMPOSTING FOOD WASTE COMPOSTING WITH ZEOLITE AND STRUVITE SALTS	76
4.1 Introduction.....	76
4.2 Materials and Methods.....	77
4.2.1 Feedstock preparation	77
4.2.1.1 Struvite formation	78
4.2.2 Composter and operation	78
4.2.3 Sampling	78
4.2.4 Treatments	79
4.2.5 Composting samples analysis	79
4.3 Results and Discussion	79
4.3.1 Change of temperature and carbon dioxide evolution.....	79
4.3.2 Changes in pH profile	82
4.3.3 Changes in electrical conductivity (EC).....	83
4.3.4 Changes in extractable ammonium nitrogen (NH ₄ ⁺ -N) and ammonia (NH ₃).....	84
4.3.5 Changes in total organic carbon (TOC), total nitrogen (TN) and C/N ratio	88
4.3.6 Loss of TOM, TOC and TN.....	91
4.3.7 Seed germination index (GI).....	92
4.3.8 Final compost characteristics	93
4.4 Conclusion	96
CHAPTER 5 - INFLUENCE OF OIL DEGRADING BACTERIAL CONSORTIUM ON FOOD WASTE COMPOSTING WITH ZEOLITE	97

5.1 Introduction.....	97
5.2 Materials and Methods.....	99
5.2.1 Feedstock Preparation	99
5.2.2 Composter and operation	100
5.2.3 Sampling	100
5.2.4 Treatments	101
5.2.5 Analyses	101
5.3 Results and Discussions	102
5.3.1 Change of temperature.....	102
5.3.2 Change of carbon dioxide evolution and cumulative carbon dioxide evolution	104
5.3.3 Changes of pH and electrical conductivity	105
5.3.4 Nitrogen dynamics	108
5.3.5 Changes in total organic carbon (TOC) and total organic matter (TOM)	111
5.3.6 Changes in seed germination index (GI).....	112
5.3.7 Changes in total lipid contents	114
5.3.8 Changes of water soluble fatty acids (VFAs)	116
5.4 Conclusion	118

CHAPTER 6 - OPTIMIZATION FOOD WASTE COMPOSTING IN FED-BATCH
COMMUNITY-COMPOSTERS: INFLUENCE OF ZEOLITE AND BACTERIAL INOCULATION 120

6.1 Introduction.....	120
6.2 Materials and Methods.....	122
6.2.1 Food wastes separation and collection	122
6.2.2 Food waste composter	122
6.2.3 Preparation of the starting culture	124
6.2.4 Preparation of composting mix input.....	125
6.2.5 Treatments	126
6.2.6 Sampling	127
6.2.7 Analyses	127
6.3 Results and Discussion	128
6.3.1 Changes of temperature, moisture, pH and EC.....	128
6.3.2 Changes of total organic carbon, total Kjeldahl nitrogen and carbon to nitrogen ratio	133
6.3.4 Changes of total lipid content during fed-batch food waste composting process.....	137
6.3.5 Changes of total bacteria count during fed-batch food waste composting process	141
6.3.6 Evaluation of ammonia emission (NH ₃) during fed-batch food waste composting process.....	142
6.3.7 Changes of water soluble fatty acids during fed-batch food waste composting process.....	145
6.4 Physicochemical properties of compost of different treatments in fed-batch operation	148
6.5 Conclusion	150

CHAPTER 7 - INFLUENCE OF FOOD WASTE COMPOSTS FROM COMMUNITY COMPOSTER ON
PLANT GROWTH..... 152

7.1 Introduction.....	152
7.2 Materials and methods	153
7.2.1 Plant species.....	153
7.2.2 Soil.....	154
7.2.3 Food waste compost	154
7.2.4 Chemical fertilizer.....	155
7.2.5 Treatments	156
7.2.6 Experimental setup and preparation	158
7.2.7 Collection of soil and plant samples	159
7.2.8 Analytical methods.....	160
7.2.8.1 Soil and compost analysis	160
7.2.8.2 Plant analysis	161

7.3 Statistical analysis	161
7.4 Results and Discussion	161
7.4.1 Properties of food waste compost and soil	161
7.4.2 Seed germination	162
7.4.3 Changes of pH	164
7.4.4 Changes of EC.....	166
7.4.5 Changes of extractable ammonium	168
7.4.6 Changes of nitrate.....	171
7.4.7 Changes of extractable phosphorus.....	173
7.4.8 Changes of total nitrogen.....	175
7.4.9 Changes of total organic matter	177
7.5 Changes of total phosphorus	179
7.5.1 Biomass production.....	181
7.5.2 Total nitrogen and total phosphorus of plant tissues.....	183
7.6 Conclusions.....	185
CHAPTER 8 - GENERAL DISCUSSION, CONCLUSIONS AND RECOMMENDATIONS.....	186
8.1 Introduction.....	186
8.2 Efficiency of Zeolite and Oil-degradative inoculum in batch mode food waste composting.....	187
8.3 Pilot scale application of Zeolite and Oil-degradative inoculum in food waste composting using fed batch composter	189
8.4 Influence of zeolite amended compost on soil and plant	190
8.5 Recommendations for further study	190
8.5.1 Feasibility of reducing dosage of 10% zeolite.....	190
8.5.2 Practicability of applying the oil degradative inoculum in large scale composter	191
REFERENCES	193
CURRICULUM VITAE	212