

MASTER'S THESIS

Applications of the subtractive hybridization method to study gene expression in rat liver after cadmium exposure

Tang, Mei Kuen

Date of Award:
1994

[Link to publication](#)

General rights

Copyright and intellectual property rights for the publications made accessible in HKBU Scholars are retained by the authors and/or other copyright owners. In addition to the restrictions prescribed by the Copyright Ordinance of Hong Kong, all users and readers must also observe the following terms of use:

- Users may download and print one copy of any publication from HKBU Scholars for the purpose of private study or research
- Users cannot further distribute the material or use it for any profit-making activity or commercial gain
- To share publications in HKBU Scholars with others, users are welcome to freely distribute the permanent URL assigned to the publication

Application of the Subtractive Hybridization Method
to study Gene Expression in Rat Liver
after Cadmium Exposure

TANG Mei Kuen

A thesis submitted in partial fulfillment of the requirements
for the degree of
Master of Philosophy

September 1994

Hong Kong Baptist College

(The Predecessor of the Hong Kong Baptist University)



Sijt
12-P-85
13328811
T-1

ABSTRACT

Induction of gene expression in male Sprague-Dawley rat liver after a sublethal dose of CdCl_2 was studied using the subtractive hybridization method. Cd-induced liver cDNA was enriched by subtractive hybridization using normal liver mRNA as driver. The subtracted cDNA was further amplified by polymerase chain reaction and labeled for screening of Cd-induced cDNA clones from the cDNA library. Around 70 positive clones were identified. One of these clones was identified as rat α -2u globulin after DNA sequence analysis of the PCR product followed by a search for homology in the gene data banks. Northern blot analysis confirmed that besides metallothionein, α -2u globulin is also inducible by Cd.

TABLE OF CONTENTS

Abstract		i
Acknowledgments		ii
Table of Abbreviations		iii
Table of Contents		iv
Part I: Introduction		
1.1	General background	1
1.2	Cadmium Toxicity	4
1.3	Metallothionein as a stress protein	7
	1.3.1 Nomenclature and occurrence	7
	1.3.2 Structure of mammalian MT	9
	1.3.3 Induction of Metallothionein	10
	1.3.4 Induction of MT after Cd exposure	12
	1.3.5 Functions of Metallothionein	12
	1.3.6 MT as a bioindicator of metal exposure	16
1.4	Subtractive Hybridization	17
	1.4.1 Background information	17
	1.4.2 Subtractive hybridization technique	18
1.5	The objective of this study	20
Part II: Materials and Methods		
2.1	Materials	22
2.2	Animal Treatment	22

2.3	Extraction of rat liver total RNA	23
	2.3.1 Monitoring the integrity of total RNA by agarose gel electrophoresis	24
2.4	Purification of poly (A)+ RNA by Oligo (dT)-cellulose column chromatography	24
2.5	Construction of Cd-induced cDNA library	25
	2.5.1 First Strand synthesis	25
	2.5.2 Second Strand synthesis	25
	2.5.3 Addition of EcoRI/NotI site onto the cDNA	27
	2.5.4 Ligation of cDNA with EcoRI digested vector λ gt11	27
2.6	Preparation of MT cDNA probe	28
	2.6.1 Preparation of competent cells	28
	2.6.2 Large-scale Preparation of Plasmid DNA containing MT cDNA	30
	2.6.3 Preparation of MT cDNA fragment by restriction enzyme digestion	31
	2.6.4 Confirmation of the complete restriction enzyme digestion by polyacrylamide gel electrophoresis	31
	2.6.5 Elution of MT cDNA fragments from polyacrylamide gel slice	32
2.7	Northern Blotting	32
2.8	Procedures for Subtractive hybridization	34
	2.8.1 Subtractive hybridization technique to pool the Cd-induced cDNA	36
2.9	<i>In vitro</i> amplification of subtracted Cd-induced cDNA	38
2.10	<i>In Vitro</i> amplification of the cDNA insert in the library	38

2.11	Preparation of ³² P-labeled subtracted Cd-induced cDNA by random primer labeling method	39
2.12	Screening of Cd-inducible clones from the rat liver cDNA library	39
2.13	Plaque hybridization	40
2.14	Asymmetric PCR for amplification of DNA template	40
2.15	DNA sequencing	41
2.16	Polyacrylamide gel electrophoresis for DNA sequence analysis	42
2.17	Data analysis	42
Part III:	Results	
3.1	Induction of MT from rat liver	43
3.2	Separation of the Cd-induced genes by the subtractive hybridization technique	54
3.3	Screening of Cd-induced cDNA clones	54
	3.3.1 Construction of Cd-treated rat liver cDNA library	54
	3.3.2 Selection of Cd-induced recombinant clones from the cDNA library	56
3.4	Identification of Rat Cd-1	57
Part IV:	Discussion and Conclusion	
4.1	Induction of MT in rat liver	68
4.2	Subtractive Hybridization technique	70
4.3	α ₂ u-globulin as a Cd-induced gene	71
4.4	Comparison of α ₂ u-globulin with Cd-binding-protein -- MT	75

4.4	Conclusion	77
	References	79
	Appendix	98
	Vita	111