

DOCTORAL THESIS

Earthworms and mycorrhizae in phytoremediation of Pb/Zn mine tailings: their effects on metal speciation, bioavailability and uptake by *Leucaena leucocephala*

Ma, Ying

Date of Award:
2003

[Link to publication](#)

General rights

Copyright and intellectual property rights for the publications made accessible in HKBU Scholars are retained by the authors and/or other copyright owners. In addition to the restrictions prescribed by the Copyright Ordinance of Hong Kong, all users and readers must also observe the following terms of use:

- Users may download and print one copy of any publication from HKBU Scholars for the purpose of private study or research
- Users cannot further distribute the material or use it for any profit-making activity or commercial gain
- To share publications in HKBU Scholars with others, users are welcome to freely distribute the permanent URL assigned to the publication

**Earthworms and Mycorrhizae in Phytoremediation of
Pb/Zn Mine Tailings: Their Effects on Metal Speciation,
Bioavailability and Uptake by *Leucaena leucocephala***

MA Ying

**A thesis submitted in partial fulfillment of the requirements
for the degree of
Doctor of Philosophy**

Principal Supervisor: Prof. WONG Ming Hung

Hong Kong Baptist University

May 2003

ABSTRACT

The present study aims to investigate the influence of earthworms and arbuscular mycorrhizae on lead and zinc speciation, bioavailability and uptake by *Leucaena leucocephala*, for phytoremediation of Pb/Zn mine tailings.

Pb/Zn mine tailings were found to be toxic to the earthworm *Pheretima*, unless the tailings were diluted by at least 20% with an artificial uncontaminated soil. LC₅₀ values showed *Pheretima guillelmi* was the more tolerant species than *Pheretima sp.* Soil pH, EC and DTPA-Pb and -Zn significantly increased following inoculation with the earthworms, and the latter two parameters were increased by up to 48.2% and 24.8% respectively.

Single and sequential chemical extraction procedures were conducted to study Pb and Zn speciation and bioavailability as affected by earthworms in a plant-soil system. It was demonstrated that earthworms significantly increased Zn bioavailability due to decreased soil pH, increased DOC, and earthworm gut-related processes. However, Pb bioavailability in soils was not increased, which was probably due to the formation of insoluble complexes with phosphates to decrease reducible-Pb in soil.

The effects of earthworms on phytoextraction of Pb/Zn mine tailings were investigated in pot experiments using Indian mustard (*Brassica juncea*). It was found that earthworms increased metal bioavailability, largely through much higher extractable metals in earthworm casts. The results indicated that accumulator plants, in combination with earthworm inoculation, may be suitable for remediating low-level metal contamination. In mine tailings, however, excluder plants or plants that retain metals in roots are likely to be more useful for ecological restoration.

The effects of earthworms on the growth of *Leucaena leucocephala* on Pb/Zn mine tailings diluted with varying amounts of mineral soil were studied. *L. leucocephala* grew successfully on tailings with a 25% (w/w) soil amendment. The presence of earthworms improved the yield of plants by 10-30%. *P. guillelmi* increased bioavailable metal concentrations in the amended tailings, accompanied by a direct increase of metal uptake by the plants, and the increased rate of metal uptake into plants from tailings ranged from 16% to 53%.

Inoculation of arbuscular mycorrhizal fungi (AMF) was found to significantly improve growth and establishment of *L. leucocephala* on Pb/Zn mine tailings. Other soil microorganisms also facilitated plant growth, although the effect was less obvious. Plant root mycorrhizal infection was positively related to soil phosphatase activity and dehydrogenase activity. Higher root mycorrhizal infection increased Pb and Zn uptake by plants under lower metal concentrations, while it decreased Pb and Zn uptake under higher metal concentrations.

A synergistic effect between earthworms and arbuscular mycorrhizae on plant

growth was observed in higher metal concentrations. Earthworms considerably improved root mycorrhizal infection and somewhat stimulated plant root nodulation, while arbuscular mycorrhizae exerted positive effects on plant nodulation and nodule efficiency. It was demonstrated that earthworms stimulated plant growth mainly through enhancing soil microbial activity, improving soil nutrient availability, and increasing plant N, P uptake. Arbuscular mycorrhizae improved plant growth through stimulating root nodule efficiency and increasing plant N, P and K uptake. Uptake of Pb and Zn by *L. leucocephala* grown in Pb/Zn mine tailings was significantly enhanced by earthworms, most probably due to the increased plant dry weight production. However, AMF inoculation did not increase metal uptake. Arbuscular mycorrhizae were proved to protect plants from metal stress through restricting metal uptake to shoots and decreasing metal bioavailability in soil.

TABLE OF CONTENTS

	P. No
DECLARATION	i
ABSTRACT	ii
ACKNOWLEDGEMENT	iv
TABLE OF CONTENTS	v
LIST OF TABLES	x
LIST OF FIGURES	xii
CHAPTER I: GENERAL INTRODUCTON	
1.1. Background of Research	
1.1.1. Current situation concerning heavy metal contaminated soil and metal mined land in China	1
1.1.2. Comparison of traditional methods for treatment of heavy metal contaminated soil	3
1.1.3. Types and advantages of phytoremediation	6
1.1.4. Trees, earthworms and mycorrhizae in phytoremediation	7
1.2. Review of Literature	
1.2.1. Decontamination of heavy metals using trees	9
1.2.2. The role of earthworms in soils and metal contaminated sites	
1.2.2.1. Effects of earthworms on soil nutrient availability and plant growth	12
1.2.2.2. Effects of heavy metal toxicity on earthworms	14
1.2.2.3. Accumulation of heavy metals in earthworms	14
1.2.2.4. Earthworm inoculation of soils and uses in reclamation	16
1.2.3. The role of mycorrhizae in soils and metal contaminated sites	
1.2.3.1. Effects of mycorrhizae on plant growth	17
1.2.3.2. Interaction between mycorrhizae and heavy metals	18
1.2.4. Conclusion	20
1.3. Objectives of Research	21
1.4. Contributions and Significance of Research	24
1.5. Framework of Research	24

CHAPTER II: TOXICITY OF LEAD/ZINC MINE TAILINGS TO EARTHWORMS AND THE EFFECT OF BURROWING ON METAL AVAILABILITY

2.1. Introduction	26
2.2. Materials and Methods	
2.2.1. Experiments	28
2.2.2. Analyses	30
2.3. Results	
2.3.1. Properties of Pb/Zn mine tailings	31
2.3.2. Earthworm survival and growth	31
2.3.3. Metal mobility	32
2.4. Discussion	44
2.5. Conclusion	48

CHAPTER III: CHEMICAL SPECIATION AND BIOAVAILABILITY OF LEAD AND ZINC AS AFFECTED BY BURROWING EARTHWORMS

3.1. Introduction	49
3.2. Materials and Methods	
3.2.1. Preparation of Pb and Zn spiked soils	51
3.2.2. Greenhouse experiment	52
3.2.3. Metal speciation of Pb and Zn	
3.2.3.1. Single chemical extraction	53
3.2.3.2. Sequential chemical extraction	54
3.2.4. Statistical analysis	56
3.3. Results	
3.3.1. General properties of test soil	57
3.3.2. Soil pH, EC and DOC as affected by earthworm activity	57
3.3.3. Single chemical extractability of soil Pb and Zn	61
3.3.4. Speciation and fractionation of soil Pb and Zn	61
3.3.5. Correlations among soil metal fractions	62
3.3.6. Plant growth and Pb and Zn bioavailability	69
3.4. Discussion	
3.4.1. Soil pH, EC, DOC and earthworms	77
3.4.2. Pb and Zn speciation and earthworms	78
3.4.3. Plant growth and earthworms	83
3.4.4. Pb and Zn bioavailability and earthworms	84
3.5. Conclusion	85

**CHAPTER IV: PHYTOREMEDIATION OF LEAD/ZINC MINE
TAILINGS: EARTHWORM-ENHANCED METAL UPTAKE**

4.1. Introduction	86
4.2. Materials and Methods	
4.2.1. Assessment of natural metal mobility by Dutch leaching test	87
4.2.2. Trial of inoculation Pb/Zn mine spoils with earthworms	88
4.2.3. Greenhouse experiment on phytoremediation of Pb/Zn mine spoils	89
4.2.4. Analyses	90
4.3. Results	
4.3.1. Properties of Pb/Zn mine spoils	91
4.3.2. Actual leachability and leaching behavior of Pb/Zn mine spoils	91
4.3.3. Effect of earthworms on DTPA-extractable Pb and Zn	91
4.3.4. Effect of earthworms on metal uptake from spoils by <i>Brassica juncea</i>	
4.3.4.1. Plant growth and biomass production	96
4.3.4.2. Removal of Pb and Zn from spoils	96
4.3.4.3. Bioavailable Pb and Zn in mine spoils	96
4.4. Discussion	
4.4.1. Leaching and extraction of metals from Pb/Zn mine spoils	105
4.4.2. Effects of earthworms on bioavailability of metals	106
4.4.3. Role of earthworms in phytoremediation	107
4.5. Conclusion	112

**CHAPTER V: REMEDIATION OF LEAD/ZINC MINE TAILINGS USING TREES:
EARTHWORMS, SOIL NUTRITION AND METAL MOBILITY**

5.1. Introduction	114
5.2. Materials and Methods	
5.2.1. Experimental materials	115
5.2.2. Experimental design	116
5.2.3. Analyses	117
5.3. Results	
5.3.1. Soils, earthworms and plant growth	119
5.3.2. Nutrients in soils and plants	127
5.3.3. Pb and Zn	134
5.4. Discussion	
5.4.1. Earthworm survival and plant growth affected by Pb/Zn mine tailings	142
5.4.2. Effects of earthworms on soil nutrition and N/P uptake by plants	143
5.4.3. Effects of earthworms on Pb and Zn uptake by plants	145
5.5. Conclusion	146

CHAPTER VI: INTERACTION OF ARBUSCULAR MYCORRHIZAE AND OTHER SOIL MICORBES ON TREE SEEDLINGS GROWTH AND METAL UPTAKE IN LEAD/ZINC MINE TAILINGS

6.1. Introduction	148
6.2. Materials and Methods	
6.2.1. Experimental materials	150
6.2.2. Experimental design	151
6.2.3. Analyses	152
6.3. Results	
6.3.1. Properties of Pb/Zn mine tailings and topsoil mixture	155
6.3.2. Plant biomass and nutrient uptake	155
6.3.3. Mycorrhizal root colonization and soil microbial characteristics	163
6.3.4. Pb and Zn uptake by <i>Leucaena leucocephala</i>	168
6.4. Discussion	
6.4.1. AMF colonization as related to plant growth and nutrient uptake	170
6.4.2. Pb and Zn uptake as affected by AMF infection	175
6.4.3. Interactions between AMF infection and soil microorganisms	179
6.5. Conclusion	181

CHAPTER VII: INTERACTIONS BETWEEN EARTHWORMS AND ARBUSCULAR MYCORRHIZAE IN TREE SEEDLING GROWTH AND METAL UPTAKE IN LEAD/ZINC MINE TAILINGS

7.1. Introduction	183
7.2. Materials and Methods	
7.2.1. Experimental materials	185
7.2.2. Experimental design	186
7.2.3. Analyses	187
7.3. Results	
7.3.1. Properties of Pb/Zn mine tailings and topsoil mixture	189
7.3.2. The pH, EC, DOC and available nutrients in plant growth substrata	190
7.3.3. Soil phosphatase activity, dehydrogenase activity, plant root nodulation and mycorrhizal root colonization	193
7.3.4. Plant growth and nutrient uptake	194
7.3.5. Pb and Zn bioavailability and uptake by <i>L leucocephala</i> and earthworms	202
7.4. Discussion	
7.4.1. Soil pH, EC and DOC as affected by earthworms and arbuscular mycorrhizae	210
7.4.2. Effects of earthworms and arbuscular mycorrhizae on plant growth related to nutrient uptake	211
7.4.3. Effects of earthworms and arbuscular mycorrhizae on Pb and Zn bioavailability	

and uptake by <i>Leucaena leucocephala</i>	216
7.4.4. Interactions between earthworms, arbuscular mycorrhizae and root nodulation of <i>L. leucocephala</i> in Pb/Zn mine tailings	218
7.5. Conclusion	222
 CHAPTER VIII: GENERAL DISCUSSION AND CONCLUSION	
8.1. Introduction	224
8.2. Earthworms in Phytoremediation of Pb/Zn Mine Tailings	
8.2.1. Earthworms and Pb and Zn speciation and bioavailability	227
8.2.2. Earthworms and plant growth	230
8.2.3. Earthworms and Pb and Zn uptake by plant	232
8.3. Arbuscular Mycorrhizae in Phytoremediation of Pb/Zn Mine Tailings	
8.3.1. Arbuscular mycorrhizae and plant growth	233
8.3.2. Arbuscular mycorrhizae and Pb and Zn uptake by plants	234
8.4. Combining Earthworms, Arbuscular Mycorrhizae and Leguminous Plant for Phytoremediation of Metal Contaminated Soils	236
8.5. Future Work	
8.5.1. Effect of soil pH on the change of metal bioavailability caused by earthworm inoculation	238
8.5.2. Characterization of metal compartmentation and distribution in earthworm tissues	239
8.5.3. Field investigation	240
8.6. General Conclusion	240
 REFERENCES	245
 PUBLICATIONS	279
 CURRICULUM VITAE	280