

MASTER'S THESIS

The role of calcium in control of animal cell proliferation: ornithine decarboxylase induction by L-asparagine in Reuber H-35 hepatoma cell

Pong, Nga Hin

Date of Award:
1991

[Link to publication](#)

General rights

Copyright and intellectual property rights for the publications made accessible in HKBU Scholars are retained by the authors and/or other copyright owners. In addition to the restrictions prescribed by the Copyright Ordinance of Hong Kong, all users and readers must also observe the following terms of use:

- Users may download and print one copy of any publication from HKBU Scholars for the purpose of private study or research
- Users cannot further distribute the material or use it for any profit-making activity or commercial gain
- To share publications in HKBU Scholars with others, users are welcome to freely distribute the permanent URL assigned to the publication

**THE ROLE OF CALCIUM IN CONTROL OF ANIMAL CELL
PROLIFERATION: ORNITHINE DECARBOXYLASE INDUCTION BY
L-ASPARAGINE IN REUBER H-35 HEPATOMA CELLS**

PONG, NGA HIN

A thesis submitted in partial fulfillment of the requirements
for the degree of Postgraduate Studies Committee
for the degree of Master of Philosophy

September 1991

Hong Kong Baptist College



Gift
10-6-92
26599

TH
M. PHIL
1991 PR

ABSTRACT

Cellular ornithine decarboxylase (ODC, L-ornithine carboxyl-lyase; EC 4.1.1.17) can be induced by each of the transport system A, ASN or N amino acids, most potently asparagine. The involvement of calcium in various cellular processes including cell proliferation has been well established. ODC is the rate-limiting enzyme in the biosynthesis of polyamines which are required for cell proliferation.

In this study, the role of Ca^{2+} in asparagine-stimulated ODC induction was investigated in Reuber H-35 hepatoma cells. Dose response effect of extracellular Ca^{2+} on enhancing ODC induction was observed. Additions of CaCl_2 above 0.1 mM to the culture medium enhanced asparagine-elicited ODC activity to maximal level. When extracellular Ca^{2+} was removed, ODC activity decreased significantly. Moreover, asparagine-stimulated ODC activity was strongly antagonized by the addition of a calcium flux blocker Lanthanum ion (La^{3+}) or a Ca^{2+} chelator EGTA (ethylene glycol (β -aminoethyl ether)-N-N' tetra-acetic acid). Additional extracellular Ca^{2+} was able to reverse the EGTA-inhibited ODC induction. These observations indicate the requirement of extracellular Ca^{2+} for a favorable ODC induction. Using a Ca^{2+} ionophore, ionomycin, and a blocker of Ca^{2+} release from intracellular store, TMB-8 (8-(N,N-diethyl amino)octyl-3,4,5,-trimethoxybenzoate), it was found that depletion of intracellular stored Ca^{2+} or blockage of intracellular Ca^{2+} release could not inhibit asparagine-stimulated ODC activity. Accordingly, we are able to conclude that extracellular Ca^{2+} , but not intracellular Ca^{2+} , is important for ODC induction.

Several types of voltage dependent calcium channel (VDCC), namely L, N and T, that mediate Ca^{2+} entry across cell membrane, were investigated. Using VDCC blockers, NiCl_2 , CdCl_2 , verapamil and nifedipine, it was found that L and N type VDCC channels are probably involved in the process of ODC induction, probably by mediating influx of extracellular Ca^{2+} .

The involvement of calmodulin in a variety of Ca^{2+} -dependent cellular functions has been well established. The effect of the calmodulin antagonist W_7 (N-(6-aminoethyl)-5-chloro-1-naphthalenesulfonamide) on ODC induction was

investigated. It was shown that the enzyme induced by asparagine was potently blocked by W_7 with 50% inhibition at $23 \mu\text{M}$, indicating that ODC induction is likely to be mediated by a Ca^{2+} -calmodulin regulatory process.

Initial investigation in the role of cyclic AMP and protein kinase C systems in ODC induction was conducted. Using cAMP modulators Rp- and Sp-cAMP, isomers of adenosine 3'-5'-monophosphorothioate, we found that Rp-cAMP, an antagonist of cAMP-dependent protein kinases, had no inhibition effect on asparagine-elicited ODC induction and that the Sp-cAMP, an agonist of cAMP-dependent protein kinases, could neither induce ODC activity by itself nor enhance asparagine effect on ODC induction. However, protein kinase C inhibitor H_7 (1-(5-isoquinolinesulfonamyl)-2-methylpiperazine), potently inhibited asparagine-induced ODC activity with 50% inhibition at $60 \mu\text{M}$ H_7 . It can be concluded that protein kinase C, but not cAMP system, is involved in ODC induction by asparagine.

It appears that the involvement of calcium in ODC induction requires participation of calmodulin regulation which collaborates with protein kinase C other than cAMP dependent protein kinases.

TABLE OF CONTENTS

	<u>Page</u>
Introduction	1
Materials and methods	36
Results	42
Discussion	68
Conclusion	87
Appendix	89
References	90
Vita	112