

DOCTORAL THESIS

Investigations of the industrial compatible aqueous purification techniques and intrinsic stabilizing factors for nattokinase

Xin, Xiong

Date of Award:
2016

[Link to publication](#)

General rights

Copyright and intellectual property rights for the publications made accessible in HKBU Scholars are retained by the authors and/or other copyright owners. In addition to the restrictions prescribed by the Copyright Ordinance of Hong Kong, all users and readers must also observe the following terms of use:

- Users may download and print one copy of any publication from HKBU Scholars for the purpose of private study or research
- Users cannot further distribute the material or use it for any profit-making activity or commercial gain
- To share publications in HKBU Scholars with others, users are welcome to freely distribute the permanent URL assigned to the publication

Abstract

Nattokinase is a potent fibrinolytic enzyme produced by *Bacillus subtilis*, which is one of the most important sources of fibrinolytic enzymes. Nattokinase has a dual function of hydrolyzing blood thrombin directly and indirectly. It was found to be active between pH 6-12 and lose fibrinolytic activity dramatically when pH is lower than 5. Currently nattokinase preparations are not pure enough to get rid of the unique smell that is repulsive to the non-Japanese. These two reasons severely blocked nattokinase to be developed for food and pharmaceutical industrial uses as supplements and active ingredients. The objective of this study was to isolate the bacteria which have ability to produce nattokinase-like fibrinolytic enzyme, purify and characterize these enzymes to determine the actual biochemical properties. An industrial compatible aqueous protocol can be applied not only for the purification of nattokinase but also for other fibrinolytic enzymes from *Bacillus* source was established to purify the enzymes to homogeneous. The intrinsic stabilizing factors in raw nattokinase fermented broth were also identified in order to make natural stable nattokinase preparation for acidic environment. 71 strains belonging to 13 different genus was determined as the fibrinolytic enzyme producing bacteria. The fibrinolytic enzyme produced by *Bacillus tequilensis*, *Bacillus amyloliquefaciens*, and *Bacillus cereus* was purified and characterized. One of these three enzymes was determined to be a new fibrinolytic enzyme that never be reported. The purification protocol established here contained 3 operation units that including one chromatographic separation step followed by membrane polishing after nattokinase was extracted with 70% ammonium sulphate from the fermented broth. The purified

nattokinase and other fibrinolytic enzymes showed single band as tested by SDS-PAGE and already pure enough to determine the N-terminal sequence directly by Edman degradation. A purification efficiency of 476.1 fold enzymatic activity increase with 48.3% recovery was obtained by using this protocol to purify nattokinase at lab level. The efficiency was 428.1-fold with 42.6% recovery at industrial compatible pilot scale (60-fold amplification). The efficiency was 329.7-fold with 42.7% recovery for purification of fibrinolytic enzyme produced by *Bacillus tequilensis*, 221.7-fold with the recovery of 32.5% for purification of fibrinolytic enzyme produced by *Bacillus amyloliquefaciens* and 288.5 fold with 38.7% recovery for purification of fibrinolytic enzyme produced by *Bacillus cereus*. The purification fold of most current protocols used for nattokinase and other fibrinolytic enzymes purification was lower than 100 and the recovery fall in between 6.28% to 80%. Thus, the protocol established in this study has a very high purification efficiency. The result of intrinsic stabilizing factor identification shown both starch and levan have the stabilizing effect on nattokinase at low pH environment. The stabilizing ability of starch is much higher than levan, and was retained even hydrolyzed by amylase. The active concentration range of starch was from 20 and up to 500 µg/ml at the pH range of 4-5.

Key words: Nattokinase; Fibrinolytic enzyme; *Bacillus*; Purification; Stabilizing factor

Table of Content

DECLARATION	I
ABSTRACT	II
ACKNOWLEDGMENTS	IV
TABLE OF CONTENT	V
LIST OF FIGURES	VII
LIST OF TABLES	IX
ABBREVIATION	XI
CHAPTER 1 GENERAL INTRODUCTION AND LITERATURE REVIEW	1
1.1 INTRODUCTION.....	1
1.2 THROMBOLYTIC AGENTS.....	4
1.3 NATTOKINASE (NK)	8
1.4 MEDICAL EFFECTS OF NK.....	10
1.5 MOLECULAR ENGINEERING AND FIBRINOLYTIC ENZYMES.....	14
1.6 EXTRACTION AND PURIFICATION OF NATTOKINASE.....	17
1.7 STABILITY OF NATTOKINASE	19
1.8 THE MEASUREMENT OF NATTOKINASE ACTIVITY	21
1.9 CURRENT ISSUES AND PROBLEMS.....	24
1.10 HYPOTHESIS	26
1.11 OBJECTIVES	27
1.12 SIGNIFICANCE AND OUTCOMES	28
CHAPTER 2 ISOLATION AND IDENTIFICATION OF BACTERIA PRODUCING FIBRINOLYTIC ENZYMES AND THE PHYLOGENETIC ANALYSIS OF BACTERIAL GENOMIC STRUCTURES	30
2.1 INTRODUCTION.....	30
2.2 MATERIALS	33
2.3 METHODS	33
2.3.1 <i>Sample collection</i>	33
2.3.2 <i>Bacteria isolation and culture</i>	33
2.3.3 <i>Bacterial identification</i>	34
2.3.4 <i>PCR purification</i>	37
2.3.5 <i>Restriction Fragment Length Polymorphism (RFLP)</i>	38
2.3.6 <i>Sequencing</i>	38
2.3.7 <i>Homologous analysis of bacteria produce fibrinolytic enzyme</i>	38
2.3.8 <i>Fibrin plate assay for fibrinolytic activity</i>	39
2.3.9 <i>Serological plate assay for fibrinolytic activity</i>	39
2.3.10 <i>Statistic and data processing</i>	40
2.4 RESULTS	41
2.4.1 <i>Isolation of bacteria</i>	41
2.4.2 <i>Bacterial DNA extraction</i>	42
2.4.3 <i>Bacterial 16S rRNA gene PCR product</i>	42
2.4.4 <i>RFLP analysis</i>	43
2.4.5 <i>Blast</i>	44
2.4.6 <i>Phylogenetic analyses of bacteria producing fibrinolytic enzymes</i>	46
2.4.7 <i>Fibrin plate assay for fibrinolytic activity</i>	48
2.4.8 <i>Serological plate assay for fibrinolytic activity</i>	52
2.5 DISCUSSION	56
2.6 CONCLUSIONS	61
DATA OF REPEATED DETERMINATIONS	62

CHAPTER 3 ESTABLISHMENT OF NK INDUSTRIAL COMPATIBLE AQUEOUS PURIFICATION TECHNIQUES AND THEIR APPLICATION IN IDENTIFICATION OF NEW FIBRINOLYTIC ENZYME PRODUCED BY *BACILLUS*74

3.1 INTRODUCTION.....	74
3.2 MATERIALS AND REAGENTS	83
3.3 METHODS	83
3.3.1 <i>The establishment of the purification protocol</i>	83
3.3.2 <i>Optimization of the purification protocol</i>	87
3.3.3 <i>Amplification of the purification protocol</i>	89
3.3.4 <i>Enzyme Activity Assay</i>	91
3.3.5 <i>Determination of Protein Concentration</i>	91
3.3.6 <i>Purify other NK like fibrinolytic enzyme from Bacillus genera bacteria use the method developed</i>	91
3.3.7 <i>Enzyme identification by MALDI-TOF MS analysis</i>	92
3.3.8 <i>N-Terminal Amino Acid Sequence Analysis</i>	93
3.3.9 <i>Biochemical properties of enzymes characterization</i>	93
3.4 RESULTS	95
3.4.1 <i>The establishment of the purification protocol</i>	95
3.4.2 <i>Optimization of the purification protocol</i>	100
3.4.3 <i>Amplification of the purification protocol</i>	104
3.4.4 <i>Generality test of the protocol</i>	108
3.4.5 <i>Enzyme identification by MALDI-TOF analysis</i>	114
3.4.6 <i>N-terminal amino acid sequence analysis of identified new fibrinolytic enzyme T</i>	117
3.4.7 <i>Biochemical properties of enzymes characterization</i>	120
3.5 DISCUSSION	138
3.6 CONCLUSION	147

CHAPTER 4 IDENTIFICATION OF THE INTRINSIC STABILIZING FACTORS OF NK..148

4.1 INTRODUCTION.....	148
4.2 MATERIALS	157
4.3 METHODS	157
4.3.1 <i>Production of NK</i>	157
4.3.2 <i>The stabilizing effect of low molecular weight compounds determination</i>	158
4.3.3 <i>The stabilizing effect of protein determination</i>	158
4.3.4 <i>The stabilizing effect of lipids determination</i>	158
4.3.5 <i>The stabilizing effect of carbohydrates determination</i>	159
4.3.6 <i>Identification of the specific intrinsic stabilizing factors</i>	160
4.3.7 <i>Determination of the stabilizing ability of the factors</i>	160
4.3.8 <i>Test of the stabilizing ability of the factors by RT-PCR</i>	161
4.4 RESULTS	164
4.4.1 <i>Fibrinolytic activity determination</i>	164
4.4.2 <i>Determination of the stabilizing effect of different compositions of the NK fermentation broths</i>	165
4.4.3 <i>Identification of the specific intrinsic stabilizing factors of NK</i>	166
4.4.4 <i>Determination of the stabilizing ability of the factors</i>	166
4.4.5 <i>Determination of the source of intrinsic stabilizing factors</i>	172
4.5 DISCUSSION	176
4.6 CONCLUSION	180

SUMMARY OF THE RESULTS181

REFERENCES184

APPENDICES.....205

CURRICULUM VITAE.....207