

MASTER'S THESIS

Characterization of hybrid incompatibilities between *Caenorhabditis briggsae* and *C. sp. 9*

Bi, Yu

Date of Award:
2014

[Link to publication](#)

General rights

Copyright and intellectual property rights for the publications made accessible in HKBU Scholars are retained by the authors and/or other copyright owners. In addition to the restrictions prescribed by the Copyright Ordinance of Hong Kong, all users and readers must also observe the following terms of use:

- Users may download and print one copy of any publication from HKBU Scholars for the purpose of private study or research
- Users cannot further distribute the material or use it for any profit-making activity or commercial gain
- To share publications in HKBU Scholars with others, users are welcome to freely distribute the permanent URL assigned to the publication

Abstract

Hybrid incompatibility (HI) refers to lethality, sterility and other reduction in fitness of hybrid progeny between related species which has been frequently observed in different taxa. Its mechanism has been generalized as the consequence of conflicts between genomes of the related species that have been subject to independent evolution. HI plays a critical role in speciation by reducing or preventing exchange of the genetic materials between related species, gradually leading to reproductive isolation. According to the widely accepted Dobzhansky-Muller model, HI is produced by incompatible epistatic interactions between multiple genes that are independently diverged between the parental species. HI has been intensively studied especially in *Drosophila* species. A number of HI loci have been mapped in various species, several of which have been molecularly cloned. However, HI remains poorly understood in other taxa.

My thesis focuses on systematic characterization of HI between two sequenced nematode species, *C. briggsae* and *C. nigoni* (*C. sp. 9*). The former has long been established as a companion species of *C. elegans* for comparative study while the latter is a recently identified species that can mate with the former and produce viable and fertile hybrids in spite of their different reproduction modes, allowing one to genetically and molecularly characterize HI between nematode species for the first time. Such a study is impossible with model organism *C. elegans* as it cannot produce viable hybrids with any other nematode species. To facilitate genome-wide mapping of HI loci between the two nematode species, over 90 dominant visible GFP markers have been randomly integrated into *C. briggsae* chromosomes, permitting repeated backcrossing the marker-associated *C. briggsae* genomic fragments into an otherwise *C. nigoni* background and definitive mapping of HI loci. Genotyping of the introgressions has

been achieved by *C. briggsae* specific PCR, with primers as close as about half a million base pairs away from each other on average.

Using the cost-effective protocol for introgression and genotyping, a genome-wide HI map between the two species has been obtained based on characterization of approximately 100 independent introgression lines. A remarkable proportion of *C. nigoni* genome was found to be replaceable by that from *C. briggsae* as judged by the fact that these introgressions are viable as a homozygote. A few HI loci critical to male inviability and sterility have been narrowed down to small intervals on the X chromosome by contrasting genotypes and HI phenotypes of independent introgression strains. Strains containing these introgressions have been examined for their gonadal structures using fluorescence microscopy. Our study provides genome-wide landscape of HI between nematodes for the first time, allowing comparative studies of HI between nematode and other species.

Acknowledgement

I would like to express my personal appreciation to those who supported and helped me in many different ways. Without them, this thesis will not be possible.

First and the foremost, I would like to express my gratitude to my supervisor, Dr. Zhao Zhongying, for providing me the opportunity to perform this meaningful project and for his patience in guiding me through these two years. Without his professional knowledge and supervision, this thesis would never be accomplished.

I am also very grateful to the Graduate School of Hong Kong Baptist University for my studentship.

I hope to express my sincere gratitude to scientific officers from Department of Biology, HKBU, including Mr. Wai Shing Chung, Ms. Chau Ching Wah, Dr. Law Yu Sheung, and Dr. Ho Yee for their technical assistance.

I am really grateful to Mr. Yan Cheung for giving me the very first introduction to basic laboratory techniques that lead me to the right track of my research. Great thanks to Dr. An Xiaomeng, Mr. Ren Xiaoliang and Mr. Xie Dongying for their contribution in this project, and to Dr. Shao Jiaofang for her technical support in bioinformatics. Thanks were also given to Ms. Wing Sze Ho for proofreading my thesis and Mr. Ming Kin Wong for his assistance in labworks. Once again I would like to thanks to my fellow labmates for their enthusiastic participation and insightful thinking in discussion, which is invaluable for me.

I would like to express my heartfelt appreciation to my friends and family who always keep in touch and support me.

In the end, I truly appreciate the teachers who spent their valuable time in reading my thesis.

Table of contents

Declaration	i
Abstract	ii
Acknowledgement	iv
Table of contents	vi
List of Tables	ix
List of Figures	x
Symbols of Abbreviation	xii
Chapter 1 Introduction	1
1.1 Overview of Hybrid Incompatibility (HI).....	1
1.2 Models explaining hybrid incompatibility.....	3
1.3 Previous studies on HI.....	6
1.4 <i>Caenorhabditis</i> species as an HI model.....	8
1.4.1 Overview of <i>Caenorhabditis</i> species.....	8
1.4.2 HI study in <i>Elegans</i> group.....	9
1.4.3 Approaches used in studying HI between <i>C. briggsae</i> and <i>C. nigoni</i>	13
Chapter 2 Materials and Methods	16
2.1 Worm growth and handling.....	16
2.1.1 Strains used in the thesis.....	16
2.1.2 Medium preparation.....	16
2.1.3 Worm maintenance.....	17

2.1.4	Decontamination of worm culture	17
2.1.5	Strain storage	18
2.2	Generation of chromosomally integrated GFP marker over <i>C. briggsae</i> genome by bombardment	18
2.2.1	Preparation of plasmid pZZ0031 for bombardment	18
2.2.2	Preparation of DNA-coated golden beads	18
2.2.3	Preparation of synchronized <i>cbr-unc-119</i> population.....	19
2.2.4	Microparticle bombardment.....	20
2.2.5	Screening for transgenic <i>C. briggsae</i>	20
2.3	Introgression of the marker-associated <i>C. briggsae</i> genomic fragment into <i>C.</i> <i>nigoni</i> by repeated backcrossing	21
2.3.1	Introgression with autosome-linked marker	21
2.3.2	Introgression with X-linked marker.....	21
2.4	Genotyping introgression boundaries through PCR.....	23
2.4.1	Primer design	23
2.4.2	Single-worm PCR	26
2.4.3	Gel electrophoresis.....	27
2.5	Methods for rendering homozygous introgression.....	30
2.6	Phenotypic characterization of HI loci	30
2.6.1	Embryonic lethality.....	30
2.6.2	Larval arrest	33
2.6.3	Sex ratio and segregation ratio of marker	33

2.6.4	Homozygous/Hemizygous sterility.....	33
2.6.5	Fertility.....	34
Chapter 3 Results.....		36
3.1	Generation of chromosomally integrated <i>C. briggsae</i> transgenic strains.....	36
3.2	Introgression strains and genotyping of introgression boundaries.....	36
3.2.1	Distribution of GFP insertions over <i>C. briggsae</i> genome.....	37
3.2.2	Distribution of introgression size.....	40
3.2.3	Effect of crossing generations on introgression size.....	41
3.3	HI phenotypes between <i>C. briggsae</i> and <i>C. nigoni</i>	43
3.3.1	Pervasive HI effect from introgressions.....	45
3.3.2	Sex-biased pattern of hybrid incompatibility.....	52
3.3.3	Large X effect supported only by fertility data.....	61
Chapter 4 Discussion.....		63
Chapter 5 Conclusion.....		66
Appendices.....		67
References.....		80
Curriculum vitae.....		90