

DOCTORAL THESIS

Development of fluorescent chemosensors based on different signal transduction mechanisms

You, Qihua

Date of Award:
2014

[Link to publication](#)

General rights

Copyright and intellectual property rights for the publications made accessible in HKBU Scholars are retained by the authors and/or other copyright owners. In addition to the restrictions prescribed by the Copyright Ordinance of Hong Kong, all users and readers must also observe the following terms of use:

- Users may download and print one copy of any publication from HKBU Scholars for the purpose of private study or research
- Users cannot further distribute the material or use it for any profit-making activity or commercial gain
- To share publications in HKBU Scholars with others, users are welcome to freely distribute the permanent URL assigned to the publication

Abstract

A series of fluorescent probes based on different signal transduction mechanisms for the detection of Fe^{3+} , Zn^{2+} , histidine and pH was designed and synthesized. Their photophysical properties, binding abilities and the further application in cell imaging were fully evaluated.

Building on the groundwork of our previous study, molecular scaffold **19** has been appended to spirobenzopyran fluorophore to furnish a highly selective and sensitive Zn^{2+} sensor. To broaden the application scope of this trifunctional receptive molecule, **19** was incorporated onto rhodamine, antipyrine and coumarin moieties to give **20**, **21** and **23**, respectively. Probe **20** operative on a chelation-enhanced fluorescence mechanism exhibited highly selective response to Fe^{3+} with 2:1 stoichiometry of **20**- Fe^{3+} complex. However, a possible tendency of probe **20** to hydrolyze induced by Fe^{3+} and the unsuccessful attempt of cell imaging would limit its application scope.

Probe **21** with *O-N-N-N-N*-ligand showed a highly selective and sensitive detection of Zn^{2+} . The probe displayed suppressed response to Cd^{2+} which is the most common interference ion in zinc metal detection. The binding of Zn^{2+} to probe **21** inhibited the photoinduced electron transfer process originating from the lone pair of the nitrogen atom in the antipyrine moiety to quinoline fluorophore. Therefore, a turn-on fluorescent probe was developed. A moderate binding constant with 1:1 stoichiometry of **21**- Zn^{2+} complex was established by fluorescence titration. The binding mechanism was fully explained by ^1H NMR


titration. To our delight, probe **21** was successfully applied for recognizing Zn^{2+} in living cells.

The preparation of probe **23** was achieved by appendage of **19** to coumarin derived fluorophore and the probe exhibited a good selectivity and fluorescent turn-off property to Cu^{2+} . The 1:1 stoichiometry of **23**- Cu^{2+} ensemble can serve as an efficient probe for the detection of histidine and biothiols. In the presence of NEM, the influence of biothiols could be eliminated. Furthermore, this sensing ensemble was also used in the detection of histidine in hard-to-transfect U87MG cells with very low cytotoxicity.


Based on our group's previous work on the spiropyran platform, a novel ratiometric near-infrared pH probe **27** operating on an excited-state intramolecular electron transfer mechanism was developed. The pK_a was calculated to be 5.9 and the ring-opening/ring-closing mechanism triggered by protons was reasonably explained by ^1H NMR titration. However, this spiropyran-based probe was found to be unsuitable for cell imaging.

To continue the innovation of pH sensing and extend its application in bioimaging, a series of ratiometric pH probes **32** and **38** characterized by their high quantum yield working in the NIR range was developed. The appendage of *N,O*-disubstituted hemiaminal ether moiety onto coumarin fluorophore with C=C double bond conferred the sensory material with the ability to display a pH-dependent ratiometric output operating on the ring-opening/ring-closing mechanism. The pK_a of **32** and **38** were 6.9 and 5.8 – 6.0, respectively, which


rendered them suitable for pH measurement in near-neutral and acidic media. A preliminary work of intracellular pH measurement was also conducted and promising results were obtained.




19




20




21




23



27



32



38-H: R = H
38-CF₃: R = CF₃
38-OMe: R = OMe

Acknowledgements

I would like to express my deepest thanks to my supervisor, Prof. Chan Wing Hong. His patient encouragement, guidance and support, enable me to develop an understanding of the subject and to joyously engage in new scientific discovery endeavor. Besides, I would like to thank Prof. Albert Wai-Ming Lee for his valuable advice and guidance throughout my study.

I would like to sincerely thank Dr. Zhu Jianfa, Dr. Li Qinghua, Dr. Li Yinhui and other labmates for their help and suggestions in my research. Collaboration and discussion with them expanded my knowledge in the area of supramolecular chemistry and organic chemistry.

My thanks also go to Ms. Elsie Chan and Ms. Yee Man Lee in the Department of Biology, Hong Kong Baptist University and Dr. Zhu Xiao-Ming in the Department of Imaging and Interventional Radiology, Prince of Wales Hospital, the Chinese University of Hong Kong (CUHK), for assistance in the cell imaging experiments and Dr. Sam C. K. Hau in the Department of Chemistry, CUHK for taking time to perform X-ray experiments.

At last I would like to thank technical staff in the Department of Chemistry for their help in HRMS, NMR, UV-visible and fluorescent spectroscopic services.

Financial support from the Research Committee of the Hong Kong Baptist University (FRG2/11-12/121) is gratefully acknowledged.

Finally, I would like to thank my family for their constant support and understanding during my PhD study.

Table of Contents

Declaration	i
Abstract	ii
Acknowledgements	v
Table of Contents	vi
List of Tables	ix
List of Figures	x
List of Schemes	xx
List of Abbreviation	xxi
Chapter 1 Introduction	1
1.1 Why fluorescence?	1
1.2 Molecular Recognition.....	3
1.3 Signal transducer parts.....	4
1.4 Signal transduction mechanisms.....	14
1.5 Objectives of the thesis work.....	23
1.6 References.....	23
Chapter 2 Rhodamine-based probe for detection of Fe³⁺	28
2.1 Introduction.....	28
2.2 Results and discussion.....	30
2.3 Experimental section.....	37
2.4 Summary.....	41

2.5 References.....	41
Chapter 3 A quinolinyl antipyrine based fluorescence probe for Zn²⁺ and its application in bioimaging.....	44
3.1 Introduction.....	44
3.2 Results and discussion.....	45
3.3 Experimental section.....	59
3.4 Summary.....	62
3.5 References.....	63
Chapter 4 Coumarin-based fluorescent probe for recognition of Cu²⁺ and for fast detection of histidine in hard-to-transfect cells by sensing ensemble approach.....	66
4.1 Introduction.....	66
4.2 Results and discussion.....	68
4.3 Experimental section.....	82
4.4 Summary.....	86
4.5 References.....	87
Chapter 5 Ratiometric spiropyran-based near-infrared fluorescent pH probe.....	92
5.1 Introduction.....	92
5.2 Results and discussion.....	94

5.3 Experimental section.....	110
5.4 Summary.....	115
5.5 References.....	115
Chapter 6 Rational design of fluorescent NIR probes for the ratiometric measurement of intracellular pH	119
6.1 Introduction.....	119
6.2 Results and discussion.....	122
6.3 Experimental section.....	133
6.4 Summary.....	147
6.5 References.....	147
Chapter 7 Summary and prospect of the research	150
Appendices	154
Curriculum Vitae	197