

## DOCTORAL THESIS

### Molecular signaling in the seed dormancy release and germination in arabidopsis

Liu, Yinggao

*Date of Award:*  
2009

[Link to publication](#)

#### General rights

Copyright and intellectual property rights for the publications made accessible in HKBU Scholars are retained by the authors and/or other copyright owners. In addition to the restrictions prescribed by the Copyright Ordinance of Hong Kong, all users and readers must also observe the following terms of use:

- Users may download and print one copy of any publication from HKBU Scholars for the purpose of private study or research
- Users cannot further distribute the material or use it for any profit-making activity or commercial gain
- To share publications in HKBU Scholars with others, users are welcome to freely distribute the permanent URL assigned to the publication

**Molecular Signaling in the Seed Dormancy Release  
and Germination in Arabidopsis**

**LIU Yinggao**

**A thesis submitted in partial fulfillment of the requirements  
for the Degree of  
Doctor of Philosophy**

**Principal supervisor: Prof. Zhang Jianhua**

**Hong Kong Baptist University**

**November 2009**

## Abstract

Seed germination is a complex process. Many signaling molecules and plant hormones participate in the breaking of its dormancy and initiation of its germination. Nitric oxide (NO) has been reported to be involved in breaking seed dormancy but its action mechanism is not clear. In this study we found that a rapid accumulation of NO induced an equally rapid decrease of abscisic acid (ABA) that is required for this action in the seed of Arabidopsis. Results of QRT-PCR and western indicate that the NO-induced ABA decrease correlates with the regulation of *CYP707A2* transcription and the (+)-abscisic acid 8'-hydroxylase (encoded by *CYP707A2*) protein expression. By analyzing *cyp707a1*, *cyp707a2* and *cyp707a3* mutants, we found that *CYP707A2* plays a major role in ABA catabolism during the first stage of imbibition. Fluorescent images demonstrate that NO is released rapidly in the early hours of the endosperm layer during imbibition. Evidently such response precedes the enhancement of ABA catabolism which is required for subsequent seed germination.

H<sub>2</sub>O<sub>2</sub> is known as a signal molecule in plant cells and also plays a role in the regulation of seeds dormancy and germination. We found that H<sub>2</sub>O<sub>2</sub> is involved in the seed dormancy and germination of Arabidopsis by regulating the ABA catabolism and GA biosynthesis during imbibition. Results of QRT-PCR showed that H<sub>2</sub>O<sub>2</sub> up-regulated ABA catabolism gene *CYP707As*, which decreased ABA content during imbibition. Such action required the participation of NO, another signal molecule. Results of QRT-PCR also indicate that H<sub>2</sub>O<sub>2</sub> up-regulate GA biosynthesis at the same time. When ABA catabolism mutant, *cyp707a2*, and over-expressing plant, *CYP707A2-OE*, were used, ABA showed a negative correlation with GA biosynthesis. Exogenously applied GA overrode the inhibition of germination by low concentration of ABA, but the effect was not significant at high concentration of ABA. Together our results

conclude that H<sub>2</sub>O<sub>2</sub> mediates the upregulation of ABA catabolism, possibly through NO, and also the GA biosynthesis. High concentration of ABA also inhibits the GA biosynthesis but a balance of these two hormones jointly controls the dormancy and germination of Arabidopsis seeds.

Function of MAPK pathways on seed germination was also investigated. After analysis 10 *MPAKKs* mutants, we found that two of them (*MKK2* and *MKK10*) are involved in the ABA-regulated seed dormancy and germination. Our results indicate that both *mkk2* and *mkk10* show ABA-insensitive during germination. By measuring some ABA-regulated gene expression (*RD29A* and *RD29B*), *MKK2* and *MKK10* are likely to affect the expression of these genes regulated by ABA to participate in the regulation of seed dormancy and germination. Furthermore, we also found that *MKK2* and *MKK10* mediate ABA-suppressed GA biosynthesis since this suppression was not found in *mkk2* and *mkk10* mutant

In summary, hormones, signaling molecules and MAPK cascades pathways constitute a complex network mediate seed dormancy break and germination.

# Table of Contents

|   |             |
|---|-------------|
| <b>Declaration</b> .....  | <b>i</b>    |
| <b>Abstract</b> .....   | <b>ii</b>   |
| <b>Acknowledgements</b> .....   | <b>iv</b>   |
| <b>Table of Contents</b> .....  | <b>v</b>    |
| <b>List of Figurers</b> .....   | <b>x</b>    |
| <b>List of Abbreviation</b> .....   | <b>xiii</b> |
| <br>  |             |
| <b>Chapter 1 General Introduction</b> .....   | <b>1</b>    |
| 1. 1 Phase of seed germination and dormancy .....   | 2           |
| 1.2 Regulating network of seed germination and dormancy.....  | 5           |
| 1.2.1 Regulation of dormancy and germination by hormones .....  | 6           |
| 1.2.1.1 Regulation of dormancy and germination by ABA.....  | 6           |
| 1.2.1.2 Regulation of dormancy and germination by GA .....  | 7           |
| 1.2.1.3 Regulation of dormancy and germination by<br>brassinosteroids and ethylene. ....                  | 8           |
| 1.2.2 Regulation of dormancy and germination by nitric oxide,<br>nitrate, and active oxygen species ..... | 9           |
| 1.2.3 Regulation of dormancy and germination by proteins. ....  | 9           |
| 1. 3 The functions of ABA .....   | 12          |
| 1.3.1 ABA biosynthesis pathway .....  | 13          |
| 1.3.2. ABA catabolism .....   | 16          |
| 1.4 The NO signaling .....  | 18          |
| 1.5 The ROS signaling .....   | 20          |
| 1.6 The MAPK cascades.....  | 22          |
| 1.6.1 Components of MAPK cascades in plant .....  | 23          |
| 1.6.2 The Role of MAPK Pathways in stress signaling .....   | 24          |

|  |           |
|--|-----------|
| 1.6.3 The role of MAPK pathways in the signaling of plant hormones.....  | 27        |
| 1.6.4. Summary and perspectives of research.....   | 28        |
| <b>Chapter 2 Nitric oxide-induced rapid decrease of ABA concentration is required in breaking seed dormancy in Arabidopsis .....</b> | <b>30</b> |
| 2.1 Introduction.....  | 30        |
| 2.2 Materials and Methods.....   | 33        |
| 2.2.1 Plant materials.....   | 33        |
| 2.2.2 T-DNA insertion line .....   | 33        |
| 2.2.3 Chemical treatments.....   | 34        |
| 2.2.4 Germination assay .....  | 34        |
| 2.2.5 Determination of NO.....   | 34        |
| 2.2.6 Laser scanning confocal microscopy.....  | 35        |
| 2.2.7 Extraction and determination of ABA .....  | 35        |
| 2.2.8 Western-blot analysis .....  | 36        |
| 2.2.9 QRT-PCR analysis.....  | 37        |
| 2.2.10 Generation of <i>CPY707A2</i> over-expressing and complementary plants .....  | 38        |
| 2.2.11 Accession numbers.....  | 38        |
| 2.3 Results .....  | 38        |
| 2.3.1 Effect of NO on seed dormancy .....  | 38        |
| 2.3.2 Effect of ABA biosynthesis and ABA catabolism on seed dormancy .....   | 43        |
| 2.3.3 Dormancy of knockout, complementary and over-expression lines of the <i>CYP707A</i> family.....                                | 45        |
| 2.3.4 <i>CYP707A2</i> gene was involved in NO-induced dormancy break .....   | 50        |
| 2.3.5 NO enhanced the tolerance to ABA in wild type, <i>cyp707a1</i>   |           |

|  |    |
|--|----|
| and <i>cyp707a3</i> mutants but not in <i>cyp707a2</i> mutant during germination .....                   | 54 |
| 2.3.6 Rapidly accumulated NO at the early stage of imbibition was required for seed dormancy break ..... | 56 |
| 2.4 Discussion .....   | 59 |

|   |           |
|---|-----------|
| <b>Chapter 3 H<sub>2</sub>O<sub>2</sub> mediates the regulations of ABA catabolism and GA biosynthesis in Arabidopsis seed dormancy and germination .....</b> | <b>63</b> |
| 3.1 Introduction .....  | 63        |
| 3.2 Materials and Methods .....   | 67        |
| 3.2.1 Plant materials .....   | 67        |
| 3.2.2 T-DNA insertion line .....  | 67        |
| 3.2.3 Chemical treatments .....   | 67        |
| 3.2.4 Germination assay .....   | 68        |
| 3.2.5 Determination of NO .....   | 68        |
| 3.2.6 Extraction and determination of ABA .....   | 68        |
| 3.2.7 Determination of H <sub>2</sub> O <sub>2</sub> .....  | 68        |
| 3.2.8 QRT-PCR analysis .....  | 69        |
| 3.2.9 Generation of <i>CPY707A2</i> over-expressing .....   | 70        |
| 3.2.10 Accession Numbers .....  | 70        |
| 3.3 Results .....   | 71        |
| 3.3.1 H <sub>2</sub> O <sub>2</sub> reduced dormancy of freshly harvested seeds of Arabidopsis .....  | 71        |
| 3.3.2 H <sub>2</sub> O <sub>2</sub> regulated ABA catabolism and GA synthesis during imbibition .....   | 74        |
| 3.3.3 H <sub>2</sub> O <sub>2</sub> needed NO participated in the regulation of ABA catabolism and seed dormancy break .....                                  | 80        |
| 3.3.4 GA biosynthesis was enhanced by H <sub>2</sub> O <sub>2</sub> .....   | 81        |
| 3.3.5 H <sub>2</sub> O <sub>2</sub> mediated GA biosynthesis and ABA suppressed GA  |           |

|  |            |
|--|------------|
| biosynthesis.....  | 85         |
| 3.4 Discussion.....  | 90         |
| <b>Chapter 4 MKK2 and MKK10 regulate ABA-mediated seed germination and root growth in Arabidopsis.....</b> | <b>95</b>  |
| 4.1 Introduction.....  | 95         |
| 4.2 Materials and Methods.....   | 96         |
| 4.2.1 Plant materials.....   | 96         |
| 4.2.3 Physiological assays.....  | 96         |
| 4.2.4 Extraction and determination of ABA.....   | 97         |
| 4.2.5 QRT-PCR analysis.....  | 97         |
| 4.2.6 Stomata aperture assay.....  | 98         |
| 4.2.7 Accession numbers.....   | 98         |
| 4.3 Results.....   | 99         |
| 4.3.1 <i>mkk2</i> and <i>mkk10</i> are insensitive to ABA in seed germination.....                         | 99         |
| 4.3.2 <i>mkk2</i> and <i>mkk10</i> show low dormancy compared with WT in freshly harvested seeds.....      | 99         |
| 4.3.3 <i>mkk2</i> and <i>mkk10</i> are also insensitive to ABA regulation in seedling growth.....          | 102        |
| 4.3.4 <i>mkk2</i> and <i>mkk10</i> do not affect leaf water loss and stomata aperture.....                 | 102        |
| 4.3.5 MKK2 and MKK10 are required for ABA-induced changes in gene expression.....                          | 107        |
| 4.3.6 ABA suppresses GA biosynthesis needs MKK2 and MKK10 participation.....                               | 111        |
| 4.3.7 <i>mkk2</i> and <i>mkk10</i> do not affect ABA biosynthesis and catabolism during imbibition.....    | 111        |
| 4.4 Discussion.....  | 114        |
| <b>Chapter 5 General discussion and conclusion.....</b>  | <b>117</b> |

**Reference:** ..... 123