

DOCTORAL THESIS

Anti-obesity effects of flavonoids and saponins from adzuki bean

Liu, Rui

Date of Award:
2014

[Link to publication](#)

General rights

Copyright and intellectual property rights for the publications made accessible in HKBU Scholars are retained by the authors and/or other copyright owners. In addition to the restrictions prescribed by the Copyright Ordinance of Hong Kong, all users and readers must also observe the following terms of use:

- Users may download and print one copy of any publication from HKBU Scholars for the purpose of private study or research
- Users cannot further distribute the material or use it for any profit-making activity or commercial gain
- To share publications in HKBU Scholars with others, users are welcome to freely distribute the permanent URL assigned to the publication

Abstract

Overweight and obesity are becoming a major public health issue. Pancreatic lipase is a key enzyme to catalyze the hydrolysis of 50-70% of dietary fat in the digestive system. Inhibition of pancreatic lipase activity can block fat absorption in gastrointestinal tract, further control obesity incidence. On the other hand, α -glucosidase is also a key enzyme to hydrolysis polysaccharides and disaccharides into glucose in small intestine. Inhibition of α -glucosidase activity can block digestion and absorption of carbohydrates, further control metabolism disorders such as diabetes and obesity. Food legumes are widely used in people's life with multiple pharmacological activities. The author extracted total phenolics and saponin components from 13 commonly consumed food legumes produced in China, did a systematic comparative study investigating their inhibitory effects against digestive enzymes (pancreatic lipase and α -glucosidase), and screened adzuki bean (*Vigna angularis* L.) as the further target bean. The results showed that the different concentrations of total phenolic extract from adzuki bean (0.25, 0.5, 0.75, 1 mg/mL) inhibited hydrolysis of triolein about 24.1, 24.0, 30.7, and 36.3%, respectively, while different concentrations of total saponin extracts from adzuki bean (0.25, 0.5, 0.75, 1 mg/mL) inhibited hydrolysis of triolein about 15.9, 23.5, 30.1, and 29.2%, respectively. On the other hand, phenolic extract of adzuki bean at the concentration from 0.25 to 1 mg/mL exhibited much more than 80% α -glucosidase inhibitory activity, while saponin extract of adzuki bean exhibited 56.4 to 68.7% α -glucosidase inhibitory activity. Based on the results of pancreatic

lipase inhibitory activity, at the concentration of 1 mg/mL, phenolics extract and saponin extract of adzuki bean (mean 32.5%) and pinto bean (mean 27%) had the stronger inhibitory effects. Moreover, phenolics extract and saponin extract of adzuki bean (mean 79.25%) and pinto bean (mean 72.85%) also had the stronger inhibitory effects on α -glucosidase activity. In addition, adzuki bean is widely used and has diverse application in foods and drugs. In a word, it indicated that adzuki bean is one of the best target beans to further study anti-obesity and anti-diabetes effects via cell and animal models.

Flavonoids and saponins in adzuki bean were obtained and characterized by high performance liquid chromatography with diode array detection and electro spray ionization-tandem multi-stage mass spectrometry in succession. Among 15 compounds identified, four flavonoids (catechin, vitexin-4''-*O*-glucoside, quercetin-3-*O*-glucoside, and quercetin-3-*O*-rutinoside) and six saponins (azukisaponin I, II, III, IV, V, and VI) in adzuki bean were further quantified by external calibration method using high performance liquid chromatography mass spectrometry with the program of time segment and extract ion chromatogram analysis.

Animal model is a good way to intuitively evaluate the anti-obesity effect of adzuki bean. So the anti-obesity effects of adzuki bean in mice fed with a high-fat diet was investigated. ICR female mice were fed with a high fat diet administrated orally with different doses of adzuki bean extracts for eight weeks. Total extract, flavonoids and saponins of adzuki bean enhanced lipolysis (166.1%, 175.6%, and

152.6%, respectively). Compared to the final body weight (33.6 g) of the high-fat diet group, oral administration (300 mg/kg per day) of total extract, flavonoids and saponins of adzuki bean significantly reduced the final body weight of mice, and significantly decreased the adipose tissue accumulation. The adzuki bean intervention also significantly reduced the levels of serum triglyceride, total cholesterol, low density lipoprotein-cholesterol, and liver lipid. Adzuki bean demonstrated the anti-obesity effects on mice fed with a high fat diet, such effects may mediated through the inhibitory effects of flavonoids and saponins from adzuki bean on α -glucosidase and pancreatic lipase activities, and lipolysis enhancement effect of active components from adzuki bean.

Obesity is characterized biologically at the cellular level by as an increase in the number of preadipocytes and an increase in the size of adipocytes differentiated from preadipocytes in adipose tissue. 3T3-L1 cell model was used to reveal the mechanism of anti-obesity effects of adzuki bean in the present study. The results showed that adzuki bean total extract, adzuki bean flavonoids, adzuki bean saponins, four mono flavonoides (catechin, vitexin-4''-O-glucoside, quercetin-3-O-glucoside, and quercetin-3-O-rutinoside) and six mono saponins (azukisaponin I, II, III, IV, V, and VI) exhibited inhibitory effect of proliferation of 3T3-L1 cells, and the inhibitory rate of proliferation of 3T3-L1 cells was about 20%. Mature adipocytes were stained by Oil Red O and the lipid accumulated exhibited red color, while the preadipocytes could not be stained. The data of the effects of adzuki bean samples on lipid accumulation during the differentiation

period of 3T3-L1 cells revealed that azukisaponin II had the highest inhibitory effect (49.72%) with a dose dependent manner from 50 to 500 $\mu\text{g/mL}$. It indicated that adzuki bean may block the differentiation of 3T3-L1 cells from preadipocytes to mature adipocytes which contribute to anti-obesity effects. Most of adzuki bean samples significantly blocked the triglyceride accumulation and enhanced lipolysis by increasing the release of glycerol during differentiation of 3T3-L1 cell. Additionally, adzuki bean samples except catechin significantly inhibited glycerol-3-phosphate dehydrogenase activity in 3T3-L1 cells, which indicated that adzuki bean samples had the ability to inhibit triglyceride synthesis.

Real-time polymerase chain reaction and Western blot methods were used to investigate the intervention effects of adzuki bean total extract, adzuki bean flavonoids, adzuki bean saponins, quercetin-3-*O*-rutinoside, and azukisaponin II isolated from adzuki bean on the related gene expressions. These adzuki bean samples inhibited differentiation of 3T3-L1 cell by decreasing the expression of peroxisome proliferator-activated receptor γ and C/EBP α (major adipocyte transcription factors) at messenger ribonucleic acid level and protein level in adipocytes. Additionally, they inhibited adipogenesis by decreasing the expression of fatty acid synthase, promoting lipolysis by increasing the expression of hormone-sensitive lipase and adipose triglyceride lipase, made energy balance by regulating the expression of leptin, adenosine monophosphate-activated protein kinase, and tumor necrosis factor α , and induced cell apoptosis by increasing the expression of bax and decreasing the expression of B-cell

lymphoma 2. These findings provide insight into the molecular mechanisms through regulation of the related gene expression in 3T3-L1 adipocytes. It also suggested that adzuki bean had a positive effect in prevention and treatment of adipogenesis-related obesity, and might be a good option of functional foods to control obesity.

Acknowledgements

First and foremost, I would like to express my sincere gratitude to my supervisor Prof. Zongwei CAI, and my co-supervisor Dr. Baojun XU for their invaluable advice and support both in lab and life. They provide me a lot of opportunity to explore and develop my potentials in scientific inquiry. Their guidance, inspiration, and continued encouragement are greatly appreciated.

I also would like to thank professors in food science program at UIC for their sincere help to improve this work, which greatly enhanced my understanding on the subjects in theoretical view of experiments.

My sincere thanks must go to my group members for their continued support and helpful discussion.

I also would like to show my appreciation to Prof. Yinan ZHENG and Prof. Wencong LIU for their help in chemical experiments and animal studies.

At last, I also would like to thank my family for their love, support and encouragement.

Table of Contents

Declaration.....	i
Abstract.....	ii
Acknowledgements.....	vii
Table of Contents.....	viii
List of tables.....	xv
List of figures.....	xvi
List of abbreviations.....	xxx
Chapter 1 Introduction.....	1
1.1. Overview on obesity.....	1
1.1.1. Occurrence of obesity.....	1
1.1.2. Causes of obesity.....	2
1.1.3. Consequences of obesity.....	2
1.1.4. Prevention and treatment of obesity.....	2
1.1.5. Pancreatic lipase inhibitors.....	3
1.1.6. α -Glucosidase inhibitors.....	4
1.1.7. 3T3-L1 cell model.....	5
1.2. Anti-obesity effects of food legumes.....	9
1.2.1. Anti-obesity of foods.....	9
1.2.2. Anti-obesity of food legumes.....	10

1.3. Adzuki bean.....	13
1.3.1. Chemical constituents	14
1.3.2. Pharmacological actions	22
1.4. Aims of the project	37
Chapter 2 Inhibitory Effect of Phenolics and Saponins from Commonly Consumed Food Legumes in China on Pancreatic Lipase Activity and α-Glucosidase Activity	43
2.1. Introduction	43
2.2. Materials and methods	45
2.2.1. Materials	45
2.2.2. Chemicals and apparatus	47
2.2.3. Methods	47
2.3. Results and discussion.....	52
2.3.1. Total phenolic content and total saponin content in 13 food legumes ..	52
2.3.2. Effects of total phenolic extracts and total saponin extracts from 13 food legumes on pancreatic lipase activity	54
2.3.3. Effects of total phenolic extracts and total saponin extracts from 13 food legumes on α -glucosidase activity.....	58
2.4. Chapter summary	61
Chapter 3 Characterization and Quantification of Flavonoids and Saponins in Adzuki Bean (<i>Vigna angularis</i> L.) by HPLC-DAD-ESI-MSⁿ Analysis	62

3.1. Introduction	62
3.2. Materials and methods	64
3.2.1. Materials	64
3.2.2. Chemicals and reagents	64
3.2.3. Preparation of saponin standards from adzuki bean	64
3.2.4. Preparation of flavonoids and saponins from adzuki bean	65
3.2.5. High performance liquid chromatography analysis	68
3.2.6. Electro spray ionization-tandem multi-stage mass spectrometry analysis	68
3.3. Results and Discussion	70
3.3.1. Optimization of sample preparation	70
3.3.2. Optimization of HPLC-DAD-ESI-MS ⁿ conditions	70
3.3.3. Analysis of flavonoids in adzuki bean by HPLC-ESI-MS ⁿ	73
3.3.4. Analysis of saponins in adzuki bean by HPLC-ESI-MS ⁿ	84
3.3.5. Quantification of flavonoids and saponins in adzuki bean	91
3.4. Chapter summary	94
Chapter 4 Saponins and flavonoids from adzuki bean (<i>Vigna angularis</i> L.) ameliorate high-fat-diet induced obesity in ICR mice	95
4.1. Introduction	95
4.2. Materials and methods	96
4.2.1. Materials	96
4.2.2. Experimental animals	96

4.2.3. Diets for ICR mice	97
4.2.4. Preparation of flavonoids and saponins from adzuki bean	97
4.2.5. Pancreatic lipase activity assay	99
4.2.6. α -Glucosidase activity assay	99
4.2.7. Lipolysis assay	99
4.2.8. Estimation of relative parameters of mice fed with a high-fat diet	100
4.2.9. Statistical analysis	102
4.3. Results and discussion.....	103
4.3.1. Effects of adzuki bean on α -glucosidase and pancreatic lipase activities in vitro.....	103
4.3.2. Effects of adzuki bean on noradrenaline-induced lipolysis of isolated fat cells from rats	106
4.3.3. Effects of adzuki bean on body weight of mice fed with a high-fat diet for eight weeks.....	108
4.3.4. Effects of adzuki bean on organ weight and adipose tissue weight of obese mice.....	110
4.3.5. Effects of adzuki bean on biochemical parameters of serum and liver of obese mice.....	114
4.4. Chapter summary	117
Chapter 5 Intervention Effects of Flavonoids and Saponins from Adzuki Bean on Proliferation and Differentiation of 3T3-L1 Preadipocyte.....	118
5.1. Introduction	118

5.2. Materials and methods	122
5.2.1. Cell line.....	122
5.2.2. Materials and cell culture media.....	122
5.2.3. MTT proliferation assay	123
5.2.4. Adipogenesis and lipid visual assay by oil red O staining.....	124
5.2.5. Triglyceride content assay	125
5.2.6. Glycerol release	126
5.2.7. Measurement of glycerol-3-phosphate dehydrogenase activity	126
5.2.8. Statistical analysis.....	127
5.3. Results and discussion.....	128
5.3.1. Effect of adzuki bean samples on proliferation of 3T3-L1 preadipocyte	128
5.3.2. Effect of adzuki bean samples on lipid accumulation during differentiation of 3T3-L1 cells.....	133
5.3.3. Effect of adzuki bean samples on triglyceride accumulation in 3T3-L1 cells.....	136
5.3.4. Effect of adzuki bean on lipolysis in differentiated 3T3-L1 adipocytes	139
5.3.5. Effect on glycerol-3-phosphate dehydrogenase activity.....	141
5.4. Chapter summary	144
 Chapter 6 Effects of Flavonoids and Saponins from Adzuki Bean on Preadipocyte Proliferation and Differentiation related Gene Expression....	145

6.1. Introduction	145
6.2. Materials and methods	151
6.2.1. Experimental materials	151
6.2.2. Cell line.....	151
6.2.3. Cell culture and treatment.....	151
6.2.4. Antibodies	151
6.2.5. Primers	151
6.2.6. Western blot analysis	152
6.2.7. Ribonucleic acid preparation and real-time polymerase chain reaction	154
6.2.8. Statistical analysis.....	159
6.3. Results and discussion.....	160
6.3.1. Effects of adzuki bean samples on PPAR γ and C/EBP α mRNA and protein expression in 3T3-L1 adipocytes.....	160
6.3.2. Effects of adzuki bean samples on the expression of FAS in 3T3-L1 adipocytes	166
6.3.3. Effects of adzuki bean samples on the expression of HSL and ATGL for lipolysis in 3T3-L1 adipocytes	169
6.3.4. Adzuki bean samples induced phosphorylation of AMPK	172
6.3.5. Adzuki bean samples induced leptin level in 3T3-L1 adipocytes	175
6.3.6. Adzuki bean samples induced apoptosis in 3T3-L1 adipocytes	178
6.4. Chapter summary	183

Chapter 7 Conclusions and Future Studies	184
References.....	189
Curriculum Vitae.....	219
Outcome of the Thesis Work	220
A. Publications	220
B. Conference and Symposium Presentations.....	221