

DOCTORAL THESIS

Co-composing of sewage sludge with coal fly ash

Fang, Min

Date of Award:
1998

[Link to publication](#)

General rights

Copyright and intellectual property rights for the publications made accessible in HKBU Scholars are retained by the authors and/or other copyright owners. In addition to the restrictions prescribed by the Copyright Ordinance of Hong Kong, all users and readers must also observe the following terms of use:

- Users may download and print one copy of any publication from HKBU Scholars for the purpose of private study or research
- Users cannot further distribute the material or use it for any profit-making activity or commercial gain
- To share publications in HKBU Scholars with others, users are welcome to freely distribute the permanent URL assigned to the publication

1563422x

TH
PH.D
1998FC

Co-Composting of Sewage Sludge with Coal Fly Ash

FANG Min

A thesis submitted in partial fulfillment of the requirements
for the degree of
Doctor of Philosophy

March 1998

Hong Kong Baptist University

Abstract

Composting is becoming a more acceptable and economical method of treating sewage sludge especially for cities with a high population density, such as Hong Kong. However, the major drawback of sewage sludge composting is the high heavy metal content in the end-product, especially for the sewage sludge derived from industrial activities. It is therefore the aim of the present study to develop a cost-effective technique using coal fly ash amendment for sewage sludge composting, which can reduce the availability of heavy metals in the composting products, and improve the decomposition activity during fly ash-sludge composting through microbiological approaches.

The first phase of the present study aims to evaluate the feasibility of using two different alkaline materials, i.e., coal fly ash and lime, as co-composting materials for sewage sludge. Sewage sludge was mixed with sawdust at 2:1 w/w (fresh weight basis) to achieve a C/N ratio of 25. Coal fly ash was amended with the sewage sludge and sawdust mixtures at 0%, 10%, 25%, and 35% w/w (dry weight basis), while amendment rates for lime were 0%, 0.63%, 1.0%, and 1.63% w/w (dry weight basis). The mixtures were composted for 100 days in a laboratory batch composting system and samples were removed periodically for the evaluation of nutrient transformations, maturation, and metal availability.

Addition of coal fly ash at an amendment rate of $\geq 25\%$ caused a significant decrease in the water-soluble and DTPA-extractable Cu, Mn, Ni, Pb, and Zn contents during the composting process. Sludge compost receiving a lime amendment rate of $\geq 1.0\%$ also decreased the water-soluble and DTPA-extractable metal contents, but was less effective than that of coal fly ash. Coal fly ash and lime amendment resulted in a low decomposition activity due to their alkalinity as indicated by the decrease in cumulative weight loss along the composting period, but significant decreases were only observed at sludge compost having 35% fly ash and 1.63% lime amendment. An increase in coal fly ash or lime amendment resulted in a significant reduction in soluble $\text{NH}_4\text{-N}$, $\text{NO}_3\text{-N}$, and $\text{PO}_4\text{-P}$ contents at the end of the composting period as compared to the control. In spite of the reduction in microbial decomposition caused by the alkaline amendment, all treatments reached maturation after 63 days of composting as indicated from the C/N ratio and cress seed germination test results, except for 35% fly ash amendment.

Any adverse effects due to coal fly ash or lime amendment on the decomposition process during sewage sludge composting would rule out their use as co-composting materials. Therefore, biological parameters including respiratory activity, microbial populations (thermophilic and mesophilic bacteria, actinomycetes, and fungi), and enzyme activities (dehydrogenase, alkaline phosphatase, β -glucosidase, and urease) were monitored during alkaline material-amended sludge composting to elucidate their effects on biological activities. Adverse effects of coal fly ash and lime on biological parameters were generally restricted to the early stage of thermophilic phase. Addition of $\leq 25\%$ coal fly ash and $< 1.0\%$ lime could be co-composted with sewage sludge without exerting a significant inhibition on the growth of bacteria, and the activities of β -glucosidase, alkaline phosphatase, and dehydrogenase at the end of composting. This, together with the higher effectiveness in reducing available heavy metal contents without any extension of the maturation period, supported the use of 25% fly ash as an appropriate amendment rate for sewage sludge composting.

Since 25% fly ash amendment exerted adverse effects on the decomposition process at the thermophilic phase of sewage sludge composting, the second phase of the study was to adopt a microbiological approach to improve the composting process. Isolation and identification of thermophilic bacteria from sludge compost with or without ash amendment was conducted according to the Biolog identification tests. *Bacillus* species including *B. brevis*, *B. coagulans*, *B. circulans*, *B. sphaericus*, *B. subtilis*, *B. licheniformis*, *B. thermoglucosidasius*, and *B. stearothermophilus* were identified and only one Gram-negative thermophile was tentatively identified as *Thermus spp.* among all isolates. *B. brevis* was the most dominant species of thermophilic bacteria in sludge or ash-sludge compost which accounted for more than 70% of total thermophilic bacterial population during thermophilic phase. Ash amendment at 25% decreased the population density and species diversity of thermophilic bacteria in sludge compost because of the high pH and salinity. Nevertheless, *B. brevis* still maintained as the most dominant species during the ash-sludge composting process.

Due to the high contents of Cu and Zn in the sludge, attempts were made to isolate Cu- and Zn-tolerant thermophilic bacteria, and inoculate their cultures in 25% fly ash-amended sludge compost in order to accelerate the decomposition activity. Three Cu- and Zn-tolerant *Bacillus* species, i.e., *B. brevis*, *B. coagulans*, and *B. licheniformis*, were isolated. The cultures of these bacilli were inoculated into ash-sludge mixture in aqueous phase and incubated for 10 days. All the three bacilli were able to decompose the organic matter more quickly as compared to the control receiving no inoculation, while no significant difference was noted among the three bacilli at an inoculation size of 10^7 CFU g⁻¹ dry sludge.

In addition to the isolated Cu- and Zn-tolerant thermophilic bacteria, a commercial decomposer and a recycled sludge compost were used as seeding materials in the composting of 25% ash-amended sludge. Inoculation of the commercial decomposer (0.6% dry weight basis) and bacterial culture (10^7 CFU g⁻¹ dry sludge) with milk powder as a substrate (1.8% dry weight basis) could improve the decomposition activity by more than 10% as compared to the control without any additives according to the results of CO₂ evolution and cumulative weight loss. Therefore, the isolated metal-tolerant *Bacillus* species were as efficient as the commercial decomposer to accelerate the decomposition activity during ash-amended sludge composting. Ash-sludge compost inoculated with 15% (dry weight basis) of recycled compost showed a slightly but insignificantly lower decomposition activity than those inoculated with bacterial culture and the commercial decomposer with milk powder. This may be due to the lower degradable organic matter in the recycled compost. Hence, the lower operation cost but an acceptable decomposition efficiency makes recycled compost a promising additive for ash-amended sludge composting.

The present study demonstrates that amendment of 25% coal fly ash in sewage sludge composting is an appropriate option in reducing the availability of heavy metals from sludge compost without affecting the maturation of the end-product. The low effective reduction of available metal contents and the high level of nitrogen loss in lime treatments render lime less attractive than coal fly ash as a co-composting material for sewage sludge. Addition of 15% recycled sludge compost can improve the decomposition activity of 25% ash-amended sludge composting. This co-composting strategy will provide a cost-effective means to produce a high quality of sludge compost, but requires further field tests and plant growth experiments for validation.

Table of Contents

	Page
Declaration	i
Abstract	iii
Acknowledgements	v
Table of Contents	vi
List of Tables	xi
List of Figures	xiv
List of Plates	xxii
List of Appendices	xxiii
Section I. Background	
Chapter 1. Aim and Objectives	1
1.1 Introduction	1
1.2 Research objectives and outline of the present study	3
Chapter 2. Literature Review: Solid Waste Composting	6
2.1 General introduction	6
2.1.1 Definition	6
2.1.2 Objective, benefits and limitations of composting	6
2.2 Basic concepts of composting	8
2.2.1 Composting process	8
2.2.2 Physical and chemical factors affecting composting process	10
2.3 Microbiology of composting	12
2.3.1 Microorganisms of decomposition	14
2.3.2 Pathogens	18
2.3.3 Inoculation	21
2.4 Carbon and nitrogen variation during composting	24
2.4.1 Carbon transformation	24
2.4.2 Nitrogen balance	27
2.5 Trace elements in compost	31

	Page	
2.5.1	Heavy metals in compost	31
2.5.2	Chemical extraction of heavy metals in compost	33
2.5.3	Effect of composting on heavy metals	36
2.5.4	Uptake of heavy metals from compost-amended soil	38
2.6	Evaluation of compost maturation	39
2.6.1	Chemical methods	41
2.6.2	Microbiological tests and activity	45
2.6.3	Plant tests	48
2.7	Sewage sludge management	51
2.7.1	Sewage sludge generation and disposal	51
2.7.2	Sewage sludge characteristics	53
2.7.3	Sewage sludge composting	56
2.8	Coal fly ash properties and disposal	58
2.9	Conclusions	62
Section II.	Co-Composting of Sewage Sludge with Alkaline Materials	
Chapter 3.	Characteristics of Raw Materials for Composting Study	64
3.1	Introduction	64
3.2	Materials and methods	65
3.3	Results and discussion	66
3.3.1	Properties of sewage sludge, coal fly ash and sawdust	66
3.4	Conclusions	69
Chapter 4.	Evaluation of Chemical Parameters during Co-Composting of Sewage Sludge with Alkaline Materials	72
4.1	Introduction	72
4.2	Materials and methods	74
4.2.1	Composting control system	74
4.2.2	Preparation of composting mass	76
4.2.3	Nutrient measurement	77

	Page	
4.2.4	Cress seed germination test	77
4.2.5	Water soluble and DTPA extractable heavy metal contents	79
4.2.6	Statistical analysis	79
4.3	Results and discussion	80
4.3.1	Changes in temperature evolution	80
4.3.2	Changes in pH and EC	82
4.3.3	Changes in ash contents and cumulative weight loss	87
4.3.4	Changes in total organic and soluble organic carbon	90
4.3.5	Changes in various forms of nitrogen	93
4.3.6	Changes in soluble organic and inorganic phosphorus	100
4.3.7	Overall mass balance	103
4.3.8	Compost maturity	103
4.3.9	Water-soluble and DTPA-extractable heavy metals	114
4.3.10	Correlation of results of chemical analysis	124
4.4	Conclusions	124
Chapter 5.	Changes in Biological Parameters during Co-Composting of Sewage Sludge with Alkaline Materials	127
5.1	Introduction	127
5.2	Materials and methods	129
5.2.1	Experimental setup	129
5.2.2	Microbiological examination	129
5.2.3	Enzyme assays	130
5.2.4	Statistical analysis	131
5.3	Results and discussion	132
5.3.1	Changes in CO ₂ evolution	132
5.3.2	Changes in microbial population	135
5.3.3	Enumeration of pathogens	145
5.3.4	Enzyme analysis	146
5.3.5	Correlation test for biological parameters	154
5.4	Conclusions	156

	Page
Section III. System Improvement of the Co-Composting Process of Sewage Sludge with Coal Fly Ash	
Chapter 6. Effect of Coal Fly Ash on Thermophilic Bacteria When Co-Composting Sewage Sludge with Coal Fly Ash	159
6.1 Introduction	159
6.2 Materials and methods	160
6.2.1 Enumeration and isolation	160
6.2.2 Biolog identification tests	161
6.3 Results and discussion	162
6.3.1 Identification of thermophilic bacteria	162
6.3.2 Effect of fly ash on <i>Bacillus</i> species	168
6.4 Conclusions	173
Chapter 7. Digestion Activity of Cu- and Zn-tolerant Thermophilic Bacteria Isolated from Ash-Amended Sewage Sludge Compost	174
7.1 Introduction	174
7.2 Materials and methods	176
7.2.1 Isolation and identification of tolerant thermophilic bacteria	176
7.2.2 Digestion test of isolated <i>Bacillus</i> strains	177
7.2.3 Statistical analysis	179
7.3 Results and discussion	180
7.3.1 Isolation of Cu- and Zn-tolerant species	180
7.3.2 Digestion activity of isolated <i>Bacillus</i> strains in ash-sludge mixture	183
7.4 Conclusions	199
Chapter 8. Seeding Effect on Sewage Sludge Composting with Coal Fly Ash	200
8.1 Introduction	200
8.2 Materials and methods	201

	Page	
8.2.1	Preparation of microbial additives	201
8.2.2	Preparation of composting mass	202
8.2.3	Chemical parameters	205
8.2.4	Bacterial enumeration	205
8.2.5	Statistical analysis	205
8.3	Results and discussion	207
8.3.1	Changes in temperature and CO ₂ evolution	207
8.3.2	Changes in thermophilic bacterial population	211
8.3.3	Changes in cumulative weight loss	214
8.3.4	Changes in pH and EC	216
8.3.5	Changes in C/N ratio	219
8.4	Conclusions	222
Section IV. General Discussion, Conclusions and Recommendations		
Chapter 9.	General Discussion, Conclusions and Recommendations.	224
9.1	Introduction	224
9.2	Efficiency of alkaline materials in reducing the availability of heavy metals	224
9.3	Effect of alkaline materials on nutrient cycling and availability	228
9.4	Effect of alkaline materials on biological properties during composting	229
9.5	Effect of alkaline materials on compost maturation	230
9.6	Thermophilic bacteria in ash-amended sludge compost	230
9.7	Seeding effect on ash-amended sludge composting	232
9.8	Strategy for large scale operation of ash-amended sludge composting	234
References		237
Appendices		266
Curriculum Vitae		274