

DOCTORAL THESIS

Investigation of environmental fate, photo-transformation and metabolism of triclosan

Wu, Jianlin

Date of Award:
2009

[Link to publication](#)

General rights

Copyright and intellectual property rights for the publications made accessible in HKBU Scholars are retained by the authors and/or other copyright owners. In addition to the restrictions prescribed by the Copyright Ordinance of Hong Kong, all users and readers must also observe the following terms of use:

- Users may download and print one copy of any publication from HKBU Scholars for the purpose of private study or research
- Users cannot further distribute the material or use it for any profit-making activity or commercial gain
- To share publications in HKBU Scholars with others, users are welcome to freely distribute the permanent URL assigned to the publication

Investigation of Environmental Fate, Photo- transformation and Metabolism of Triclosan

WU Jianlin

A thesis submitted in partial fulfillment of the requirements

for the degree of

Doctor of Philosophy

Supervisor: Prof. Zongwei CAI

Hong Kong Baptist University

September 2009

Abstract

Triclosan, 5-chloro-2-(2,4-dichlorophenoxy)phenol, is an active ingredient in many household disinfectants and has been extensively used in improving environmental hygiene. The chemical can be found as an antiseptic component in medical products such as hand disinfecting soaps, medical skin creams, dental products and many household cleansers. Triclosan has been found as a contaminant of rivers, lakes, coastal water, breast milk and human urine and so on. In Europe, about 350 tonnes of triclosan are produced annually for commercial applications. Recent studies suggested that triclosan can be undergone cyclization to form 2,8-dichlorodibenzo-*p*-dioxin (2,8-DCDD) in aqueous solution under UV irradiation. In order to understand triclosan and its relative components of photo-transformation and metabolism in environmental water, mouth wash, toothpaste, breast milk, rat urine and plasma comprehensively, we have performed several projects and their main results are shown as below.

In this thesis, triclosan in the waste, river and coastal water samples collected in Hong Kong was analyzed by GC-ITMS. $^{13}\text{C}_{12}$ -triclosan was used as internal standard in the quantitative analysis. Water samples were extracted and cleaned-up with a C18 solid-phase extraction cartridge. The recoveries of triclosan in spiked coastal water at three different concentrations were 83 % to 110 %. The method detection limit was 0.25 ng/l for triclosan in one-liter water, and the relative standard deviation and relative error were less than 11.0 % and 12.3 %, respectively ($n = 3$). The method was successfully applied to analyze water samples collected from rivers, coastal water bodies and wastewater treatment plants at ng/l levels.

The formation of dioxins including 2,7/2,8-dichlorodibenzo-*p*-dioxin (2,7/2,8-DCDD), 2,3,7-trichlorodibenzo-*p*-dioxin (2,3,7-TrCDD), 1,2,8-trichlorodibenzo-*p*-dioxin

(1,2,8-TrCDD), 2,3,7,8-tetrachlorodibenzo-*p*-dioxin (2,3,7,8-TeCDD), 1,2,3,8-tetrachlorodibenzo-*p*-dioxin (1,2,3,8-TeCDD) and chlorinated triclosan were studied in the presence of low concentration of active chlorine in filtered seawater under UV irradiation (365nm) and dark condition. 2,3,7,8-TeCDD was not detected under dark condition when concentration of triclosan was 0.2 mg/ml and active chlorine was 2.0 mg/ml while 2,7/2,8-DCDD, 2,3,7-TrCDD, 1,2,8-TrCDD, 1,2,3,8-TeCDD and chlorinated trilosan were determined. Under the same concentration of triclosan and active chlorine, 2,3,7,8-TeCDD was detected after the UV irradiation from 120 to 960 min. Formation rate and levels of dioxins and chlorinated triclosan were compared for the experiments with different concentration of triclosan and active chlorine.

A GC-MS method was developed for the analysis of 2,8-DCDD in toothpaste and mouthwash. After liquid-liquid extraction of aqueous phase with n-hexane, the extract was cleaned by a silica-bonded C18 solid-phase extraction (SPE) cartridge and analyzed. The detection limit was 0.96 ng/g and 0.83 ng/g for 1.0 g toothpaste and mouthwash, respectively. The recovery of 2,8-DCDD ranged from 87.5% to 104.2%. The relative errors were less than 12.5 %, and the intra-day as well as inter-day precisions represented by RSD were less than 11.2 % and 10.6 %, respectively. The highest concentration of 2,8-DCDD was 56.3 ± 8.5 ng/g in toothpaste and 180.6 ± 19.2 ng/g in mouthwash. Photo-degradation of spiked triclosan to 2,8-DCDD in toothpaste and mouthwash under UV irradiation was performed to validate the formation mechanism of 2,8-DCDD in toothpaste and mouthwash.

A method involving liquid-liquid extraction, gel permeation chromatography clean up and GC-MS analysis was developed for the determination of triclosan in the breast milk. The calibration curve gave a linear dynamic range of 1.0 to 150 ng/g with correlation coefficient of 0.9996 and the recoveries ranged from 86.2 % to 96.6 % for

the spiked milk samples containing triclosan at three different concentrations. The relative standard deviation and relative error were 0.75% to 9.0% and $-7.2 \pm 3.4\%$ to $9.7 \pm 0.9\%$, respectively. The limit of quantification of the method was 0.15 ng/g milk weight or 4.8 ng/g lipid. The method was successfully applied for the determination of triclosan in human breast milk. The results of triclosan concentration in 148 breast milk samples were ranged from not detected to 53.04 ng/g (lipid based).

In vitro metabolic study of triclosan was performed using Sprague-Dawley (SD) rat liver S9 and microsome, while the oral metabolism was investigated on SD rat. Twelve metabolites were identified by using in-source fragmentation from HPLC-APCI/ITMS. Compared to ESI/MS and tandem mass spectrometry that gave little fragmentation for triclosan and its metabolites, the in-source fragmentation under APCI provided intensive fragmentations for structural identifications. The obtained results indicated that the glucuronidation and sulfonation was the major pathway of phase II metabolism and the hydroxylated products were the major phase I metabolites. Glucose, mercapturic acid and cysteine conjugates of triclosan were also observed in the urine samples of rat oral administrated with triclosan. The *in vitro* metabolic rates of triclosan and its major glucuronidation metabolite were determined.

Plasma samples from the SD rats after oral administration of 5 mg/kg triclosan were analyzed by ultra performance liquid chromatography-triple quadrupole mass spectrometry methods for pharmacokinetic study of triclosan. The limit of quantification of the developed analytical method was 10.8 ng/ml. The recovery, accuracy, precision and repeatability were satisfactory. The half time of elimination was 48.5 ± 10.5 hours. Triclosan and its metabolites, including two hydroxylated and sulfated triclosan, one glucuronidated triclosan and one sulfated triclosan were detected in rat plasma.

Table of Contents

Declaration.....	i
Abstract.....	ii
Acknowledgements.....	v
Table of Contents.....	vi
List of Tables	xii
List of Figures.....	xiv
Chapter 1 General introduction.....	1
1.1. Chemical and physical properties	1
1.2. Bioactive property.....	3
1.3. Toxicity of triclosan.....	5
1.4. Contamination and risk assessment of triclosan in biota	6
1.5. Formation of dioxins and chlorophenol from triclosan	12
1.6. Analytical methods	19
1.7. Objective of this project.....	22
Chapter 2 Determination of triclosan in environmental water related to wastewater treatment by using GC-ITMS	25
2.1. Introduction.....	25
2.2. Experimental	26
2.2.1. Reagents and chemicals	26
2.2.2. Water sample collected and pretreatment	28
2.2.3. GC-MS/MS analysis	28
2.2.4. Method development	31

2.2.5. Quality control	32
2.3. Results and discussion	33
2.3.1. GC-MS/MS analysis of triclosan	33
2.3.2. Method development and validation.....	37
2.3.3. Determination of triclosan in environmental water and wastewater treatment	44
2.4. Chapter Summary	48
Chapter 3 Determination of dioxins from photo-transformation of triclosan by using GC-MS and HRGC-HRMS	50
3.1. Introduction.....	50
3.2. Experimental section.....	52
3.2.1. Apparatus	52
3.2.2. Chemicals and reagents.....	55
3.2.3. Procedure of photo-transformation reaction	55
3.2.4. Liquid-liquid extraction	56
3.2.5. Column chromatography clean-up.....	57
3.2.6. GC-MS analysis	57
3.2.7. HRGC-HRMS analysis.....	58
3.2.8. Method validation	59
3.3. Results and discussion	60
3.3.1. Calibration curves, detection limits, recovery, accuracy and precision..	60
3.3.2. Identification of photo-transformation products by using GC-MS and HRGC-HRMS.....	63
3.3.3. Quantitative analysis of photo-transformation products.....	72

3.3.4. Proposed photo-transformation pathways.....	78
3.4. Chapter Summary	82
Chapter 4 Determination of 2,8-DCDD in toothpaste and mouthwash by using GC-MS	
.....	83
4.1. Introduction.....	83
4.2. Experimental.....	84
4.2.1. Chemicals and reagents.....	84
4.2.2. Toothpaste and mouthwash samples.....	84
4.2.3. GC-MS conditions	86
4.2.4. Sample pretreatment	86
4.2.5. Photo-degradation of spiked triclosan in toothpaste and mouthwash.....	87
4.2.6. Method validation	88
4.3. Results and discussion	89
4.3.1. Identification of 2,8-DCDD in toothpaste and mouthwash	89
4.3.2. Calibration curve and limit of quantification.....	91
4.3.3. Recovery and repeatability	93
4.3.4. Determination of 2,8-DCDD in toothpaste and mouthwash samples.....	95
4.4. Chapter Summary	99
Chapter 5 Determination of triclosan in human breast milk by using GC-MS	100
5.1. Introduction.....	100
5.2. Experimental section.....	101
5.2.1. Apparatus	101
5.2.2. Chemicals and reagents.....	101

5.2.3. Sample pre-treatment	102
5.2.4. GC-MS condition.....	106
5.3. Results and discussion	107
5.3.1. Identification of triclosan in milk using GC-MS analysis	107
5.3.2. Calibration curve.....	113
5.3.3. Recovery and detection limits.....	115
5.3.4. Accuracy and precision.....	117
5.3.5. Results of breast milk samples analysis.....	124
5.3.6. Fat content in breast milk.....	134
5.3.7. Triclosan level in breast milk and in fat of breast milk	137
5.4. Chapter Summary	140
Chapter 6 Identification of triclosan metabolites by using in-source fragmentation	
HPLC-APCI/ITMS	141
6.1. Introduction.....	141
6.2. Experimental	142
6.2.1. Chemicals.....	142
6.2.2. <i>In vitro</i> study	143
6.2.3. <i>In vivo</i> study	143
6.2.4. LC-MS analysis	144
6.3. Results and discussion	145
6.3.1. APCI-MS analysis of triclosan and its metabolites with in-source fragmentation	145
6.3.2. Quantitative analysis of triclosan and major metabolites	157
6.4. Chapter Summary	162

Chapter 7 Metabolism and pharmacokinetics studies of triclosan in rat plasma by using	
UPLC-TQMS.....	163
7.1. Introduction.....	163
7.2. Experimental.....	164
7.2.1. Chemicals and animals	164
7.2.2. UPLC-MS conditions	164
7.2.3. Oral administrations and the preparation of plasma samples	165
7.2.4. Method validation	166
7.3. Results and discussion	167
7.3.1. Optimization of conditions of UPLC and mass spectrometry	167
7.3.2. Method validation	168
7.3.3. Pharmacokinetic studies.....	173
7.3.4. Detection of metabolites in rat plasma.....	178
7.4. Chapter Summary	182
General conclusion	183
References.....	187
Curriculum Vitae	200
Outcome of the Thesis Work	201
I. Publications	201
II. Conference and Symposium Presentations	202

III. Scholarship and Awards 204