

DOCTORAL THESIS

Enhanced methane gas generation by reutilization of acidogenic off-gas during two-phase anaerobic digestion of food waste

Yan, Binghua

Date of Award:
2015

[Link to publication](#)

General rights

Copyright and intellectual property rights for the publications made accessible in HKBU Scholars are retained by the authors and/or other copyright owners. In addition to the restrictions prescribed by the Copyright Ordinance of Hong Kong, all users and readers must also observe the following terms of use:

- Users may download and print one copy of any publication from HKBU Scholars for the purpose of private study or research
- Users cannot further distribute the material or use it for any profit-making activity or commercial gain
- To share publications in HKBU Scholars with others, users are welcome to freely distribute the permanent URL assigned to the publication

**Enhanced Methane Gas Generation by Reutilization of Acidogenic
Off-gas during Two-phase Anaerobic Digestion of Food Waste**

YAN Binghua

**A thesis submitted in partial fulfillment of the requirements
for the degree of
Doctor of Philosophy**

Principal Supervisor: Prof. WONG Jonathan W C

Hong Kong Baptist University

January 2015

DECLARATION

I hereby declare that this thesis represents my own work which has been done after registration for the degree of PhD at Hong Kong Baptist University, and has not been previously included in a thesis or dissertation submitted to this or any other institution for a degree, diploma or other qualifications.

Signature: _____

Date: January 2015

ABSTRACT

Mass balance analysis of two-phase AD indicated that off-gas (H_2 and CO_2) produced in acidogenic reactor represent up to 30% of the consumed substrate and under most circumstances, this part of energy was not utilized leading to low overall energy recovery. Hence, the objective of this study was to enhance overall energy recovery during two-phase AD of food waste through reutilization of acidogenic off-gas and to further optimize the processes through manipulating the metabolic pathways and controlling acidogenic off-gas production.

In the first phase, feasibility of reutilizing acidogenic off-gas in methanogenic reactor and contribution of acidogenic off-gas to overall energy recovery was investigated. Acidogenic off-gas diversion increased the methane gas (CH_4 , 0.28 L/g VS_{added}) production up to 38.6%, of which ~8% was contributed by acidogenic off-gas. Both higher hydrolysis rate and COD production were also achieved with off-gas diversion.

Metabolic pathway determines the distributions of intermediate soluble products, which constitute the quality of acidogenic leachate. Therefore, two experiments focusing on manipulating metabolic pathways were performed. Firstly, the effects of four levels of headspace pressures, 6-12 psi (T1), ~3-6 psi (T2), ~3 psi (T3) and ambient pressure (T4) were investigated. Mixed acids metabolic pathways prevailed in all the treatments with butyrate as the single major component. Then, four different levels of H_2 partial pressure (PH_2) were set the next experiment, self-generated PH_2 (T1, control), 80% of H_2 (T2), 60% of H_2 (T3) and 0.04% of H_2 , while the headspace pressure was kept at 3.3 psi. Typical butyrate fermentation pathways dominated in T4 whereas mixed acid fermentation pathways were prevailing in the other three treatments. Because of the improved hydrolysis/acidogenesis and higher quality of acidogenic products, overall CH_4 recovery in T4 (301.0 L/kg VS_{added}) was 44.6% higher than the control.

In Phase III, strategies to enhance acidogenic off-gas production were investigated. First, four types of neutralization modes including daily pH adjustment of leachate to 6.0, methanogenic effluent recirculation, and initial addition of NaOH and lime separately at a dosage of 20.0 and 14.0 g/kg food waste, respectively, were investigated. Obviously, a H_2 production rate of 3.0 and 2.1 L/d with lime and NaOH addition was much higher than 0.7 and 0.4 L/d with effluent recirculation and daily

adjustment, respectively. Also, addition of alkali agents could enhance the COD leaching of food waste, especially with NaOH. A CH₄ production of 11.24 L/d could be attributed to both the elevated leachate quality and the acidogenic off-gas with lime addition. Another experiment investigated the effect of different carbohydrate contents in the substrates on acidogenic H₂ production. Anaerobic hydrolysis of wastes sourced from bakery (T1), Chinese-style restaurant (T2), western-style restaurant (T3) and wet market were performed in LBRs. Food waste collected from western-style restaurant with a carbohydrate content of 69.5% achieved the highest H₂ production of 61.0 L/kg VS_{added}. The highest specific CH₄ production rate at 0.42 L/gVS_{added} was also achieved with western restaurant food waste.

Finally, the possible redirection of fluxes associated with shift of metabolic pathways from the experiment of PH₂ was proposed. Significant increase in the production of butyrate in treatment T4 with PH₂ of 3.3 psi × 0.04% indicated the channeling of electrons towards the production of butyrate. Dynamics of the microbial community were correlated with the distribution of metabolites. In T1 without external gas flushing, lactic acid fermentation was dominant during the initial 7-days. Accordingly, phylotypes affiliated to the genus *Lactobacillus* sp. were detected. A heterlatic fermentation pathway was observed in in both T2 and T4 during first four days, and thereafter the fermentation pathways shifted towards acetate and butyrate as dominant products, which were accompanied by changing the microbial community with phylotypes of *Clostridium* sp. and *Bifidobacterium* sp. becoming dominant.

To conclude, reutilization of acidogenic off-gas by diversion to methanogenic phase is a promising strategy for enhancing overall energy recovery during two-phase AD of food waste. However, improvement of the short-lived acidogenic H₂ production and H₂/CO₂ ratio needs further investigation.

ACKNOWLEDGEMENT

I would like to express my sincere appreciation to my principal supervisor Professor Jonathan W. C. Wong, for his time and generous guidance throughout my PhD study and research. I want to thank him for the support, advice and encouragement he has given me and also for helping me to solve problems I encountered during my research. But what I treasure most is that he taught the way to solve problem rather than solving the problem. I would also like to thank my co-supervisor Professor N. K. Mak, for his valuable suggestion on my study.

I would also like to thank my thesis examiners, Prof. Zong Wei CAI, Prof. Po Keung WONG, Prof. Dr. Michael Nelles, Dr Kwok Keung SHIU, Dr. Jian Wen QIU and Dr. Ka Man LAI. Your thoughtful comments and suggestions promoted this thesis in many aspects, and I also learned a lot.

I would also like to thank Dr. Ammaiyappan Selvam, for his generous support and invaluable discussion on the experimental problem. Further, kind advice and comments on my thesis revision are greatly appreciated. I thank other members of our research group, Dr. K. Murugesan, Dr. S. Y. Xu, Dr. Muiyar, Dr. Mukesh, Dr. G. Y. Zheng, Dr Ankur, Mr. X. Wang, Ms. Y. Zhou, Ms. M.T. Chan, Ms. C. Y. He and Ms. J. Y. Long for their help and encouragement.

My appreciation also goes to Ms. Josephine L.S. Mok, for her excellent technical assistance throughout my study period; and other technical officers including Mr. W. K. Ip, Mr. K. K. Ma, Mr. K. W. Chan, Mr. C. K. Leung, Dr. H. Y. Yeung, Ms. O. C. W. Chan and Ms L. L. H. Ng for their support in using GC-FID, HPLC-PDA, GC-TCD, PCR, Q-PCR and other instruments.

Finally, a special thank goes to my Mum and family members for their unconditional support during the past three years.

TABLE OF CONTENTS

Declaration	i
Abstract	ii
Acknowledgment	iv
Table of Contents	v
List of Tables	x
List of Figures	xi
Abbreviations	xv
CHAPTER ONE	1
OBJECTIVES AND OUTLINE	1
1.1 Background	1
1.2 Research Gaps	5
1.3 Objective and Outline of the Thesis	5
CHAPTER TWO	8
LITERATURE REVIEW	8
2.1 Introduction of Anaerobic Digestion	8
2.1.1 Hydrolysis/acidogenesis and homoacetogenesis	9
2.1.2 Methanogenesis	11
2.1.3 Two-phase anaerobic digestion	18
2.2 Anaerobic Digestion of Food Waste	20
2.2.1 Food waste in Hong Kong	20
2.2.2 Two-phase dry anaerobic digestion of food waste	21
2.3 Energy Recovery and Mass Balance	22
2.3.1 H ₂ and CO ₂ production	23
2.3.2 Methane gas production	26
2.3.3 Enhancement of CH ₄ production	33
2.3.4 Mass balance	45
2.4 Utilization of H ₂ and CO ₂	45
2.4.1 Homoacetogenic pathway	46
2.4.2 Hydrogenotrophic methanogenesis	51
2.5 Further Improvement of Energy Recovery	53
2.5.1 Metabolic pathways	54
2.5.2 Acidogenic biogas production	57
2.5.3 Bioefficiency of different products	60
2.6 Summary	62
CHAPTER THREE	64
RESEARCH METHODOLOGY	64
3.1. Reactor Design	64
3.2 Substrate and Inoculum	65

3.3 Analytical Methods	66
3.3.1 Physicochemical Analyses	66
3.3.2 Phylogenetic analysis.....	68
3.3.3 Functional gene analysis	70
CHAPTER FOUR.....	73
FEASIBILITY OF REUTILIZATION OF ACIDOGENIC OFF-GAS BY DIVERTING TO METHANOGENIC REACTOR DURING TWO-PHASE ANAEROBIC DIGESTION OF FOOD WASTE	73
Abstract.....	73
4.1 Introduction	74
4.2 Materials and Methods.....	76
4.2.1 Substrate and inoculum.....	76
4.2.2 Experimental set up.....	77
4.2.3 Diversion of acidogenic off-gas to UASB	77
4.2.4 Evaluating the contribution of acidogenic off-gas.....	79
4.2.5 Mass balance analysis	79
4.2.6 Analytical methods.....	80
4.3 Results and Discussion.....	80
4.3.1 Enhanced hydrolysis/acidogenesis rate of food waste.....	80
4.3.2 Changes in acidogenic metabolic pathways	83
4.3.3 Increased methane production	86
4.3.4 Mass balance analysis of CH ₄ production	88
4.4 Summary	89
CHAPTER FIVE	91
REGULATION OF ACIDOGENIC METABOLIC PATHWAY DURING TWO- PHASE ANAEROBIC DIGESTION OF FOOD WASTE: EFFECT OF HEADSPACE PRESSURE IN ACIDOGENIC REACTOR.....	91
Abstract.....	91
5.1 Introduction	92
5.2 Experiment Design.....	94
5.3 Materials and Methods.....	94
5.3.1 Substrate and inoculum.....	94
5.3.2 Experimental set up and treatments	95
5.3.3 Analyses	95
5.3.4 Statistical analysis	96
5.4 Results and Discussion.....	96
5.4.1 Pressure profile	96
5.4.2 Performance of LBR.....	97
5.4.3 Metabolic pathways	105
5.4.4 Biogas production	110
5.4.5 Overall performance	114
5.5 Summary	115
CHAPTER SIX	117
REGULATION OF ACIDOGENIC METABOLIC PATHWAY DURING TWO-	

PHASE ANAEROBIC DIGESTION OF FOOD WASTE: EFFECT OF HEADSPACE H₂ PARTIAL PRESSURE AND COMPOSITION.....	117
Abstract.....	117
6.1 Introduction.....	118
6.2 Experimental Design.....	122
6.3 Materials and Methodology.....	122
6.3.1 Substrate and inoculum.....	122
6.3.2 Experimental set-up.....	123
6.3.3 Analytical methods.....	123
6.3.4 Thermodynamic analysis.....	124
6.3.5 Q-PCR.....	125
6.4 Results and Discussion.....	126
6.4.1. General performance of the digesters.....	126
6.4.2 Metabolic pathways and distribution of soluble metabolites.....	130
6.4.3 Functional gene analysis.....	134
6.4.4 Thermodynamics.....	140
6.4.5 Recovery of methane gas.....	141
6.4.6 Overall Performance.....	143
6.5 Conclusion.....	143
CHAPTER SEVEN.....	145
ENHANCED ACIDOGENIC HYDROGEN PRODUCTION DURING TWO PHASE ANAEROBIC DIGESTION OF FOOD WASTE: EFFECT OF DIFFERENT NEUTRALIZATION MODES.....	145
Abstract.....	145
7.1 Introduction.....	146
7.2 Experimental Design.....	149
7.3 Materials and Methodology.....	149
7.3.1 Substrate and seed sludge.....	149
7.3.2 Calculation of H ₂ production.....	150
7.3.3 Analyses.....	151
7.4 Results and Discussion.....	151
7.4.1 Acidogenic H ₂ production.....	151
7.4.2 Distribution of soluble products.....	156
7.4.3 General performance of the digesters.....	163
7.4.4 Overall performacne.....	168
7.5 Conclusion.....	169
CHAPTER EIGHT.....	171
ENHANCED BIOHYDROGEN PRODUCTION DURING TWO-PHASE ANAEROBIC DIGESTION OF FOOD WASTE: EFFECT OF CARBOHYDRATE CONTENTS IN THE SUBSTRATE.....	171
Abstract.....	171
8.1 Introduction.....	172
8.2 Experimental Design.....	174
8.3 Materials and Methods.....	174

8.3.1 Food waste and seed sludge	174
8.3.2 Experimental set up and treatments	175
8.3.3 Calculation of H ₂ production	175
8.3.4 Activity of hydrogenase	176
8.3.5 Analyses	176
8.4 Results and Analysis	177
8.4.1 Characterization of food waste and acidogenic H ₂ production.....	177
8.4.2 Distributions of soluble products	183
8.4.3 Performance of LBR-UASB	188
8.5 Conclusion.....	194
CHAPTER NINE	195
CONTROLLING THE ACIDOGENIC METABOLIC PATHWAYS TOWARDS	
THE PRODUCTION OF TARGET PRODUCTS.....	195
Abstract.....	195
9.1 Introduction	196
9.2 Materials and Methods.....	198
9.2.1 Experimental setup and operation.....	198
9.2.2 End product analysis	199
9.2.3 Microbial community analysis.....	200
9.3 Results and Discussion.....	201
9.3.1 End-product analysis.....	201
9.3.2 Dynamics of microbial community	202
9.3.3 Ordination analysis	213
9.3.4 Redirection of carbon and electron flux.....	217
Summary	220
CHAPTER TEN.....	222
CONCLUSIONS AND RECOMMENDATIONS FOR FURTHER RESEARCH	
.....	222
10.1. Introduction.....	222
10.2 Feasibility of Diversion of Acidogenic (LBR) Off-gas to Methanogenic UASB for Reutilization	226
10.3 Optimization through Manipulation of Metabolic Pathways in Acidogenic Reactor	227
10.4. Enhanced CH ₄ Recovery through Controlling Acidogenic Biogas Production	228
10.5 Dynamics of Microbial Community and Proposed Redirection of Electron Flux	229
10.6 Recommendations for Further Research.....	230
REFERENCES	232
CURRICULUM VITAE	294