

DOCTORAL THESIS

Synthesis, characterization and photophysical properties of metal complexes with schiff-base and porphyrin ligands

Zhang, Jie

Date of Award:
2012

[Link to publication](#)

General rights

Copyright and intellectual property rights for the publications made accessible in HKBU Scholars are retained by the authors and/or other copyright owners. In addition to the restrictions prescribed by the Copyright Ordinance of Hong Kong, all users and readers must also observe the following terms of use:

- Users may download and print one copy of any publication from HKBU Scholars for the purpose of private study or research
- Users cannot further distribute the material or use it for any profit-making activity or commercial gain
- To share publications in HKBU Scholars with others, users are welcome to freely distribute the permanent URL assigned to the publication

**Synthesis, Characterization and Photophysical Properties of
Metal Complexes with Schiff-Base and Porphyrin Ligands**

ZHANG Jie

**A thesis submitted in partial fulfillment of the requirements
for the degree of
Doctor of Philosophy**

Principal Supervisor: Prof. WONG Wai Kwok

Hong Kong Baptist University

June 2012

Abstract

A series of a new class of D- π -A conjugated nonsymmetrical metallosalophen complexes have been synthesized and characterized. The two photon absorption cross-section $\sigma^{(2)}$ reached level as high as to 1754 GM, as measured by a femtosecond open-aperture Z-scan, which is among the highest value known for metallosalophen complexes to date. The similar $\sigma^{(2)}$ values 1465 GM of symmetric metallosalophen indicates the attachment of D and A fragments doesn't have much effect on electronic properties of this kind of nonsymmetrical metallosalophen complexes. DFT calculations also show the two photon absorption is slightly affected by D and A fragments. Such kind of D- π -A conjugated metallosalophen complexes can be further modified along the Y axis. These salophen complexes with moderate fluorescence quantum yields, larger two photon absorption cross-sections $\sigma^{(2)}$ and high stability are potential two-photon fluorophores applicable in tissue imaging which can be excited by NIR or longer wavelengths to increase penetration depth and reduce cell damage.

We synthesized a series of new di-nuclear Schiff-base (**L**) Complexes **ZnLnL** (**Ln**=La, Gd, Nd, Yb, Er, Eu, Tb, Ho) heterometallic iodine substituted Schiff-base complexes. Some of these complexes have been structurally characterized. The structures of **ZnEuL** and **ZnNdL** were ascertained by X-ray crystallography. Effective energy transfer was observed from the triplet state of the **ZnL** complex to the excited

state of lanthanide(III) ions, which emitted in the visible (Ln = Tb, Ho) and NIR (Ln = Nd, Er, Yb) region. The NIR luminescence of lanthanide ions can be enhanced by putting a heavy-atom such as iodine at the phenyl rings. The Eu(III) centered emission was quenched possibly because of the electron transfers from excited state of **ZnL** to the Eu^{3+} .

A series of platinum(II) Schiff-base complexes were synthesized and fully characterized. The low-energy absorption bands (400-600 nm) of the complexes were assigned to metal-to-ligand charge-transfer (MLCT): $^1[\text{Pt}(5d) \rightarrow \pi^*(\text{L})]$ mixed with $^1[\text{L}(\text{phenoxide}) \rightarrow \pi^*(\text{imine})]$ transitions. These complexes showed high emission quantum yields and good thermal stability. HOMO (-5.09 to -5.40 eV) and LUMO (-3.26 to -3.43 eV) energy levels had been determined by cyclic voltammetry. The energy gaps of these complexes were from 1.66-2.13 eV. Yellow and red organic light-emitting devices (OLEDs) were fabricated using these complexes as emitting materials. The maximum luminous efficiency, power efficiency, and brightness for the complexes doped OLEDs were 23 cd/A, 17 lm/W and 11106 cd/m^2 , respectively. Aggregation was successfully inhibited and the emission was not changed even at high concentrations in OLEDs device. The WOLEDs is further fabricated with maximum external quantum efficiency of 7.7%, maximum luminance efficiency of 21 cd A^{-1} and maximum power efficiency of 17.7 lm W^{-1} . This phenomenon indicates that these

platinum(II) Schiff-base complexes are good candidates for phosphorescence materials used in OLED devices.

We have synthesized porphyrin-salen-porphyrin triads (**H₆L³**, **ZnL³**, **H₆L⁴**, **ZnL⁴**, **H₂L⁵** and **ZnL⁵**). These products were characterized by ¹H NMR, IR, MALDI-TOF-MS and elemental analysis. The UV-visible absorption results of **H₆L³**, **ZnL³**, **H₆L⁴**, **ZnL⁴**, **H₂L⁵** and **ZnL⁵** supported that the porphyrin and Schiff-base moiety exist a profound electronic communication in the ground state. Comparing to the triads of **H₆L³**, **ZnL³**, **H₆L⁴** and **ZnL⁴**, the emission and absorption peak of **H₂L⁵** and **ZnL⁵** were red-shifted. This is probably due to the *meso*-position linked porphyrin-salen-porphyrin triads have a better conjugation than β -position linked porphyrin-salen-porphyrin triads. Energy transfer studies showed that the energy was transferred from Schiff-base (donor) to porphyrin (acceptor) effectively. HOMO (-4.95 to -5.31 eV) and LUMO (-3.25 to -3.40 eV) energy levels had been determined by cyclic voltammetry, which showed these products were low band gap materials. Furthermore, the highest PCE of 0.32% was achieved for **ZnL³** with a $V_{oc} = 0.65$ V, $J_{sc} = 1.72$ mA cm⁻² and $FF = 0.29$ under illumination of an AM 1.5 solar cell simulator in a 1000 rpm spinning speed. The light energy absorbed by the triads/PCBM blend film was to some extent converted into electricity. The present work demonstrates the potential application of porphyrin-salen-porphyrin triads for visible and near-infrared light power generation. However, the absorption

bands (Soret band) of these triads at 450 nm are narrow, and the Q-bands are still weak in the film absorption spectra. Therefore, the photovoltaic efficiency could be increased by improving the absorption property of porphyrin containing triads in the visible and near-infrared region.

Table of contents

Declaration	i
Abstract	ii
Acknowledgements	vi
Table of contents	viii
List of Tables	xiii
List of Figures	xvi
List of Schemes	xxvi
List of Symbols and Abbreviation	xxvii
Chapter 1	
Introduction	1
1.1 Schiff-Base.....	1
1.1.1 Background.....	1
1.1.2 Schiff -Base Metal Complexes.....	2
1.1.2.1 Schiff-Base Transition Metal Complexes Used in Nonlinear Optics....	3
1.1.2.2 Schiff-Base Transition Metal Complexes Used in Organic Light-emitting Diodes.....	7
1.1.3 Schiff-Base Lanthanide Metal Complexes	12
1.2 Porphyrin.....	16

1.2.1	Background.....	16
1.2.2	Porphyrin Used in Bulk Heterojunction Solar Cells.....	16
1.3	Object of the Thesis.....	20
1.4	References.....	23
Chapter 2	Synthesis and Two-photon Absorption Properties of Unsymmetric	
	Metallosalophen Complexes.....	34
2.1	Introduction.....	34
2.2	Results and Discussion.....	36
2.2.1	Synthesis of Unsymmetric Metallosalophen Complexes.....	36
2.2.2	¹ H NMR of Unsymmetric Metallosalophen Complexes.....	37
2.2.3	IR and MS of Unsymmetric Metallosalophen Complexes.....	42
2.2.4	Thermal Stability of Unsymmetric Metallosalophen Complexes.....	47
2.2.5	Photophysical Properties of Unsymmetric Metallosalophen Complexes.....	48
2.2.6	Theoretical Calculations of Unsymmetric Metallosalophen Complexes.....	58
2.3	Conclusion.....	62
2.4	Experimental Section.....	63
2.5	References.....	80

Chapter 3	Heterometallic Zn(II)- Ln(III) Iodine- Substituted Schiff-Base Complexes: Synthesis, Structures and Photoluminescence.....	87
3.1	Introduction.....	87
3.2	Results and Discussion.....	88
3.2.1	Synthesis of Schiff-Base Ligand and Its Complexes.....	88
3.2.2	¹ H NMR of Schiff-Base Ligand and Its Complexes.....	90
3.2.3	IR and MS of Schiff-Base Ligand and Its Complexes.....	93
3.2.4	Crystal Structures of ZnEuL and ZnNdL.....	97
3.2.5	Photophysical Properties of the Schiff-Base Ligand and Its Complexes.....	101
3.3	Conclusion.....	113
3.4	Experimental Section.....	114
3.5	References.....	121
Chapter 4	Efficient Organic Light-emitting Devices Based on Phosphorescent Platinum(II) Schiff-Base Complexes.....	126
4.1	Introduction.....	126
4.2	Results and Discussion.....	129
4.2.1	Synthesis of Platinum(II) Schiff-Bases Complexes.....	129
4.2.2	¹ H NMR of Schiff-Base Ligands and Platinum(II) Complexes.....	133

4.2.3	IR and MS of Schiff-Base Ligands and Platinum(II) Complexes.....	136
4.2.4	UV-visible Absorption of Schiff-Base Ligands and Platinum(II) Complexes.....	141
4.2.5	Emission of Schiff-Base Ligands and Platinum(II) Complexes.....	144
4.2.5	Electrochemical Properties of Schiff-Base Ligands and Platinum(II) Complexes.....	148
4.2.6	Thermal Stability of Platinum(II) Schiff-Base Complexes.....	151
4.2.7	OLED Devices Performance of Platinum(II) Schiff-Base Complexes.....	152
4.3	Conclusion.....	172
4.4	Experimental Section.....	173
4.5	References.....	189
Chapter 5	Porphyrin-Salen-Porphyrin Triads: Synthesis, Photophysical Properties and Performance in Photovoltaic Device.....	194
5.1	Introduction.....	194
5.2	Results and Discussion.....	197
5.2.1	Synthesis of Porphyrin-Salen-Porphyrin Triads.....	197
5.2.2	¹ H NMR of Porphyrin-Salen-Porphyrin Triads.....	202

5.2.3	IR and MS of Porphyrin-Salen-Porphyrin Triads.....	205
5.2.4	UV-visible absorption of Porphyrin-Salen-Porphyrin Triads.....	209
5.2.5	Fluorescence of Porphyrin-Salen-Porphyrin Triads.....	214
5.2.6	Electrochemical Properties of Porphyrin-Salen-Porphyrin Triads...	219
5.2.7	Photovoltaic Behavior of Porphyrin-Salen-Porphyrin Triads.....	224
5.3	Conclusion.....	229
5.4	Experimental Section.....	230
5.5	References.....	253
Chapter 6	Summary and Future work.....	261
Appendix		265
Curriculum Vitae		303