

## A Diffusion Perturbed Risk Process with Stochastic Return on Investments

Yin, Chuancun; CHIU, Sung Nok

*Published in:*  
Stochastic Analysis and Applications

*DOI:*  
[10.1081/SAP-120028594](https://doi.org/10.1081/SAP-120028594)

Published: 01/03/2004

[Link to publication](#)

*Citation for published version (APA):*  
Yin, C., & CHIU, S. N. (2004). A Diffusion Perturbed Risk Process with Stochastic Return on Investments. *Stochastic Analysis and Applications*, 22(2), 341-353. <https://doi.org/10.1081/SAP-120028594>

### General rights

Copyright and intellectual property rights for the publications made accessible in HKBU Scholars are retained by the authors and/or other copyright owners. In addition to the restrictions prescribed by the Copyright Ordinance of Hong Kong, all users and readers must also observe the following terms of use:

- Users may download and print one copy of any publication from HKBU Scholars for the purpose of private study or research
- Users cannot further distribute the material or use it for any profit-making activity or commercial gain
- To share publications in HKBU Scholars with others, users are welcome to freely distribute the permanent publication URLs

# A DIFFUSION PERTURBED RISK PROCESS WITH STOCHASTIC RETURN ON INVESTMENTS

**Chuancun Yin**

Department of Mathematics, Qufu Normal University,

Shandong 273165, P. R. of China

and Department of Mathematics, Hong Kong Baptist University

Kowloon Tong, Hong Kong, P. R. of China

e-mail: chuancunyin@sina.com

**S. N. Chiu\***

Department of Mathematics, Hong Kong Baptist University,

Kowloon Tong, Hong Kong, P. R. of China

e-mail: snchiu@math.hkbu.edu.hk

## ABSTRACT

A general risk model that allows for stochastic return on investments as well as perturbation by diffusion is studied. Integro-differential equations for the distributions of the time of ruin, the surplus prior to ruin and the deficit at ruin of this model are established. In particular, we consider a diffusion perturbed risk model with interest force in details.

*Keywords:* Risk process; Time of ruin; Ruin probability; Stochastic return

---

\* Corresponding author.

## 1. INTRODUCTION

Paulsen (1) introduced the following risk process  $U_t$  that allows for a stochastic rate of return on investments as well as a stochastic rate of inflation:

$$U_t = \frac{\mathcal{E}(R)_t}{\mathcal{E}(I)_t} \left( u + \int_0^t \frac{\mathcal{E}(I)_{t-}}{\mathcal{E}(R)_{t-}} dP_s \right).$$

The notation  $\mathcal{E}(A)$  denotes the Doléans-Dade exponential of  $A$  given as the solution of the stochastic growth equation  $\mathcal{E}(A)_t = 1 + \int_0^t \mathcal{E}(A)_{s-} dA_s$ , and  $P_t, I_t$  and  $R_t$  are all semi-martingales representing the surplus generating process, the inflation generating process and the return on investment generating process, respectively. The initial values are  $P_0 = u, I_0 = 0$  and  $R_0 = 0$ . Paulsen and Gjessing (2) simplified the model above by assuming that there is no inflation and both the surplus generating process  $P_t$  and the return on investment generating process  $R_t$  are the classical risk processes perturbed by Brownian motions. Paulsen (3) considered a risk process  $U_t$  given by

$$U_t = u + P_t + \int_0^t U_{s-} dR_s, \text{ with } P_0 = R_0 = 0, \quad (1.1)$$

where  $P_t$  and  $R_t$  are independent Lévy processes. Using the above notation (1.1) can be written as  $U_t = \mathcal{E}(R)_t(u + \int_0^t \mathcal{E}(R)_{s-}^{-1} dP_s)$ . For further references see (4).

The present paper generalizes the model given in (1.1) by considering that  $P_t$  is a general jump-diffusion process, whilst  $R_t$  is still a Lévy process. The organization of the paper is as follows. Section 2 presents a precise formulation of our model. Integro-differential equations for the distributions of time of ruin, surplus prior to ruin and deficit at ruin are established in Section 3. The last section studies the model with interest force.

## 2. THE MODEL

Assume that all processes and random variables are defined on the probability space  $(\Omega, \mathcal{F}, \mathcal{F}_t, P)$ , where  $\mathcal{F}_t$  is right-continuous and P-complete. Let  $\mathbb{R}$  and  $\mathbb{R}^+$  denote the real line and the non-negative half line endowed with the Borel  $\sigma$ -algebra  $\mathcal{B}$  and  $\mathcal{B}^+$ , respectively. A risk process can be described as a marked point process  $\{T_n, Z_n\}_{n \geq 1}$ , where  $T_i$ 's are non-negative and represent the times when claims occur, and the marks  $Z_i$ 's are the corresponding claim amounts, which are assumed to be i.i.d. and positive. In risk theory (e.g. (5)), one often defines the aggregate claim amount at time  $t$  as  $S_t = \sum_{i=1}^{\infty} \mathbf{1}(T_i \leq t) Z_i$ . Møller (6, 7) considered a more general form of the aggregate claim amount, and introduced the risk reserve model of the form

$$V_t = V_0 + \int_0^t b(s, V_s) ds - \int_0^t \int_0^{\infty} f(s, z) N(ds, dz), \quad (2.1)$$

where  $b : \mathbb{R}^+ \times \mathbb{R} \rightarrow \mathbb{R}$  is a piecewise continuous function representing the premium income,  $f : \mathbb{R}^+ \times \mathbb{R}^+ \rightarrow \mathbb{R}^+$  a Borel measurable function,  $N(t, D) := \sum_{i=1}^{\infty} \mathbf{1}(T_i \leq t, Z_i \in D)$  a locally finite (not necessarily Poisson) counting measure that counts the number of the claims over  $(0, t]$  with claim amounts taking values in  $D \in \mathcal{B}^+$ . The last term in the right-hand side of (2.1) represents the total amount of claims in the time interval  $(0, t]$ , which can be written as  $\sum_{i=1}^{N(t, \mathbb{R}^+)} f(T_i, Z_i)$ , where each claim amount may be multiplied by a discount factor.

We define the risk process  $U_t$  as the solution of the stochastic differential equation

$$\begin{aligned} U_t = u + \int_0^t b(s, U_{s-}) ds + \int_0^t a(s, U_{s-}) dB_s \\ - \int_0^t \int_0^{\infty} f(s, U_{s-}, z) N(ds, dz) + \int_0^t U_{s-} dR_s, \quad t \geq 0, \end{aligned} \quad (2.2)$$

where  $u$  is the initial assets,  $\{B\}_{t \geq 0}$  is a standard Brownian motion independent of the inhomogeneous Poisson counting measure  $N(\cdot, \cdot)$  and  $a, b : \mathbb{R}^+ \times \mathbb{R} \rightarrow \mathbb{R}$ ,  $f : \mathbb{R}^+ \times \mathbb{R} \times$

$\mathbb{R}^+ \rightarrow \mathbb{R}^+$  are measurable functions satisfying the condition that for positive  $m$  and  $M$ , there exists a positive constant  $K(m, M)$  such that for  $t \leq m$  and  $|x|, |y| \leq M$ ,

$$\begin{cases} |a(t, x) - a(t, y)| + |b(t, x) - b(t, y)| \\ \quad + \int_0^\infty |f(t, x, u) - f(t, y, u)| \lambda(t, du) \leq K(m, M)|x - y|, \\ |a(t, x)| + |b(t, x)| + \int_0^\infty |f(t, x, u)| \lambda(t, du) \leq K(m, M)(1 + |x|), \end{cases} \quad (2.3)$$

where  $\lambda(t, D)$  denotes the compensator of the point process  $\{N(t, D)\}$ . The process  $\{R_t\}_{t \geq 0}$  is a Lévy process independent of  $\{P_t\}_{t \geq 0}$  and having the Lévy–Itô decomposition (see (3))

$$R_t = rt + \sigma_R B_{R,t} + \int_0^t \int_{-\infty}^\infty x(\mu_R(ds, dx) - \mathbf{1}(|x| \leq 1)K_R(dx)ds),$$

where  $r$  and  $\sigma_R$  are positive constants,  $\{B_{R,t}\}_{t \geq 0}$  is another standard Brownian motion independent of  $\mu_R$  which is the Poisson counting measure associated with the jumps of  $R_t$ ,  $K_R(dx)ds$  is the compensator of  $\mu_R(ds, dx)$  and such that  $K_R(\{0\}) = 0$  and  $\int_{-\infty}^\infty \min(1, x^2)K_R(dx) < \infty$ . For simplicity, assume that  $K_R((-\infty, -1]) = 0$ .

Because the quadratic variational processes of  $\int_0^t a(s, U_{s-})dB_s + \int_0^t \sigma_R U_{s-}dB_{R,s}$  and  $\int_0^t \sqrt{a^2(s, U_{s-}) + \sigma_R^2 U_{s-}^2}dW_s$  are the same, where  $\{W\}_{t \geq 0}$  is a standard Brownian motion independent of the compound Poisson processes involved, by Ikeda and Watanabe [(8), p.85], they have the same distributions. Thus, in distribution, we have

$$\begin{aligned} U_t = u &+ \int_0^t (b(s, U_s) + rU_s)ds \\ &+ \int_0^t \sqrt{a^2(s, U_{s-}) + \sigma_R^2 U_{s-}^2}dW_s - \int_0^t \int_0^\infty f(s, U_{s-}, z)N(ds, dz) \\ &+ \int_0^t \int_{-\infty}^\infty xU_{s-}(\mu_R(ds, dx) - \mathbf{1}(|x| \leq 1)K_R(dx)ds). \end{aligned} \quad (2.4)$$

If  $a(t, x) = a(t)$ ,  $b(t, x) = b(t)$  and  $f(t, x, z) = f(t, z)$ , then the solution of (2.2) is

$$U_t = \mathcal{E}(R)_t \left( u + \int_0^t \mathcal{E}(R)_{s-}^{-1} dQ_s \right),$$

where

$$Q_t = u + \int_0^t b(s)ds + \int_0^t a(s)dB_s - \int_0^t \int_0^\infty f(s, z)N(ds, dz),$$

$$\mathcal{E}(R)_t = e^{R_t - \frac{1}{2}\sigma_R^2 t} \prod_{0 \leq s \leq t} (1 + R_s - R_{s-})e^{-(R_s - R_{s-})}.$$

A natural generalization of (2.4) is the time inhomogeneous risk process obtained by replacing the integrands of the first two integrals in the right-hand side of (2.4) by two general measurable functions, say,  $q$  and  $\sigma$ , i.e.

$$U_t = u + \int_0^t q(s, U_s)ds + \int_0^t \sigma(s, U_s)dW_s - \int_0^t \int_0^\infty f(s, U_{s-}, z)N(ds, dz)$$

$$+ \int_0^t \int_{-\infty}^\infty xU_{s-}(\mu_R(ds, dx) - \mathbf{1}(|x| \leq 1)K_R(dx)ds). \quad (2.5)$$

If  $q, \sigma$  and  $f$  satisfy condition (2.3), then  $U_t$  defined by (2.5) is a strong Markov process (see e.g. (9), Theorem 32).

Using Itô's formula, we can prove that the infinitesimal generator  $\mathcal{L}$  of  $\{U_t\}_{t \geq 0}$  in (2.5) is given by

$$\mathcal{L}_{t,y}g(t, y) = \frac{\partial g}{\partial t} + \frac{1}{2}\sigma^2(t, y)\frac{\partial^2 g}{\partial y^2} + q(t, y)\frac{\partial g}{\partial y} + \int_0^\infty [g(t, y - f(t, y, z)) - g(t, y)]\lambda(t, dz)$$

$$+ \int_{-\infty}^\infty [g(t, y + yx) - g(t, x) - \mathbf{1}(|x| \leq 1)xy\frac{\partial g}{\partial y}]K_R(dx). \quad (2.6)$$

### 3. INTEGRO-DIFFERENTIAL EQUATIONS

Let  $T(s) = \inf\{t \geq s : U_t < 0\}$  with the usual convention that  $\inf \emptyset = \infty$ . For  $\alpha \geq 0$ , denote  $\psi_1(s, u) = \mathbb{P}(T(s) < \infty | U_s = u)$ ,  $\psi_2(s, u, t) = \mathbb{P}(T(s) < t | U_s = u)$ ,  $\psi_3(s, \alpha, u) = \mathbb{E}[e^{-\alpha T(s)} | U_s = u]$ ,  $H_1(s, u; x, y) = \mathbb{P}(T(s) < \infty, U_{T(s)-} \leq x, -U_{T(s)} \leq y | U_s = u)$ ,  $H_2(s, u; x, y, t) = \mathbb{P}(T(s) < t, U_{T(s)-} \leq x, -U_{T(s)} \leq y | U_s = u)$ , and  $H_3(s, u; x, y) = \mathbb{E}(e^{-\alpha T(s)} \mathbf{1}(T(s) < \infty, U_{T(s)-} \leq x, -U_{T(s)} \leq y) | U_s = u)$ . We will derive integro-differential equations for  $H_i$ 's.

**Theorem 3.1.** *Assume that  $H_i$ 's have bounded continuous derivatives with respect to  $t$  and  $s$  and bounded continuous first and second derivatives with respect to  $u \geq 0$ . We have*

(i)  $H_1(s, u) := H_1(s, u; x, y)$  satisfies  $\mathcal{L}_{s,u}H_1(s, u) = 0$  in  $(0, \infty) \times (0, \infty)$ , together with the boundary conditions, for  $s, x, y > 0$ ,

$$\begin{aligned} H_1(s, u; x, y) &= \mathbf{1}(-y \leq u), & u \leq 0, \\ H_1(s, u; x, y) &\rightarrow 0, & u \rightarrow \infty. \end{aligned}$$

(ii)  $H_2(s, u, t) := H_2(s, u; x, y, t)$  satisfies  $-\frac{\partial H_2(s, u, t)}{\partial t} + \mathcal{L}_{s,u}H_2(s, u, t) = 0$  in  $(0, \infty) \times (0, \infty) \times (s, \infty)$ , together with the boundary and initial conditions, for  $s, x, y > 0$ ,

$$\begin{aligned} H_2(s, u; x, y, t) &= \mathbf{1}(-y \leq u), & t > s, u \leq 0, \\ H_2(s, u; x, y, t) &\rightarrow 0, & t > s, u \rightarrow \infty, \\ H_2(s, u; x, y, t) &\rightarrow 0, & t \rightarrow s, u < 0. \end{aligned}$$

(iii)  $H_3(s, u) := H_3(s, u; x, y)$  satisfies  $\mathcal{L}_{s,u}H_3(s, u) = \alpha H_3(s, u)$  in  $(0, \infty) \times (0, \infty)$ , together with the boundary conditions, for  $s, x, y > 0$ ,

$$\begin{aligned} H_3(s, u; x, y) &= \mathbf{1}(-y \leq u), & u \leq 0, \\ H_3(s, u; x, y) &\rightarrow 0, & u \rightarrow \infty. \end{aligned}$$

*Proof.* (i) By the strong Markov property of  $U_t$ , it can be proved that  $H_1(t \wedge T(s), U_{t \wedge T(s)})$  is an  $\mathcal{F}_t$ -martingale on  $t \in [s, \infty)$ . Applying Itô's formula gives

$$H_1(t \wedge T(s), U_{t \wedge T(s)}) - H_1(s, u) = \int_0^{t \wedge T(s)} \mathcal{L}_{r,u}H_1(r, U_r)dr + \text{martingale}.$$

It follows that  $\int_0^{t \wedge T(s)} \mathcal{L}_{r,u}H_1(r, U_r)dr$  is a zero-mean  $\mathcal{F}_t$ -martingale. Hence,  $\int_0^{t \wedge T(s)} \mathcal{L}_{r,u}H_1(r, U_r)dr = 0$ , and thus  $\mathcal{L}_{s,u}H_1(s, u) = 0$ , since  $t$  is arbitrary and  $\mathcal{L}_{s,u}H_1(s, u)$  is continuous. The boundary conditions are obvious.

(ii) and (iii) can be proved by using the same argument as in (i).

*Remark 3.1.* Notice that  $\psi_1(s, u) = H_1(s, u; \infty, \infty)$ ,  $\psi_2(s, u, t) = H_2(s, u; \infty, \infty, t)$  and  $\psi_3(s, u) = H_3(s, u; \infty, \infty)$ . Thus, from Theorem 3.1 we can obtain the integro-differential equations for  $\psi_i$ 's. In particular, for the homogeneous Markov process  $\{U_t\}_{t \geq 0}$  in (2.5) with  $f(t, y, z) = z$ ,  $q(s, \cdot) = q(\cdot)$ ,  $\sigma(s, \cdot) = \sigma(\cdot)$  and  $\lambda(t, dz) = \lambda dG(z)$ , where  $G$  is the common distribution function of the claim sizes and  $\lambda$  is the constant intensity of the homogeneous Poisson process  $N$ , we obtain the integro-differential equations for  $\psi_1(0, u)$ ,  $\psi_2(0, u, t)$  and  $\psi_3(0, u)$  that have been considered in (2).

#### 4. RISK MODEL WITH INTEREST FORCE

We have shown in the last section that the probability of ultimate ruin and the distributions of the time of ruin, the surplus immediately prior to ruin and the deficit at ruin satisfy certain integro-differential equations. However, the equations involved are so complicated that explicit solutions are generally hard to get even for particular cases. Examples of the solutions for some non-trivial special cases can be found in (2). In this section we give the Laplace–Stieltjes transform of  $H_1$  as well as its estimation for the compound Poisson risk model with a constant interest force.

Consider the risk process  $U_t$  in (2.5) with  $q(s, x) = c + \rho x$ ,  $\sigma(s, x) = \sigma$ ,  $\mu_R \equiv 0$  and  $K_R \equiv 0$ , i.e.  $dU_t = cdt + \rho U_t dt + \sigma dW_t - dX_t$ , its solution is given by

$$U_t = ue^{\rho t} + c \int_0^t e^{\rho y} dy + \sigma \int_0^t e^{\rho(t-s)} dW_s - \int_0^t e^{\rho(t-v)} dX_v,$$

where  $X_t = \int_0^t \int_0^\infty z N(ds, dz)$ ,  $u > 0$  is the initial capital,  $c > 0$  the premium rate and  $\rho > 0$  the interest force. In order to indicate the role of  $\rho$  and  $\sigma$ , we let  $\psi_{\rho, \sigma}(u) := \psi_1(u) = P(\inf_{t \geq 0} U_t < 0 | U_0 = u)$  denote the ultimate ruin probability for this risk



process. For the claim distribution  $G$  with mean  $\mu$ , denote by  $L(x) = \mu^{-1} \int_0^x (1-G(y))dy$  the integrated tail of the claim size distribution. Moreover, let

$$Q(\alpha, x, y) := \int_0^x e^{-\alpha u} dL(u) - e^{\alpha y} \int_y^{y+x} e^{-\alpha u} dL(u), \quad \alpha, x, y > 0.$$

The Laplace-Stieltjes transform of a function is denoted by putting a hat on the top of the function, for example,  $\hat{G}(\beta) = \int_0^\infty e^{-\beta u} dG(u)$ . Because the process  $U_t$  is homogeneous, we consider  $H_1(0, u; x, y)$  instead of  $H_1(s, u; x, y)$  and denote it by  $H_1(u; x, y)$ .

**Theorem 4.1.** *Assume that  $H_1(u) := H_1(u; x, y)$  has bounded continuous first and second derivatives. Then*

(i) *If  $\sigma = 0$ , then*

$$\hat{H}_1(\beta) = \int_\beta^\infty \frac{\lambda\mu}{\rho} \left( H_1(0)\hat{L}(\alpha) - Q(\alpha, x, y) \right) \exp\left(-\frac{a_1(\alpha) - a_1(\beta)}{\rho}\right) d\alpha,$$

where  $a_1(\beta) = c\beta - \lambda\mu \int_0^\beta \hat{L}(\alpha) d\alpha$ , and

$$H_1(0) = \frac{\lambda\mu \int_0^\infty Q(\alpha, x, y) \exp\left(-\frac{a_1(\alpha)}{\rho}\right) d\alpha}{\rho + \lambda\mu \int_0^\infty \hat{L}(\alpha) \exp\left(-\frac{a_1(\alpha)}{\rho}\right) d\alpha}. \quad (4.1)$$

(ii) *If  $\sigma \neq 0$ , then*

$$\hat{H}_1(\beta) = -\frac{1}{\rho} \int_\beta^\infty \left( \frac{1}{2}\sigma^2 h_2'(0) - \lambda\mu \hat{L}(\alpha) + \lambda\mu Q(\alpha, x, y) \right) \exp\left(-\frac{a_2(\alpha) - a_2(\beta)}{\rho}\right) d\alpha,$$

where  $a_2(\beta) = (c + \frac{1}{2}\sigma^2)\beta - \lambda\mu \int_0^\beta \hat{L}(\alpha) d\alpha$ , and

$$\frac{1}{2}\sigma^2 h_2'(0) = \frac{\rho + \lambda\mu \int_0^\infty \exp\left(-\frac{a_2(\alpha)}{\rho}\right) \left( \hat{L}(\alpha) - Q(\alpha, x, y) \right) d\alpha}{\int_0^\infty \exp\left(-\frac{a_2(\alpha)}{\rho}\right) d\alpha}.$$

*Proof.* It follows from Theorem 3.1(i) that  $H_1(u)$  satisfies

$$\begin{aligned} \frac{1}{2}\sigma^2 H_1''(u) + (\rho u + c)H_1'(u) + \lambda \int_0^u H_1(u-z)dG(z) \\ + \lambda(G(u+y) - G(u))\mathbf{1}(u \leq x) = \lambda H_1(u), \quad u > 0. \end{aligned} \quad (4.2)$$

Integrating (4.2) gives

$$\begin{aligned} & \frac{1}{2}\sigma^2(H_1'(u)-H_1'(0)) + (\rho u + c)H_1(u) - c + \lambda \int_0^{u \wedge x} (G(y+v) - G(v))dv \\ & = \rho \int_0^u H_1(v)dv + \lambda \int_0^u H_1(u-z)(1-G(z))dz, \quad u > 0. \end{aligned} \quad (4.3)$$

(i) Consider the case  $\sigma = 0$ . To avoid complications with the jump of  $H_1(\cdot)$  at 0, just as Sundt and Teugels (10), we introduce the auxiliary distribution

$$h_1(u) := h_1(u; x, y) := 1 - \frac{H_1(u; x, y)}{H_1(0; x, y)},$$

from which we have  $H_1(u; x, y) = H_1(0; x, y)(1 - h_1(u; x, y))$  and  $h_1(0; x, y) = 0$ . In terms of  $h_1(u)$ , equation (4.3) can be written as

$$\begin{aligned} ch_1(u) &= -\rho \int_0^u v dh_1(v) + \lambda \mu (h_1 * L)(u) \\ &\quad - \lambda \mu L(u) + \frac{\lambda}{H_1(0)} \int_0^{u \wedge x} (G(y+v) - G(v))dv, \end{aligned} \quad (4.4)$$

where  $*$  denotes the Stieltjes convolution.

Taking the Laplace-Stieltjes transform of (4.4) gives

$$c\hat{h}_1(\beta) = \rho\hat{h}_1'(\beta) + \lambda\mu\hat{h}_1(\beta)\hat{L}(\beta) - \lambda\mu\hat{L}(\beta) + \frac{\lambda\mu}{H_1(0)}Q(\beta, x, y). \quad (4.5)$$

The solution of (4.5) with the boundary condition  $\lim_{\beta \rightarrow \infty} \hat{h}_1(\beta) = 0$  is given by

$$\hat{h}_1(\beta) = -\frac{\lambda\mu}{\rho} \int_{\beta}^{\infty} \left( \hat{L}(\alpha) - \frac{Q(\alpha, x, y)}{H_1(0)} \right) \exp\left(-\frac{a_1(\alpha) - a_1(\beta)}{\rho}\right) d\alpha, \quad (4.6)$$

where  $a_1(\beta) = c\beta - \lambda\mu \int_0^{\beta} \hat{L}(\alpha) d\alpha$ .

Letting  $\beta \rightarrow 0$  in (4.6) yields (4.1), and because  $\hat{H}_1(\beta) = -H_1(0)\hat{h}_1(\beta)$ , the result follows.

(ii) Consider  $\sigma \neq 0$ . In this case  $H_1(0) = 1$ , and so the auxiliary distribution is

$$h_2(u) := h_2(u; x, y) := 1 - H_1(u; x, y).$$

In terms of  $h_2(u)$ , equation (4.3) can be written as

$$\begin{aligned} \frac{1}{2}\sigma^2(h_2'(u) - h_2'(0)) + ch_2(u) &= -\rho \int_0^u v dh_2(v) + \lambda\mu(h_2 * L)(u) \\ &\quad - \lambda\mu L(u) + \lambda \int_0^{u \wedge x} (G(y+v) - G(v)) dv. \end{aligned} \quad (4.7)$$

Taking the Laplace-Stieltjes transform of (4.7) gives

$$\frac{1}{2}\sigma^2(-h_2'(0) + \beta \hat{h}_2(\beta)) + c\hat{h}_2(\beta) = \rho \hat{h}_2(\beta) + \lambda\mu \hat{h}_2(\beta) \hat{L}(\beta) - \lambda\mu \hat{L}(\beta) + \lambda\mu Q(\beta, x, y). \quad (4.8)$$

The solution of (4.8) with the boundary condition  $\lim_{\beta \rightarrow \infty} \hat{h}_2(\beta) = 0$  is given by

$$\hat{h}_2(\beta) = \frac{1}{\rho} \int_{\beta}^{\infty} \left( \frac{1}{2}\sigma^2 h_2'(0) - \lambda\mu \hat{L}(\alpha) + \lambda\mu Q(\alpha, x, y) \right) \exp\left(-\frac{a_2(\alpha) - a_2(\beta)}{\rho}\right) d\alpha, \quad (4.9)$$

where  $a_2(\beta) = (c + \frac{1}{2}\sigma^2)\beta - \lambda\mu \int_0^{\beta} \hat{L}(\alpha) d\alpha$ .

Letting  $\beta \rightarrow 0$  in (4.9) yields the result.

**Theorem 4.2.** *Suppose that the safety loading  $c - \lambda\mu$  is positive and  $\rho, x, y > 0$ . We have*

$$\frac{B_{\sigma}(u; x, y)}{u} - \int_u^{\infty} \frac{A_{\sigma}(z; x, y)}{z^2} dz \leq H_1(u; x, y) \leq \frac{A_{\sigma}(u, x, y)}{u} - \int_u^{\infty} \frac{B_{\sigma}(z; x, y)}{z^2} dz, \quad (4.10)$$

where

$$\begin{aligned}
A_\sigma(u, x, y) &= \frac{\lambda}{\rho(1 - \psi_{0,\sigma}(u))} \left( \int_u^\infty (G(v+y) - G(v))dv - \psi_{0,\sigma}(u) \right. \\
&\quad \left. \times \int_0^\infty (G(v+y) - G(v))dv + \psi_{0,\sigma}(u) * \int_0^u (G(v+y) - G(v))dv \right), \\
B_\sigma(u; x, y) &= \frac{\lambda}{\rho + u^{-1}c} \int_u^\infty (G(v+y) - G(v))dv, \\
1 - \psi_{0,0}(u) &= \left( 1 - \frac{\lambda\mu}{c} \right) \sum_{n=0}^\infty \left( \frac{\lambda\mu}{c} \right)^n L^{n,*}(u), \\
1 - \psi_{0,\sigma}(u) &= \left( 1 - \frac{\lambda\mu}{c} \right) \sum_{n=0}^\infty \left( \frac{\lambda\mu}{c} \right)^n (J^{(n+1),*} * L^{n,*})(u),
\end{aligned}$$

with  $J(u) = (1 - \exp(-\frac{2c}{\sigma^2}u)) \mathbf{1}(u > 0)$ ,  $L^{n,*}$ , for  $n \geq 1$ , denotes the  $n$ -fold convolution of  $L$  with itself and  $L^{0,*}$  is the distribution function corresponding to the Dirac measure at zero.

*Proof.* Consider the case  $\sigma = 0$ . By (4.5) we have

$$\hat{h}_1(\beta) = \sum_{n=0}^\infty \left( \frac{\lambda\mu}{c} \hat{L}(\beta) \right)^n \left( \frac{\rho}{c} \hat{h}'_1(\beta) - \frac{\lambda\mu}{c} \hat{L}(\beta) + \frac{\lambda\mu}{cH_1(0)} Q(\beta, x, y) \right).$$

Inverting it gives

$$\begin{aligned}
h_1(u) &= \frac{\rho}{c} \sum_{n=0}^\infty \left( \frac{\lambda\mu}{c} \right)^n (L^{n,*} * k_1)(u) - \sum_{n=0}^\infty \left( \frac{\lambda\mu}{c} \right)^{n+1} L^{(n+1),*}(u) \\
&\quad + \frac{\lambda}{cH_1(0)} \sum_{n=0}^\infty \left( \frac{\lambda\mu}{c} \right)^n L^{n,*}(u) * \int_0^u (G(v+y) - G(v))dv,
\end{aligned} \tag{4.11}$$

where  $k_1(u) = \int_u^\infty zdh_1(z)$ .

Letting  $u \rightarrow \infty$  yields

$$H_1(0) = \frac{\lambda \int_0^\infty (G(v+y) - G(v))dv}{c + \rho \int_0^\infty udh_1(u)}. \tag{4.12}$$

We can rewrite (4.11), in terms of  $\psi_{0,0}$ , as

$$\begin{aligned} h_1(u) &= \frac{\rho}{c - \lambda\mu} ((1 - \psi_{0,0}) * k_1)(u) - \frac{\lambda\mu}{c - \lambda\mu} ((1 - \psi_{0,0}) * L)(u) \\ &\quad + \frac{\lambda}{H_1(0)(c - \lambda\mu)} ((1 - \psi_{0,0}(u)) * \int_0^u (G(v + y) - G(v))dv). \end{aligned}$$

Because  $1 \geq h_1(u)$  and  $((1 - \psi_{0,0}) * k_1)(u) \geq (1 - \psi_{0,0}(u))(k_1(u) - k_1(0))$ , it follows that

$$\begin{aligned} \rho(1 - \psi_{0,0}(u))k_1(u) &\leq \frac{\lambda}{H_1(0)} \int_u^\infty (G(v + y) - G(v))dv \\ &\quad - \psi_{0,0}(u) \frac{\lambda \int_0^\infty (G(v + y) - G(v))dv}{H_1(0)} \\ &\quad + \frac{\lambda}{H_1(0)} \psi_{0,0}(u) * \int_0^u (G(v + y) - G(v))dv, \end{aligned} \quad (4.13)$$

where we have used (4.12). By (4.4),

$$\begin{aligned} \frac{1}{c}k_1(u) &= \frac{\rho}{c}k_1(0) + h_1(u) + \frac{\lambda\mu}{c}L(u) - \frac{\lambda\mu}{c}(h_1 * L)(u) \\ &\quad - \frac{\lambda}{cH_1(0)} \int_0^u (G(y + v) - G(v))dv. \end{aligned} \quad (4.14)$$

Letting  $u \rightarrow \infty$  gives

$$\frac{\rho}{c}k_1(0) + 1 - \frac{\lambda}{cH_1(0)} \int_0^\infty (G(v + y) - G(v))dv = 0. \quad (4.15)$$

Because  $h_1(u) \geq 1 - u^{-1}k_1(u)$  and  $L(u) - (h_1 * L)(u) \geq 0$ , it follows from (4.14) and (4.15) that

$$k_1(u) \geq \frac{\lambda}{H_1(0)(\rho + cu^{-1})} \int_u^\infty (G(v + y) - G(v))dv. \quad (4.16)$$

Note that

$$H_1(u) = H_1(0) \left( \frac{k_1(u)}{u} - \int_u^\infty \frac{k_1(v)}{v^2} dv \right),$$

the result given in (4.10) follows easily from (4.13) and (4.16).

Next, consider the case  $\sigma \neq 0$ . Since the proof is similar to the proof of the first case, we only indicate the key steps. Inverting the solution of  $\hat{h}_2(\beta)$  obtained from (4.8) yields

$$h_2(u) = \frac{1}{c - \lambda\mu} \left( \frac{\sigma^2}{2} h_2'(0)(1 - \psi_{0,\sigma}(u)) + \rho((1 - \psi_{0,\sigma}) * k_2)(u) - \lambda\mu((1 - \psi_{0,\sigma}) * L)(u) + \lambda(1 - \psi_{0,\sigma}(u)) * \int_0^u (G(v+y) - G(v))dv \right),$$

where  $k_2(u) = \int_u^\infty v dh_2(v)$ .

Letting  $\beta \rightarrow 0$  in (4.5) gives

$$\frac{\sigma^2}{2c} h_2'(0) = 1 + \frac{\rho}{c} k_2(0) - \frac{\lambda}{c} \int_0^\infty (G(v+y) - G(v))dv. \quad (4.17)$$

Similar to (4.13) we have

$$\begin{aligned} \rho(1 - \psi_{0,\sigma}(u))k_2(u) &\leq \lambda \int_u^\infty (G(v+y) - G(v))dv - \lambda\psi_{0,\sigma}(u) \int_0^\infty (G(v+y) - G(v))dv \\ &\quad + \lambda\psi_{0,\sigma}(u) * \int_0^u (G(v+y) - G(v))dv. \end{aligned} \quad (4.18)$$

Using (4.7) and (4.17), we get

$$k_2(u) \geq \frac{\lambda}{\rho + cu^{-1}} \int_u^\infty (G(v+y) - G(v))dv. \quad (4.19)$$

The result follows from (4.18) and (4.19) for the case  $\sigma \neq 0$ .

### ACKNOWLEDGMENTS

This work was supported by a grant from Research Grants Council of Hong Kong Special Administrative Region (Project No. HKBU/2075/98p) and by the National Natural Science Foundation of China (Project No. 19801020).

## REFERENCES

1. Paulsen, J. Risk theory in a stochastic economic environment. *Stochastic Process. Appl.* **1993**, 46, 327-361.
2. Paulsen, J.; Gjessing, H. K. Ruin theory with stochastic return on investments. *Adv. Appl. Prob.* **1997**, 29, 965-985.
3. Paulsen, J. Sharp conditions for certain ruin in a risk process with stochastic return on investments. *Stochastic Process. Appl.* **1998**, 75, 135-148.
4. Paulsen, J. Ruin theory with compounding assets—a survey. *Insurance: Math. Econ.* **1998**, 22, 3-16.
5. Rolski, T.; Schmidli, H.; Schmidt, V.; Teugels, J. *Stochastic Processes for Insurance and Finance*. Wiley: New York, 1999.
6. Møller, C. M. Stochastic differential equations for ruin probabilities. *J. Appl. Prob.* **1995**, 32, 74-89.
7. Møller, C. M. Integral equations for compound distribution functions. *J. Appl. Prob.* **1996**, 33, 399-399.
8. Ikeda, N.; Watanabe, S. *Stochastic Differential Equations and Diffusion Processes*. North-Holland Publishing Company: Amsterdam, 1981.
9. Protter, P. *Stochastic Integration and Differential Equations* (2nd Edition). Springer-Verlag: Berlin, 1992.
10. Sundt, B.; Teugels, J. L. Ruin estimates under interest force. *Insurance: Math. Econ.* **1995**, 16, 7-22.