

## K-resonance in toroidal polyhexes

Shiu, Wai Chee; Lam, Peter Che Bor; Zhang, Heping

*Published in:*  
Journal of Mathematical Chemistry

*DOI:*  
[10.1007/s10910-004-6897-4](https://doi.org/10.1007/s10910-004-6897-4)

Published: 01/11/2005

*Document Version:*  
Peer reviewed version

[Link to publication](#)

*Citation for published version (APA):*  
Shiu, W. C., Lam, P. C. B., & Zhang, H. (2005). K-resonance in toroidal polyhexes. *Journal of Mathematical Chemistry*, 38(4), 451-466. <https://doi.org/10.1007/s10910-004-6897-4>

### General rights

Copyright and intellectual property rights for the publications made accessible in HKBU Scholars are retained by the authors and/or other copyright owners. In addition to the restrictions prescribed by the Copyright Ordinance of Hong Kong, all users and readers must also observe the following terms of use:

- Users may download and print one copy of any publication from HKBU Scholars for the purpose of private study or research
- Users cannot further distribute the material or use it for any profit-making activity or commercial gain
- To share publications in HKBU Scholars with others, users are welcome to freely distribute the permanent publication URLs

# *k*-resonance in toroidal polyhexes

Wai Chee Shiu<sup>a</sup>, Peter Che Bor Lam<sup>a</sup>, Heping Zhang<sup>b</sup>

<sup>a</sup>Department of Mathematics, Hong Kong Baptist University, Kowloon Tong, Hong Kong, P. R. China,  
e-mails: wshiu@hkbu.edu.hk, cblam@hkbu.edu.hk

<sup>b</sup>School of Mathematics and Statistics, Lanzhou University, Lanzhou, Gansu 730000, P. R. China,  
e-mail: zhanghp@lzu.edu.cn

Running Title : *k*-resonance in toroidal polyhexes

All correspondence should be sent to:

Wai Chee Shiu  
Department of Mathematics,  
Hong Kong Baptist University,  
224 Waterloo Road, Kowloon Tong,  
Hong Kong, China.

# *k*-resonance in toroidal polyhexes\*

Wai Chee Shiu<sup>a</sup>, Peter Che Bor Lam<sup>a</sup>, Heping Zhang<sup>b</sup>

<sup>a</sup>Department of Mathematics, Hong Kong Baptist University, Kowloon Tong, Hong Kong, P. R. China,  
e-mails: wshiu@hkbu.edu.hk, cblam@hkbu.edu.hk

<sup>b</sup>School of Mathematics and Statistics, Lanzhou University, Lanzhou, Gansu 730000, P. R. China,  
e-mail: zhanghp@lzu.edu.cn

## Abstract

This paper considers the *k*-resonance of a toroidal polyhex (or toroidal graphitoid) with a string  $(p, q, t)$  of three integers ( $p \geq 2, q \geq 2, 0 \leq t \leq p - 1$ ). A toroidal polyhex  $G$  is said to be *k*-resonant if, for  $1 \leq i \leq k$ , any  $i$  disjoint hexagons are mutually resonant, that is,  $G$  has a Kekulé structure (perfect matching)  $M$  such that these hexagons are  $M$ -alternating (in and off  $M$ ). Characterizations for 1, 2 and 3-resonant toroidal polyhexes are given respectively in this paper.

**Keywords** : Fullerene, toroidal polyhex, Kekulé structure, *k*-resonance.

**AMS 2000 MSC** : 05C10, 05C70, 05C90

## 1 Introduction

The discovery of the fullerene molecules and nanotubes has stimulated much interests in other possibilities for carbons. Classical fullerene is an all-carbon molecule in which the atoms are arranged on a pseudospherical framework made up entirely of pentagons and hexagons. Its molecular graph is a finite trivalent graph embedded on the surface of a sphere with only hexagonal and (exactly 12) pentagonal faces. Deza et al. [4] considered fullerene's extension to other closed surfaces and showed that only four surfaces are possible: sphere, torus, Klein bottle and projective plane. Unlike spherical fullerenes, toroidal and Klein bottle's fullerenes have been regarded as tessellations of entire hexagons on their surfaces since they must contain no pentagons [4, 11]. For the theoretical consideration and detailed classifications of hexagonal tilings (dually, triangulations) on the torus and the Klein bottle, see [17, 22]. Toroidal fullerenes are likely to have direct experimental relevance since "crop circles fullerenes" discovered by Liu et al. [14] in 1997 are presumably torus-shaped.

A *toroidal polyhex* (or *toroidal graphitoid*, *torene*) is a toroidal fullerene that can be described by a string  $(p, q, t)$  of three integers ( $p \geq 1, q \geq 1, 0 \leq t \leq p - 1$ ); its definition is referred to the next section. Some features of toroidal polyhexes with chemical relevance were discussed [9, 10]. For example, a systematic coding and classification scheme were given for the enumeration of isomers of toroidal polyhexes, the calculation of the spectrum and the count for spanning trees. There have been a few work in the enumeration of perfect matchings of toroidal polyhexes by applying various techniques, such as transfer-matrix [12, 21], permanent of the adjacency matrix [1], and Pfaffian orientation [8].

This paper considers the *k*-resonance of toroidal polyhexes. The concept of *k*-resonance originates from Clar's aromatic sextet theory [3] and Randić's conjugated circuit model [18, 19, 20]. In the former Clar found that various electronic properties of polycyclic aromatic hydrocarbons can be predicted by defining mutually resonant sextets [7] (i.e., disjoint hexagons that are all alternating with respect to a Kekulé structure). In Randić's model a conjugated hexagon has the largest contribution of the resonance energy among all  $4n + 2$  conjugated circuits (a cycle is said

---

\*This work is supported by FRG, Hong Kong Baptist University; NSFC and TRAPOYT.

to be conjugated or resonant if it is alternating with respect to a Kekulé structure). A benzenoid system is said to be  $k$ -resonant if, for  $1 \leq i \leq k$ , any  $i$  disjoint hexagons are mutually resonant. For a recent survey on  $k$ -resonant benzenoids and  $k$ -cycle resonant graphs, see [5]. Zhang and Chen [24] characterized completely 1-resonant benzenoid systems, solved such a problem proposed by Gutman [6] and showed its equivalence to normal benzenoid systems (i.e., each edge is contained in a Kekulé structure). The similar results were extended to coronoid systems (benzenoids with holes) [2] and plane bipartite graphs [27]. Later, Zheng [28, 29] characterized general  $k$ -resonant benzenoid systems. In particular he showed that any 3-resonant benzenoid systems are also  $k$ -resonant for any integer  $k \geq 3$  and gave their systematic construction. The same result and similar construction are still valid for coronoid systems [2, 13] and open-ended carbon nanotubes [23].

In this paper mutually resonant hexagons and  $k$ -resonance are naturally extended to toroidal polyhexes  $H(p, q, t)$  ( $p \geq 2, q \geq 2$ ) (in some degenerated cases, for instance,  $H(1, q, 0)$ ,  $H(p, 1, 0)$  and  $H(p, 1, p - 1)$ , a hexagonal face is not bounded by a cycle). We point out several hexagon-preserving automorphisms of toroidal polyhexes and thus show the vertex- and hexagon-transitivity of this kind of graphs. Then we give a sufficient condition for some disjoint hexagons being mutually resonant whereby we show that all toroidal polyhexes are 1-resonant only except for  $(2, 2, 0)$ , though they are all elementary bipartite graphs. This exception demonstrates a great difference with the plane situation. Further a simple characterization for 2-resonant toroidal polyhexes are given. Finally we completely characterize 3-resonant toroidal polyhexes.

## 2 Toroidal polyhex with symmetry

A *toroidal polyhex* is a 3-regular (cubic or trivalent) graph embedded on the torus such that each face is a hexagon, described by three parameters  $p, q$  and  $t$ , denoted by  $H(p, q, t)$  [16, 22], and drawn in the plane (equipped with the regular hexagonal lattice  $L$ ) using the representation of the torus by a  $p \times q$ -parallelogram  $P$  with the usual boundary identification (see Fig. 1): each side of  $P$  connects the centers of two hexagons, and is perpendicular to an edge-direction of  $L$ , both top and bottom sides pass through  $p$  vertical edges of  $L$  while two lateral sides pass through  $q$  edges. First identify its two lateral sides, then rotate the top cycle  $t$  hexagons, finally identify the top and bottom at their corresponding points. From this we get a toroidal polyhex  $H(p, q, t)$  with the torsion  $t$  ( $0 \leq t \leq p - 1$ ). In fact there is at least two ways to get such a toroidal polyhex. For example, see [8, 16].




Figure 1: A toroidal polyhex  $H(p, q, t)$  for  $p = 7, q = 5, t = 2$ .

We easily know that  $H(p, q, t)$  has  $pq$  hexagons,  $2pq$  vertices and  $3pq$  edges. The graph lying in the interior of the parallelogram  $P$  has a proper 2-coloring (white-black): the vertices incident with a downward vertical edge and with two upwardly oblique edges are colored black, and the

other vertices white (see Fig. 2). Such a 2-coloring is a proper 2-coloring of  $H(p, q, t)$ , i.e., the end-vertices of each edge receive distinct colors. Hence we have

**Proposition 2.1**  $H(p, q, t)$  is a bipartite graph.




Figure 2: Labeling for the hexagons and vertices of toroidal polyhex  $H(p, q, t)$  for  $p = 7, q = 5, t = 2$ .

For convenience, the hexagons and vertices of  $H(p, q, t)$  are labeled as an ordered pair of non-negative integers in the sense that we take the first component to be congruent modulo  $p$  and the second modulo  $q$ . We first establish an affine coordinate system  $XOY$  (see Fig. 2): take the bottom side as  $x$ -axis, a lateral side as  $y$ -axis, their intersection as the origin  $O$  such that both sides form an angle of  $60^\circ$ , and  $P$  lies on non-negative region. For any positive integer  $n$ , we shall use  $\mathbb{Z}_n$  to denote the set  $\{0, 1, \dots, n-1\}$  with arithmetic modulo  $n$ . The distance between a pair of parallel edges in a hexagon is a unit length. Each hexagon is labeled by the coordinates  $(x, y)$  of its center, where  $x \in \mathbb{Z}_p$  and  $y \in \mathbb{Z}_q$ . Hence such a hexagon is denoted by its label  $(x, y)$  (or  $h_{xy}$ ,  $h_{x,y}$ ). For each hexagon  $h_{xy}$ , choosing the upper one of a pair of parallel edges perpendicular to  $y$ -axis, the black end-vertex is named by  $b_{xy}$  (or  $b_{x,y}$ ) and the white end-vertex by  $w_{xy}$  (or  $w_{x,y}$ ) (for example, see Fig. 2). In this notations, each  $w_{0,y}$  is adjacent to  $b_{0,y}$  and each  $w_{x,0}$  to  $b_{x+t+1,q-1}$ ; the  $y$ -th layer is the even cycle  $w_{0,y}b_{1,y}w_{1,y}b_{2,y} \cdots w_{p-1,y}b_{0,y}w_{0,y}$ ,  $0 \leq y \leq q-1$ .

An automorphism  $\phi$  of a simple graph is a bijection from the vertex-set to itself so that both  $\phi$  and the inverse  $\phi^{-1}$  preserve the adjacency between vertices. We now define three types of automorphisms of toroidal polyhex  $H(p, q, t)$  as follows: the r-l shift  $\phi_{rl}$  moves horizontally backward every vertex a unit, i.e.,

$$\phi_{rl}(w_{x,y}) = w_{x-1,y} \text{ and } \phi_{rl}(b_{x,y}) = b_{x-1,y}, \text{ for every pair } (x, y). \quad (1)$$

The t-b shift  $\phi_{tb}$  moves downwards every vertex a unit from the  $y$ -axis, but the  $x$ -coordinates may change. More precisely,

$$\phi_{tb}(w_{x,y}) = w_{x,y-1}, \text{ and } \phi_{tb}(b_{x,y}) = b_{x,y-1} \text{ for } 1 \leq y \leq q-1, \quad (2)$$

and

$$\phi_{tb}(w_{x,0}) = w_{x+t,q-1} \text{ and } \phi_{tb}(b_{x,0}) = b_{x+t,q-1}. \quad (3)$$

It can be seen that both shifts  $\phi_{lr}$  and  $\phi_{tb}$  are hexagon-preserving automorphism of  $H(p, q, t)$ . Further  $\phi_{rl}$  and  $\phi_{tb}$  generate a subgroup of the automorphism group of  $H(p, q, t)$ , denoted by  $\langle \phi_{rl}, \phi_{tb} \rangle$ , which acts transitively on the set of hexagons of  $H(p, q, t)$ ; that is, for each pair of hexagons  $h$  and  $h'$  there is a hexagon-preserving automorphism  $g \in \langle \phi_{rl}, \phi_{tb} \rangle$  so that  $g(h) = h'$ . Hence we have the following result.

**Lemma 2.2**  $H(p, q, t)$  is hexagon-transitive.

Finally, let  $R_2$  be the rotation of  $180^\circ$  about the center of the parallelogram  $P$ . Then  $R_2$  is also a hexagon-preserving automorphism of  $H(p, q, t)$  that interchanges the black and white vertices. The generated subgroup  $\langle \phi_{rl}, \phi_{tb}, R_2 \rangle$  is transitive on the vertex-set of  $H(p, q, t)$ .

**Lemma 2.3**[22, 16]  $H(p, q, t)$  is vertex-transitive.

### 3 1- and 2-Resonance

A *perfect matching* or *1-factor*  $M$  (Kekulé structure in chemistry) of a graph  $G$  is a set of pairwise disjoint edges of  $G$  such that every vertex of  $G$  is incident with an edge in  $M$ . A bipartite graph is called *elementary* [15] if it is connected and each edge is contained in a perfect matching. We easily know that any toroidal polyhex is an elementary bipartite graph since it is 3-regular. In fact the edge-set of a toroidal polyhex can be decomposed into three perfect matchings so that each consists of all edges with the same edge-direction [8].

The concept for  $k$ -resonance is now extended to toroidal polyhexes. The set of some disjoint hexagons of  $H(p, q, t)$  is called a *resonant pattern*, or these hexagons are *mutually resonant*, if  $H(p, q, t)$  has a perfect matching  $M$  such that these hexagons are all  $M$ -alternating (in and off  $M$ ) cycles. In some degenerated cases of toroidal polyhexes, for instance,  $H(1, q, 0)$ ,  $H(p, 1, 0)$  and  $H(p, 1, p-1)$ , a hexagonal face is not bounded by a cycle. So we only consider toroidal polyhexes  $H(p, q, t)$  with  $p \geq 2$  and  $q \geq 2$ . In such cases disjoint hexagons of  $H(p, q, t)$  are mutually resonant if and only if the subgraph obtained from  $H(p, q, t)$  by deleting the vertices of the hexagons either has a perfect matching or is empty (since each hexagonal face is bounded by a cycle with length 6).

**Definition 3.1** For some positive integer  $k$ , a toroidal polyhex  $H(p, q, t)$  is called  *$k$ -resonant* if for any  $i$  ( $\leq k$ ) disjoint hexagons of  $H(p, q, t)$  are mutually resonant.

**Remark 3.1** By the definition above, a toroidal polyhex  $H(p, q, t)$  is  $k$ -resonant if it is  $(k-1)$ -resonant and does not contain any  $k$  disjoint hexagons.

To obtain our characterizations (the main results in this paper) for  $k$ -resonance of toroidal polyhexes, the following provides us a crucial approach. Let  $\mathcal{S}$  be a subgraph of a toroidal polyhex  $H(p, q, t)$  for which each component is either a hexagon or an edge with the end-vertices. Then  $\mathcal{S}$  is called a *Clar cover* [26] if  $\mathcal{S}$  includes all vertices of  $H(p, q, t)$ ; an *ideal configuration* if  $\mathcal{S}$  is alternately incident with white and black vertices along any direction of each layer.

**Lemma 3.1** Any ideal configuration  $\mathcal{S}$  of a toroidal polyhex  $H(p, q, t)$  can be extended to a Clar cover, and the hexagons in  $\mathcal{S}$  are thus mutually resonant.

**Proof.** Let  $L_y$  be the  $y$ -th layer of  $H(p, q, t)$ ,  $0 \leq y \leq q-1$ . Let  $L_y - \mathcal{S}$  be the subgraph obtained from  $L_y$  by deleting all vertices of  $\mathcal{S}$  together with their incident edges. If  $\mathcal{S}$  contains a vertex of

$L_y$ , each component of  $L_y - \mathcal{S}$  is a path with odd length for which the end-vertices have different colors because  $\mathcal{S}$  is alternately incident with white and black vertices along the cycle  $L_y$ . Hence each  $L_y - \mathcal{S}$  has a perfect matching. This implies that  $H(p, q, t) - \mathcal{S}$  has a perfect matching. Hence  $\mathcal{S}$  can be extended to a Clar cover, and its hexagons are thus mutually resonant.  $\square$

In this section we first give 1-resonant toroidal polyhexes, and then characterize 2-resonant toroidal polyhexes.

**Theorem 3.2** *A toroidal polyhex  $H(p, q, t)$  ( $p, q \geq 2$ ) is 1-resonant if and only if  $(p, q, t) \neq (2, 2, 0)$ .*

**Proof:** Since  $H(p, q, t)$  is hexagon-transitive (Lemma 2.2), it suffices to check whether any given hexagon is resonant or not.

**Case 0:**  $p = 2 = q$ . Choose the hexagon  $h_{11}$  (see Fig. 3).  $H(2, 2, 0) - h_{11}$  consists of exactly two isolated vertices  $w_{00}$  and  $b_{01}$ . This implies that  $H(2, 2, 0)$  is not 1-resonant.  $H(2, 2, 1) - h_{11}$  consists of exactly two adjacent vertices  $w_{00}$  and  $b_{01}$ . Hence  $H(2, 2, 1)$  is 1-resonant.



Figure 3: (a)  $H(2, 2, 0)$

(b)  $H(2, 2, 1)$ .

**Case 1:**  $p \geq 2$  and  $q \geq 3$ . Choose the hexagon  $h_{11}$  (i.e., the cycle  $b_{11}w_{11}b_{20}w_{10}b_{10}w_{01}b_{11}$ , which is hatched in Fig. 4 (left)), and the vertical edges  $w_{00}b_{t+1,q-1}$ ,  $w_{12}b_{21}$ ,  $w_{13}b_{22}$ ,  $\dots$ ,  $w_{1,q-1}b_{2,q-2}$  (indicated by thick lines in Fig. 4 (left)). The chosen hexagon and vertical edges form an ideal configuration since it is incident with vertices:  $w_{00}, b_{10}, w_{10}$  and  $b_{20}$  in the 0-th layer;  $w_{01}, b_{11}, w_{11}$  and  $b_{21}$  in the first layer;  $w_{1y}, b_{2,y}$  in the  $y$ -th layers,  $2 \leq y \leq q - 2$ ; and  $w_{1,q-1}, b_{t+1,q-1}$  in the  $(q - 1)$ -th layer. Hence by Lemma 3.1  $h_{11}$  is a resonant hexagon.




Figure 4: Illustration for the proof of Theorem 3.2.

**Case 2:**  $p \geq 3$  and  $q \geq 2$ . For  $t \neq 0$ , the same arguments as Case 1 can be made. For  $t = 0$ , we choose the hexagon  $h_{2,1}$  (i.e., the cycle  $b_{21}w_{21}b_{30}w_{20}b_{20}w_{11}b_{21}$ ), and vertical edges  $w_{00}b_{1,q-1}, w_{02}b_{11}, \dots, w_{0,q-1}b_{1,q-2}$  (indicated by thick lines, see Fig. 4 (right)). By the same reason as Case 1 the chosen hexagon and vertical edges form an ideal configuration and the hexagon  $h_{2,1}$  is resonant.  $\square$

**Lemma 3.3**  $H(p, q, t)$  is 2-resonant for  $p \geq 3$  and  $q \geq 3$ .

**Proof:** It suffices to prove that any pair of disjoint hexagons  $(x_1, y_1)$  and  $(x_2, y_2)$  are mutually resonant, where  $x_i \in \mathbb{Z}_p$  and  $y_i \in \mathbb{Z}_q$ . By hexagon-preserving automorphisms of  $H(p, q, t)$  we need only to consider the following situation:

- (i)  $1 = y_1 \leq y_2 \leq \frac{q}{2} + 1 (\leq q - 1)$ , and
- (ii)  $1 = \min(x_1, x_2) \leq \max(x_1, x_2) \leq \frac{p}{2} + 1 (\leq p - 1)$ .

We first show this point. Without loss of generality we may assume that  $0 \leq y_1 \leq y_2 \leq q - 1$ . If  $0 = y_1 \leq y_2 \leq \frac{q}{2}$ , then we make the reversion of the t-b shift operation  $\phi_{tb}$  (see Eq. (2)) to  $H(p, q, t)$ , i.e.,  $\phi_{tb}^{-1}(x_1, 0) = (x_1, 1)$  and  $\phi_{tb}^{-1}(x_2, y_2) = (x_2, y_2 + 1)$ . The resulting hexagons satisfy statement (i). If  $y_2 - y_1 > \frac{q}{2}$ , then we make the t-b shift operation  $y_2 - 1$  times (see Eqs. (2) and (3)), i.e.,  $(x'_2, y'_2) = \phi_{tb}^{y_2-1}(x_2, y_2) = (x_2, y_2 - (y_2 - 1)) = (x_2, 1)$ , and  $(x'_1, y'_1) = \phi_{tb}^{y_2-1}(x_1, y_1) \equiv (x_1, y_1 - (y_2 - 1)) \pmod{q}$ . So  $y'_1 = y_1 - (y_2 - 1) + q \in \mathbb{Z}_q$ . Further  $1 = y'_2 \leq y'_1 = q - (y_2 - y_1) + 1 < \frac{q}{2} + 1$ . Otherwise,  $y_1 \geq 1$  and  $y_2 - y_1 \leq \frac{q}{2}$ . Similarly we can reduce this case to (i) by making the t-b shift operation  $y_1 - 1$  times. Then, by applying l-r shift operation  $\phi_{lr}$  that makes the  $y$ -coordinates to remain unchanged, similarly we can reduce all cases to the situation (ii).

To prove that  $(x_1, y_1)$  and  $(x_2, y_2)$  are mutually resonant, we will apply mainly the technique in Lemma 3.1: construct an ideal configuration containing such two hexagons by choosing a series of additional edges. We now distinguish the following four cases.

**Case 1:**  $x_1 = y_1 = x_2 = 1$  and  $3 \leq y_2 \leq \frac{q}{2} + 1$ . If  $y_2 \leq q - 2$ , then we choose vertical edges  $w_{00}b_{t+1, q-1}; w_{2,i}b_{3, i-1}, i = 2, \dots, y_2 - 1; w_{1,j}b_{2, j-1}, j = y_2 + 1, \dots, q - 1$ , (indicated by thick lines, see Fig. 5 (left)). If  $y_2 = q - 1$ , then  $q = 4$  only. For  $t \neq 0$ , we choose two edges  $w_{00}b_{t+1, 3}$  and  $w_{2,2}b_{3,1}$ . For  $t = 0$ ,  $w_{00}b_{1,3}$  is replaced by  $w_{20}b_{3,3}$  and the edge  $w_{22}b_{31}$  remain unchanged (see Fig. 5 (right)). It is easy to see that the chosen hexagons and vertical edges form an ideal configuration.




Figure 5: Illustration for Case 1 in the proof of Lemma 3.3.

**Case 2:**  $1 = x_1 < x_2 \leq \frac{p}{2} + 1$  and  $1 = y_1 < y_2 \leq \frac{q}{2} + 1$ . If  $y_2 \leq q - 2$ , then we choose vertical edges  $w_{00}b_{t+1, q-1}; w_{x_2-1, i}b_{x_2, i-1}, i = 2, \dots, y_2 - 1; w_{x_2, j}b_{x_2+1, j-1}, j = y_2 + 1, \dots, q - 1$ , which are indicated by thick lines in Fig. 6 (left). If  $y_2 = q - 1$ , then  $q = 3$  or  $4$ . For  $t \neq x_2 - 1$ , we choose edges  $w_{00}b_{t+1, q-1}$  and  $w_{x_2-1, 2}b_{x_2, 1}$  if  $q = 4$ , and we only choose the former if  $q = 3$  since the edge  $w_{00}b_{t+1, q-1}$  is disjoint with the hexagon  $(x_2, y_2)$ . For  $t = x_2 - 1$ ,  $w_{00}b_{t+1, q-1}$  is replaced by  $w_{20}b_{x_2+2, q-1}$  since  $x_2 + 2 \not\equiv x_2 \pmod{p}$ , and the other choices remain unchanged (see Fig. 6 (right)). For each subcase, the subgraph formed by the chosen hexagons and vertical edges is an ideal configuration.






Figure 6: Illustration for Case 2 in the proof of Lemma 3.3.

**Case 3:**  $1 = x_2 < x_1 \leq \frac{p}{2} + 1$  and  $1 = y_1 < y_2 \leq \frac{q}{2} + 1$ . If  $y_2 \leq q - 2$ , then we choose vertical edges  $w_{00}b_{t+1,q-1}$ ;  $w_{x_1,i}b_{x_1+1,i-1}$ ,  $i = 2, \dots, y_2 - 1$ ;  $w_{1,j}b_{2,j-1}$ ,  $j = y_2 + 1, \dots, q - 1$ , (indicated by thick lines, see Fig. 7 (left)). If  $y_2 = q - 1$ , then  $q = 3$  or  $4$ . For  $t \neq 0$ , we still choose such edges  $w_{00}b_{t+1,q-1}$  and  $w_{x_1,i}b_{x_1+1,i-1}$ ,  $i = 2, 3, \dots, y_2 - 1$ , since the edge  $w_{00}b_{t+1,q-1}$  is disjoint with the hexagon  $(1, y_2)$ . For  $t = 0$ ,  $w_{00}b_{1,q-1}$  is replaced by  $w_{10}b_{2,q-1}$  and the other choices remain unchanged (see Fig. 7 (right)). For each subcase, the chosen hexagons together with such vertical edges form an ideal configuration.




Figure 7: Illustration for Case 3 in the proof of Lemma 3.3.

**Case 4:**  $x_1 = y_1 = y_2 = 1$  and  $3 \leq x_2 \leq \frac{p}{2} + 1$ . This situation is more complicated than the previous three cases. Obviously,  $p \geq 4$  and  $1 \leq q - 2$ . We first choose the vertical edges  $w_{1,i}b_{2,i-1}$ ,  $w_{x_2,i}b_{x_2+1,i-1}$ ,  $i = 2, \dots, q - 1$  (indicated by thick lines, see Fig. 8).

For  $t = p - 1, 0, 1, \dots, x_2 - 3$ , we further choose two edges  $w_{20}b_{t+3,q-1}$  and  $w_{x_2+1,0}b_{x_2+2+t,q-1}$ . Since  $2 \leq t + 3 \leq x_2$  and  $x_2 + 1 \leq x_2 + 2 + t \leq 2x_2 - 1 \leq p + 1$  (note that  $t + 3, x_2 + 2 + t \in \mathbb{Z}_p$ ), the chosen part has the incident vertices  $w_{1,q-1}$ ,  $b_{t+3,q-1}$ ,  $w_{x_2,q-1}$  and  $b_{x_2+2+t,q-1}$  in the  $(q - 1)$ -th layer that alternate between white and black in one direction of this layer. The same fact holds obviously for the other layers.

For  $t = x_2 - 2, x_2 - 1, \dots, p - 2$ , we further choose two edges  $w_{20}b_{t+3,q-1}$  and  $w_{x,0}b_{x+1+t,q-1}$ , where  $x$  will be determined below. If  $t = p - 2$ , then put  $x := x_2 + 1$ . Since  $x_2 < 3 + t = p + 1$  and  $x + 1 + t \equiv x_2 \pmod{p}$ , the chosen part has the four incident vertices  $w_{1,q-1}$ ,  $b_{x_2,q-1}$ ,  $w_{x_2,q-1}$ ,  $b_{3+t,q-1}$ , ordered in the  $(q - 1)$ -th layer's direction. If  $x_2 - 2 \leq t \leq p - 3$ , then put  $x := x_2 - 2 - t \in \mathbb{Z}_p$ . Since  $x_2 + 1 \leq x \leq p$  and  $x + 1 + t = x_2 - 1$ , the chosen part has the incident vertices in the 0-th layer as ordered  $b_{10}$ ,  $w_{10}$ ,  $b_{20}$ ,  $w_{20}$ ,  $b_{x_2,0}$ ,  $w_{x_2,0}$ ,  $b_{x_2+1,0}$ ,  $w_{x,0}$ ; in the  $(q - 1)$ -th layer as  $w_{1,q-1}$ ,  $b_{x+1+t,q-1}$ ,  $w_{x_2,q-1}$ ,  $b_{3+t,q-1}$ , which alternate between white and black.




Figure 8: Illustration for Case 4 in the proof of Lemma 3.3.

For each subcase mentioned above the chosen hexagons  $(x_1, y_1)$  and  $(x_2, y_2)$  together with a series of corresponding vertical edges compose of an ideal configuration.

Therefore, by Lemma 3.1 any pair of disjoint hexagons are mutually resonant. Namely,  $H(p, q, t)$  is 2-resonant for  $p \geq 3$  and  $q \geq 3$ .  $\square$

**Lemma 3.4** For  $p \geq 4$ ,  $H(p, 2, t)$  is 2-resonant if and only if  $t$  is neither 0 nor  $p - 2$ .

**Proof:** Suppose  $t = 0$  or  $p - 2$ . We are going to show that  $H(p, 2, t)$  is not 2-resonant. Choose two disjoint hexagons  $h_{11}$  and  $h_{31}$ . Then vertices  $b_{20}$ ,  $b_{30}$  and  $b_{3+t,1}$  are neighbors of  $w_{20}$ . They lie in the hexagon  $h_{11}$  or  $h_{31}$ . Hence  $w_{20}$  is an isolated vertex of  $H(p, 2, t) - h_{11} - h_{31}$ . This shows that such two hexagons are not mutually resonant.

For the other cases, i.e.,  $t \notin \{0, p - 2\}$ , we will show that  $H(p, 2, t)$  is 2-resonant. By a similar argument as in the proof of Lemma 3.3 we consider a pair of disjoint hexagons  $(x_1, y_1)$  and  $(x_2, y_2)$ , where  $x_i \in \mathbb{Z}_p$  and  $y_i \in \mathbb{Z}_2$ . If  $y_1 \neq y_2$ , then both hexagons compose of an ideal configuration. Hence they are mutually resonant by Lemma 3.1. So we only consider the case  $x_1 = y_1 = y_2 = 1$  and  $3 \leq x_2 \leq \frac{p}{2} + 1 (\leq p - 1)$  by hexagon-preserving automorphisms of  $H(p, q, t)$ .

If  $t = p - 1, 1, 2, \dots, x_2 - 4$ , then we choose  $w_{20}b_{t+3,1}$  and  $w_{x_2+1,0}b_{x_2+2+t,1}$ . Since  $2 \leq t + 3 \leq x_2 - 1$  and  $x_2 + 1 \leq x_2 + 2 + t \leq 2x_2 - 2 \leq p$ , the end-vertices of both chosen edges in the 0-th and 1-st layers are separated by hexagons  $(1, 1)$  and  $(x_2, 1)$  (see Fig. 9). If  $t = x_2 - 2, x_2 - 1, \dots, p - 3$ , then we choose  $w_{20}b_{t+3,1}$  and  $w_{x,0}b_{x+1+t,1}$ , where  $x = x_2 - 2 - t + p$ . Since  $x_2 + 1 \leq 3 + t \leq p$ ,  $x_2 + 1 \leq x \leq p$  and  $x + 1 + t = p + x_2 - 1$ , the above result also holds. For the last case  $t = x_2 - 3$ , we have that  $x_2 \geq 4$  since  $t \neq 0$ . Since  $t + 1 = x_2 - 2 \geq 2$  and  $x_2 + 1 \leq x_2 + t = 2x_2 - 3 \leq p - 1$ , we choose  $w_{00}b_{t+1,1}$  and  $w_{x_2-1,0}b_{x_2+t,1}$ . For each subcase, the chosen hexagon  $(1, 1)$  and  $(x_2, 1)$  with the chosen edges form an ideal configuration. Therefore, by Lemma 3.1 the hexagons  $(1, 1)$  and  $(x_2, 1)$  are mutually resonant.  $\square$




Figure 9: Illustration for the proof of Lemma 3.4 for  $p = 8$ ,  $x_2 = 5$  and  $t = 1$ .

**Lemma 3.5** For  $q \geq 4$ ,  $H(2, q, t)$  is not 2-resonant.

**Proof:** Consider the disjoint hexagons  $h_{11}$  and  $h_{13}$ . The neighbors of the vertex  $w_{02}$ , which are  $b_{11}, b_{12}$  and  $b_{02}$ , belong to the chosen hexagons. Hence  $H(2, q, t) - h_{11} - h_{13}$  has no perfect matching. Namely hexagons  $h_{11}$  and  $h_{13}$  are not mutually resonant.  $\square$

From Lemma 3.2 we know that  $H(2, 2, 0)$  is not 1-resonant,  $H(2, 2, 1)$ ,  $H(3, 2, t)$  ( $t = 0, 1, 2$ ) and  $H(2, 3, t)$  ( $t = 0, 1$ ) are 1-resonant. We can see that  $H(2, 2, 1)$ ,  $H(3, 2, 0)$ ,  $H(3, 2, 1)$ ,  $H(2, 3, 0)$  and  $H(2, 3, 1)$  do not contain two disjoint hexagons. So automatically they are 2-resonant. Further  $H(3, 2, 2)$  is 2-resonant since it has exactly 12 vertices, which are all included in any pair of two disjoint hexagons. Combining these facts and Lemmas 3.3 to 3.5, we summarize the following characterization for the 2-resonance of toroidal polyhexes.

**Theorem 3.6** *For  $p \geq 2$  and  $q \geq 2$ , a toroidal polyhex  $H(p, q, t)$  is 2-resonant if and only if one of the following cases appears:*

1.  $\min(p, q) \geq 3$ ,
2.  $p \geq 4$ ,  $q = 2$  and  $t \notin \{0, p - 2\}$ ,
3.  $(p, q) = (3, 2)$  or  $(2, 3)$ ,
4.  $(p, q, t) = (2, 2, 1)$ .

## 4 3-Resonance

In this section we further consider the 3-resonance of toroidal polyhexes. The following lemma shows that most toroidal polyhexes are not 3-resonant.

**Lemma 4.1** *For  $p, q \geq 4$ ,  $H(p, q, t)$  is not 3-resonant.*

**Proof:** Choose three disjoint hexagons  $h_{11}$ ,  $h_{13}$  and  $h_{31}$ . Then vertices  $w_{11}$ ,  $w_{12}$  and  $w_{21}$  are neighbors of  $b_{21}$ . They lie in hexagons  $h_{11}$ ,  $h_{13}$  and  $h_{31}$ , respectively. Hence  $H(p, q, t) - h_{11} - h_{13} - h_{31}$  has an isolated vertex  $b_{21}$ . This implies that such three hexagons are not mutually resonant.  $\square$

**Lemma 4.2** *For  $q \geq 2$ ,  $H(3, q, t)$  is 3-resonant.*

**Proof:** By Lemma 3.3 or Theorem 3.6  $H(3, 2, t)$  and  $H(3, 3, t)$  are 2-resonant. We can easily see that they are 3-resonant since either they contain no three disjoint hexagons or their three disjoint hexagons contain all vertices. So we may assume  $q \geq 4$ .




Figure 10: An ideal configuration of  $H(3, 7, t)$  for (a)  $y_2 = 3, y_3 = 5$  and  $t = 1$ , (b)  $y_2 = 5, y_3 = 3$  and  $t = 0$ .

It suffices to show that any three disjoint hexagons  $(x_i, y_i)$ ,  $x_i \in \mathbb{Z}_3$  and  $y_i \in \mathbb{Z}_q$ ,  $i = 1, 2, 3$ , are mutually resonant. By l-r and t-b shift operations there are three cases to be considered.

**Case 1:**  $x_1 = 0, x_2 = 1, x_3 = 2$  and  $y_1 = 1$ . If  $2 \leq y_2 < y_3 \leq q - 1$ , then we choose vertical edges  $w_{0j}b_{1,j-1}, j = 2, \dots, y_2 - 1; w_{1j}b_{2,j-1}, j = y_2 + 1, \dots, y_3 - 1; w_{2j}b_{0,j-1}, j = y_3 + 1, \dots, q - 1$ ; we further choose one vertical edge  $w_{10}b_{t+2,q-1}$  when  $t = 1$  and 2, and  $w_{20}b_{0,q-1}$  when  $t = 0$  (see Fig. 10(a)). If  $2 < y_3 < y_2 \leq q - 1$ , then we choose vertical edges  $w_{0j}b_{1,j-1}, j = 2, \dots, y_3 - 1, y_3 + 1, \dots, y_2 - 1; w_{2j}b_{0,j-1}, j = y_2 + 1, \dots, q - 1$ ; we further choose one vertical edge  $w_{10}b_{t+2,q-1}$  when  $t = 0$  and 1, and  $w_{20}b_{2,q-1}$  when  $t = 2$  (see Fig. 10(b)). It can be seen that the three hexagons together with the chosen edges for any subcase form an ideal configuration. So by Lemma 3.1 such three hexagons are mutually resonant.

**Case 2:**  $x_1 = x_2 = x_3 = 1$  and  $1 = y_1 < y_2 < y_3 \leq q - 1$ . We choose vertical edges  $w_{2j}b_{0,j-1}, j = 2, \dots, y_2 - 1, y_2 + 1, \dots, y_3 - 1, y_3 + 1, \dots, q - 1$ . We further choose one vertical edge  $w_{00}b_{t+1,q-1}$  when  $t = 1$  and 2, and  $w_{20}b_{0,q-1}$  when  $t = 0$ . By the same reason as the above such three hexagons are mutually resonant.

**Case 3:**  $\{x_1, x_2, x_3\} = \{1, 2\}$  and  $1 = y_1 < y_2 < y_3 \leq q - 1$ . If  $x_1 = 1, x_2 = 2$  and  $x_3 = 1$ , then we choose vertical edges  $w_{1j}b_{2,j-1}, j = 2, \dots, y_2 - 1; w_{2j}b_{0,j-1}, j = y_2 + 1, \dots, y_3 - 1, y_3 + 1, \dots, q - 1$  ( $y_2 + 1 < y_3$ ). We further choose one vertical edge  $w_{00}b_{t+1,q-1}$  when  $t = 1$  and 2, and  $w_{20}b_{0,q-1}$  when  $t = 0$ . The chosen hexagons and edges form an ideal configuration. Similarly, we can show the remaining subcases to have an ideal configuration containing the chosen hexagons. Hence such three hexagons are mutually resonant.  $\square$

**Lemma 4.3** For  $p \geq 4$ ,  $H(p, 3, t)$  is 3-resonant if and only if  $t = 0, p - 3, p - 2$  or  $p - 1$ .

**Proof:** For  $p \geq 5$  and  $t \in \mathbb{Z}_p \setminus \{0, p - 3, p - 2, p - 1\}$ , we choose three disjoint hexagons  $(1, 1)$ ,  $(p - 1, 1)$  and  $(t + 1, 2)$ . Then the vertex  $w_{00}$ 's neighbors  $b_{10}, b_{00}$  and  $b_{t+1,2}$  are vertices in hexagons  $(1, 1)$ ,  $(p - 1, 1)$  and  $(t + 1, 2)$ , respectively. Hence the chosen hexagons are not mutually resonant. That says that  $H(p, 3, t)$  is not 3-resonant if  $t \notin \{0, p - 3, p - 2, p - 1\}$  for  $p \geq 4$ .

For  $t \in \{0, p - 3, p - 2, p - 1\}$ , we now show that  $H(p, 3, t)$  is 3-resonant. It suffices to show that any three disjoint hexagons  $(x_i, y_i), x_i \in \mathbb{Z}_p$  and  $y_i \in \mathbb{Z}_3, i = 1, 2, 3$ , are mutually resonant. By l-r and t-b shift operations there are three cases to be considered.

**Case 1:**  $y_1 = 0, y_2 = 1$  and  $y_3 = 2$ . Such three hexagons form an ideal configuration and are thus mutually resonant by Lemma 3.1.




Figure 11: An ideal configuration of  $H(8, 3, 0)$  for  $x_2 = 3, x_3 = 6$  and  $t = 0$ .

**Case 2:**  $y_1 = y_3 = 1, y_2 = 2$  and  $1 = x_1 < x_2 < x_3 \leq p - 1$ . If  $t = 0$  and  $p - 1$ , we choose  $w_{x_i-1,0}b_{x_i+t,2}, i = 1, 3$ , and  $w_{p-1,2}b_{0,1}$  as distinguished edges, see Fig. 11. We only list

the incident vertices of the chosen hexagons and vertical edges on the 2nd layer as  $b_{1+t,2}$ ,  $w_{x_2-1,2}$ ,  $b_{x_2,2}$ ,  $w_{x_2,2}$ ,  $b_{x_3+t,2}$ ,  $w_{p-1,2}$  in one direction of the cycle; If  $t = p - 2$  and  $p - 3$ , choose  $w_{x_i+1,0}b_{x_i+t+2,2}$ ,  $i = 1, 3$ , and  $w_{p-1,2}b_{0,1}$  as distinguished edges. The corresponding incident vertices are ordered as  $b_{3+t,2}$ ,  $w_{x_2-1,2}$ ,  $b_{x_2,2}$ ,  $w_{x_2,2}$ ,  $b_{x_3+t+2,2}$ ,  $w_{p-1,2}$  (note that  $x_2 < x_3 - 1$ ). As for the 0-th and 1-st layers, the similar facts are obvious. Hence to each subcase the three hexagons and all chosen edges compose of an ideal configuration. By Lemma 3.1 such three hexagons are mutually resonant.

**Case 3:**  $y_1 = y_2 = y_3 = 1$  and  $1 = x_1 < x_2 < x_3 \leq p - 1$ . We choose vertical edges  $w_{x_i,2}b_{x_i+1,1}$ ,  $i = 1, 2, 3$ . If  $t = 0, p - 1$ , we further choose  $w_{x_i-1,0}b_{x_i+t,2}$ ,  $i = 1, 2, 3$ . If  $t = p - 2, p - 3$ , we further choose  $w_{x_i+1,0}b_{x_i+t+2,2}$ ,  $i = 1, 2, 3$ . To each subcase the three hexagons and all chosen edges compose of an ideal configuration. By Lemma 3.1 such three hexagons are mutually resonant.  $\square$

**Lemma 4.4** For  $p \geq 3$ ,  $H(p, 2, t)$  is 3-resonant if and only if  $t = 1, p - 3$  or  $p - 1$ .

**Proof:** For  $t \in \mathbb{Z}_p \setminus \{1, p - 3, p - 1\}$  ( $p \geq 4$ ), we choose disjoint hexagons  $(1, 1)$ ,  $(t + 1, 1)$  and  $(p - 1, 1)$  of  $H(p, 2, t)$  (note that if  $t = 0$  and  $t = p - 2$ , then two of them are coincidence). The vertex  $w_{00}$  has three neighbors:  $b_{10}$ ,  $b_{00}$  and  $b_{t+1,1}$ , which are all included in the chosen hexagons. This implies that  $H(p, 2, t) - h_{1,1} - h_{t+1,1} - h_{p-1,1}$  has no perfect matching. Hence  $H(p, 2, t)$  ( $p \geq 3$ ) is not 3-resonant for  $t \notin \{1, p - 3, p - 1\}$ .

We now show that if  $t \in \{1, p - 1, p - 3\}$ ,  $H(p, 2, t)$  ( $p \geq 3$ ) is 3-resonant. Since it is 2-resonant (Lemma 3.4 or Theorem 3.6), it suffices to show any three disjoint hexagons  $(x_i, y_i)$ ,  $x_i \in \mathbb{Z}_p$  and  $y_i \in \mathbb{Z}_2$ ,  $i = 1, 2, 3$ , are mutually resonant. By the l-r and t-b shift operations (Eqs. (1) to (3)) there are two cases to be considered.

**Case 1:**  $y_1 = y_2 = y_3 = 1$  and  $1 = x_1 < x_2 < x_3 \leq p - 1$ . If  $t = 1$  or  $p - 1$ , then we choose the edges  $w_{x_i-1,0}b_{x_i+t,1}$ ,  $i = 1, 2, 3$ ; if  $t = p - 3$ , then we choose the edges  $w_{x_i+1,0}b_{x_i+t+2,1}$  for  $i = 1, 2, 3$ . Then the chosen hexagons and edges form an ideal configuration and such three hexagons are thus mutually resonant by Lemma 3.1.

**Case 2:**  $y_1 = y_3 = 1, y_2 = 0$  and  $1 = x_1 < x_2 < x_3 \leq p - 1$ . If  $t = p - 1$ , then we only choose the edge  $w_{00}b_{01}$  (see Fig. 12 (right)). If  $t = 1$ , then  $x_3 \geq x_2 + 3$ . We choose edges  $w_{x_i-1,0}b_{x_i+1,1}$ , for  $i = 1, 3$ , and  $w_{x_2,1}b_{x_2+1,0}$  (see Fig. 12 (left)). If  $t = p - 3$ , then  $x_2 - x_1 \geq 4$ . We choose the edges  $w_{x_i+1,0}b_{x_i-1,1}$  for  $i = 1, 3$ , and  $w_{x_2-2,1}b_{x_2-1,0}$ . In any subcase the chosen hexagons and vertical edges form an ideal configuration and such chosen hexagons are mutually resonant by Lemma 3.1.  $\square$




Figure 12: Illustration for Case 2 in the proof of Lemma 4.4 ( $t = 1$  and  $t = p - 1$ ).

Automatically  $H(2, 3, t)$  and  $H(2, 2, 1)$  are 3-resonant. Combining Lemmas 3.5 and 4.1 to 4.4, we obtain the following criterion for the 3-resonance of toroidal polyhexes.

**Theorem 4.5** For  $p \geq 2$  and  $q \geq 2$ , a toroidal polyhex  $H(p, q, t)$  is 3-resonant if and only if one of the following cases appears

1.  $(p, q, t) = (2, 2, 1)$ ,
2.  $p = 2$  and  $q = 3$ ,
3.  $p = 3$  and  $q \geq 2$ ,
4.  $p \geq 4$ ,  $q = 2$  and  $t \in \{1, p - 3, p - 1\}$ ,
5.  $p \geq 4$ ,  $q = 3$  and  $t \in \{0, p - 3, p - 2, p - 1\}$ .

## References

- [1] G.G. Cash, Simple means of computing the Kekulé structure count for toroidal polyhex fullerenes, *J. Chem. Inf. Comput. Sci.*, **38** (1998), 58-61.
- [2] R. Chen, X. Guo,  $k$ -coverable coronoid systems, *J. Math. Chem.*, **12** (1993), 147-162.
- [3] E. Clar, *The Aromatic Sextet*, (Wiley: London, 1972).
- [4] M. Deza, P.W. Fowler, A. Rassat and K.M. Rogers, Fullerenes as tilings of surfaces, *J. Chem. Inf. Comput. Sci.*, **40** (2000), 550-558.
- [5] X. Guo and F. Zhang,  $k$ -resonant benzenoid systems and  $k$ -cycle resonant graphs, *J. Chem. Inf. Comput. Sci.*, **41** (3) (2001), 480-483.
- [6] I. Gutman, Covering hexagonal systems with hexagons, *Proc. 4th Yugoslav Seminar on Graph Theory*, Novi Sad, 1983, 151-159.
- [7] H. Hosoya and T. Yamaguchi, Sextet polynomials. A new enumeratin and proof technique for the resonance theory applied to the aromatic hydrocarbons, *Tetrahedron Lett.*, **52** (1975), 4659-4662.
- [8] P.E. John, Kekulé count in toroidal hexagonal carbon cages, *Croat. Chem. Acta*, **71** (3) (1998), 435-447.
- [9] E.C. Kirby, R.B. Mallion and P. Pollak, Toridal polyhexes, *J. Chem. Soc. Faraday Trans.*, **89** (12) (1993), 1945-1953.
- [10] E.C. Kirby and P. Pollak, How to enumerate the connectional isomers of a toridal polyhex fullerene, *J. Chem. Inf. Comput. Sci.*, **38** (1998), 66-70.
- [11] D.J. Klein, Elemental benzenoids, *J. Chem. Inf. Comput. Sci.*, **34** (1994), 453-459.
- [12] D.J. Klein and H. Zhu, Resonance in elemental benzenoids, *Discrete Appl. Math.*, **67** (1996), 157-173.
- [13] K. Lin and R. Chen,  $k$ -coverable polyhex graphs, *Ars Combin.*, **43** (1996), 33-48.
- [14] J. Liu, H.J. Dai, J.H. Hafner, D.T. Colbert, D.T. R.E. Smalley, S.J. Tans and C. Dekker, Fullerene 'crop circles', *Nature*, **385** (1997), 780-781.
- [15] L. Lovász and M.D. Plummer, *Matching Theory, Annals of Discrete Math.* **29**, (North-Holland, Amsterdam, 1986).

- [16] D. Marušič and T. Pisanski, Symmetries of hexagonal molecular graphs on the torus, *Croat. Chem. Acta*, **73** (4) (2000), 969-981.
- [17] S. Negami, Uniqueness and faithfulness of embedding of toroidal graphs, *Discrete Math.*, **44** (1983), 161-180.
- [18] M. Randić, Aromaticity of polycyclic conjugated hydrocarbons, *Chem. Reviews*, **103** (9) (2003), 3449-3605.
- [19] M. Randić, Conjugated circuits and resonance energies of benzenoid hydrocarbons, *Chem. Phys. Lett.*, **38** (1976), 68-70.
- [20] M. Randić, Aromaticity and conjugation, *J. Am. Chem. Soc.*, **99** (1977), 444-450.
- [21] T.G. Schmalz, W.A. Seitz, D.J. Klein and G.E. Hite, Elemental carbon cages, *J. Am. Chem. Soc.*, **110** (1988), 1113-1127.
- [22] C. Thomassen, Tilings of the torus and the Klein bottle and vertex-transitive graphs on a fixed surface, *Trans. Amer. Math. Soc.*, **323** (1991), 605-635.
- [23] F. Zhang and L. Wang,  $k$ -resonance of open-ended carbon nanotubes, *J. Math. Chem.*, **35** (2) (2004), 87-103.
- [24] F. Zhang and R. Chen, When each hexagon of a hexagonal system covers it, *Discrete Appl. Math.*, **30** (1991), 63-75.
- [25] F. Zhang and M. Zheng, Generalized hexagonal systems with each hexagon being resonant, *Discrete Appl. Math.*, **36** (1992), 67-73.
- [26] H. Zhang and F. Zhang, The Clar covering polynomial of hexagonal systems I, *Discrete Appl. Math.*, **69** (1996), 147-167.
- [27] H. Zhang and F. Zhang, Plane elementary bipartite graphs, *Discrete Appl. Math.*, **105** (2000), 291-311.
- [28] M. Zheng,  $k$ -resonant benzenoid systems, *J. Mol. Struct. (Theochem)*, **231** (1991), 321-334.
- [29] M. Zheng, Construction of 3-resonant benzenoid systems, *J. Mol. Struct. (Theochem)*, **277** (1992), 1-14.