

On localizing and concentrating electromagnetic fields

Harrach, Bastian; Lin, Yi Hsuan; LIU, Hongyu

Published in:
SIAM Journal on Applied Mathematics

DOI:
[10.1137/18M1173605](https://doi.org/10.1137/18M1173605)

Published: 01/01/2018

[Link to publication](#)

Citation for published version (APA):
Harrach, B., Lin, Y. H., & LIU, H. (2018). On localizing and concentrating electromagnetic fields. *SIAM Journal on Applied Mathematics*, 78(5), 2558-2574. <https://doi.org/10.1137/18M1173605>

General rights

Copyright and intellectual property rights for the publications made accessible in HKBU Scholars are retained by the authors and/or other copyright owners. In addition to the restrictions prescribed by the Copyright Ordinance of Hong Kong, all users and readers must also observe the following terms of use:

- Users may download and print one copy of any publication from HKBU Scholars for the purpose of private study or research
- Users cannot further distribute the material or use it for any profit-making activity or commercial gain
- To share publications in HKBU Scholars with others, users are welcome to freely distribute the permanent publication URLs

ON LOCALIZING AND CONCENTRATING ELECTROMAGNETIC FIELDS*

BASTIAN HARRACH[†], YI-HSUAN LIN[‡], AND HONGYU LIU[§]

Abstract. We consider field localizing and concentration of electromagnetic waves governed by the time-harmonic anisotropic Maxwell system in a bounded domain. It is shown that there always exist certain boundary inputs which can generate electromagnetic fields with the energy localized/concentrated in a given subdomain while nearly vanishing in another given subdomain. The theoretical results may have potential applications in telecommunication, inductive charging, and medical therapy. We also derive a related Runge approximation result for the time-harmonic anisotropic Maxwell system with partial boundary data.

Key words. electromagnetic waves, localizing and concentration, anisotropic Maxwell system, Runge approximation, partial data

AMS subject classifications. Primary, 35Q61; Secondary, 78A25, 78A45

DOI. 10.1137/18M1173605

1. Introduction.

1.1. Background and motivation. The electromagnetic (EM) phenomena are ubiquitous and they lie at the heart of many scientific and technological applications including radar and sonar, geophysical exploration, medical imaging, information processing, and communication. In this paper, we are mainly concerned with the mathematical study of field localizing and concentration of electromagnetic waves governed by the time-harmonic Maxwell system in a bounded anisotropic medium. More specifically, we show that there always exist certain boundary inputs which can generate the desired electromagnetic fields that are localized/concentrated in a given subdomain while nearly vanishing in another given subdomain.

The localizing and concentration of electromagnetic fields can have many potential applications. In telecommunication [44], one common means of transmitting information between communication participants is via the electromagnetic radiation. In a certain practical scenario, say, secure communication, one may intend the information to be transmitted mainly to a partner located at a certain region, while avoiding the transmission to another region. Clearly, if the information is encoded into the electromagnetic waves that are localized and concentrated in the region where the partner is located while nearly vanishing in the undesired region, then one can achieve the expected telecommunication effect. In the setup of our proposed study, one can easily obtain the aforementioned communication effect, in particular if the communication participants transmit and receive information on some surface patches.

*Received by the editors March 2, 2018; accepted for publication (in revised form) July 23, 2018; published electronically September 25, 2018.

<http://www.siam.org/journals/siap/78-5/M117360.html>

Funding: The third author's work was supported by the FRG and startup grants from Hong Kong Baptist University, and Hong Kong RGC General Research Funds, 12302415 and 12302017.

[†]Institute for Mathematics, Goethe University Frankfurt, Frankfurt, 60325, Germany (harrach@math.uni-frankfurt.de).

[‡]Institute for Advanced Study, The Hong Kong University Science and Technology, Clear Water Bay, Kowloon, Hong Kong (yihhsuanlin3@gmail.com).

[§]Department of Mathematics, Hong Kong Baptist University, Kowloon, Hong Kong SAR, China (hongyu.liuip@gmail.com, hongyuliu@hkbu.edu.hk).

Concentrating electromagnetic fields can also be useful in inductive charging, also known as wireless charging or cordless charging [43], which is an emerging technology that can have significant impact on real life. It uses electromagnetic fields to transfer energy between two objects through electromagnetic induction. Clearly, the energy transfer would be more efficient and effective if the corresponding electromagnetic fields are concentrated around the charging station. The localizing of electromagnetic fields can also have potential application in electromagnetic therapy. Though it is mainly considered to be pseudoscientific with no affirmative evidence, electromagnetic therapy has been widely practiced and claims to treat disease by applying electromagnetic radiation to the body. If electromagnetic therapy shall be proven to be effective, then through the use of certain purposely designed sources, one can generate electromagnetic fields that are concentrated around the diseased area.

The above conceptual and potential applications make the study of field concentration and localizing very appealing. Nevertheless, it is emphasized that in the current article, we are mainly concerned with mathematical and theoretical study. We achieve some substantial progress on this interesting topic, though the corresponding study is by no means complete. It is also interesting to note that the localizing of resonant electromagnetic fields has been used to produce invisibility cloaking and has received significant attention in the literature in recent years [2, 3, 6, 31, 34, 37]. The corresponding study is mainly based on the use of plasmonic materials to induce so-called anomalous localized resonance.

Our mathematical argument for proving the existence of localized and concentrated electromagnetic fields is mainly based on combining the unique continuation property (UCP) for the anisotropic Maxwell system with a functional analytic duality argument developed in [12]. By a similar argument, we also obtain a related Runge approximation property.

The use of blow up solutions has a long tradition in the study of inverse boundary value problems; cf. [1, 25, 27, 28] for early seminal works on this topic. Moreover, the combination of localized fields and monotonicity relations has led to the development of monotonicity-based methods for obstacle/inclusion detection; cf. [22, 39] for the origins and mathematical justification of this approach, [5, 7, 9, 10, 11, 16, 17, 18, 19, 21, 23, 33, 38, 40, 42, 45] for further recent contributions, and the recent works [13, 20] for the Helmholtz equation. Theoretical uniqueness results for inverse coefficient problems have also been obtained by this approach in [4, 14, 15, 21, 24].

In this work, we show the existence of localized electromagnetic fields for the more challenging case of a time-harmonic anisotropic Maxwell system with partial data. We also derive a Runge approximation result, which shows that every solution in a subdomain can be approximated by a solution on the whole domain. In that context let us note the famous equivalence theorem from Peter Lax [29]: the weak UCP is equivalent to the Runge approximation property for the second order elliptic equation. In our study, we affirmatively verify that this property still holds for the anisotropic Maxwell system.

The rest of this section is devoted to the mathematical description of the setup of our study and the statement of the main result.

1.2. Mathematical setup and statement of the main result. Let Ω be a simply connected domain in \mathbb{R}^3 with a Lipschitz connected boundary $\partial\Omega$. Let $\epsilon = (\epsilon_{ij})_{1 \leq i, j \leq 3}$ and $\mu = (\mu_{ij})_{1 \leq i, j \leq 3}$ be two 3×3 real matrix-valued functions on Ω satisfying the following:

- Strong ellipticity: There exist constants $\mu_0 > 0$ and $\epsilon_0 > 0$ verifying

$$(1.1) \quad \begin{cases} \mu_0 |\xi|^2 \leq \sum_{i,j=1}^3 \mu_{ij}(x) \xi_i \xi_j \leq \mu_0^{-1} |\xi|^2, \\ \epsilon_0 |\xi|^2 \leq \sum_{i,j=1}^3 \epsilon_{ij}(x) \xi_i \xi_j \leq \epsilon_0^{-1} |\xi|^2 \end{cases} \quad \text{for any } x \in \Omega \text{ and } \xi \in \mathbb{R}^3.$$

- Smoothness: ϵ and μ are Lipschitz continuous.
- Symmetry: ϵ and μ are symmetric matrices, that is, $\epsilon_{ij} = \epsilon_{ji}$ and $\mu_{ij} = \mu_{ji}$ for all $i, j = 1, 2, 3$.

The functions ϵ and μ , respectively, signify the electric permittivity and magnetic permeability of the medium in Ω . Consider the time-harmonic electromagnetic wave propagation in Ω . With the e^{-ikt} time-harmonic convention assumed, we let $E(x)$ and $H(x)$, respectively, denote the electric and magnetic fields. Here, $k \in \mathbb{R}_+$ signifies a circular frequency. Then the electromagnetic wave propagation is governed by the following Maxwell system:

$$(1.2) \quad \begin{cases} \nabla \times E - ik\mu H = 0 & \text{in } \Omega, \\ \nabla \times H + ik\epsilon E = 0 & \text{in } \Omega, \\ \nu \times E = \begin{cases} f & \text{on } \Gamma, \\ 0 & \text{otherwise} \end{cases} & \text{on } \partial\Omega, \end{cases}$$

where Γ is an arbitrary nonempty relatively open subset of $\partial\Omega$ and ν is the unit outer normal vector on $\partial\Omega$. It is assumed that $k > 0$ is not an eigenvalue (or nonresonance, see section 2) for (1.2) and $f \in C_c^\infty(\Gamma)$ throughout this paper.

The main result concerning the localized electromagnetic fields for the anisotropic Maxwell system (1.2) is contained in the following theorem.

THEOREM 1.1. *Let $\Omega \subset \mathbb{R}^3$ be a bounded Lipschitz domain and $\Gamma \subseteq \partial\Omega$ be a relatively open subset of the boundary. Let $\epsilon, \mu \in L^\infty(\Omega, \mathbb{R}^{3 \times 3})$ be real-valued, piecewise Lipschitz continuous functions satisfying (1.1) and $k > 0$ be a nonresonant frequency. Let $D \Subset \Omega$ be a closed set with a connected complement $\Omega \setminus D$. For every open set $M \subseteq \Omega$ with $M \not\subseteq D$ (see Figure 1 for the schematic illustration), there exists a sequence $\{f^{(\ell)}\}_{\ell \in \mathbb{N}} \subset C_c^\infty(\Gamma)$ such that the electromagnetic fields fulfill*

$$\int_M (|E^{(\ell)}|^2 + |H^{(\ell)}|^2) dx \rightarrow \infty \quad \text{and} \quad \int_D (|E^{(\ell)}|^2 + |H^{(\ell)}|^2) dx \rightarrow 0 \quad \text{as } \ell \rightarrow \infty,$$




FIG. 1. Schematic illustration of the field localizing and concentration.

where, for $\ell \in \mathbb{N}$, $(E^{(\ell)}, H^{(\ell)}) \in H(\text{curl}, \Omega) \times H(\text{curl}, \Omega)$ is a solution of

$$\begin{cases} \nabla \times E^{(\ell)} - ik\mu H^{(\ell)} = 0 & \text{in } \Omega, \\ \nabla \times H^{(\ell)} + ik\epsilon E^{(\ell)} = 0 & \text{in } \Omega \end{cases}$$

with the boundary data

$$\nu \times E^{(\ell)}|_{\partial\Omega} = \begin{cases} f^{(\ell)} & \text{on } \Gamma, \\ 0 & \text{otherwise.} \end{cases}$$

Remark 1.1. We denote the sequence $\{(E^{(\ell)}, H^{(\ell)})\}_{\ell \in \mathbb{N}}$ in Theorem 1.1 to be the localized electromagnetic fields.

The rest of the paper is organized as follows. In section 2, we present results on the well-posedness of the time-harmonic anisotropic Maxwell system. We also provide the UCP for the anisotropic Maxwell system, whenever the coefficients μ and ϵ are piecewise Lipschitz continuous matrix-valued functions. In section 3, we demonstrate that there exist localized electromagnetic fields, which proves Theorem 1.1. The method relies on certain functional analysis techniques. In section 4 we prove a related Runge approximation property for the anisotropic Maxwell system with partial boundary data.

2. The anisotropic Maxwell system in a bounded domain. In this section, we summarize some useful results of the Maxwell system, including the unique solvability and a UCP. Throughout this section we let $\Omega \subset \mathbb{R}^3$ be a bounded Lipschitz domain.

2.1. Spaces and traces. We introduce the spaces

$$\begin{aligned} H(\text{div}, \Omega) &:= \{E \in L^2(\Omega)^3; \quad \nabla \cdot E \in L^2(\Omega)\}, \\ H(\text{curl}, \Omega) &:= \{E \in L^2(\Omega)^3; \quad \nabla \times E \in L^2(\Omega)^3\}, \end{aligned}$$

and the tangential trace operators

$$\begin{aligned} \gamma_t : H(\text{curl}, \Omega) &\rightarrow H^{-1/2}(\text{div}_{\partial\Omega}, \partial\Omega), \quad E \mapsto \gamma_t E := \nu \times E|_{\partial\Omega}, \\ \gamma_T : H(\text{curl}, \Omega) &\rightarrow H^{-1/2}(\text{curl}_{\partial\Omega}, \partial\Omega), \quad E \mapsto \gamma_T E := \nu \times (E|_{\partial\Omega} \times \nu), \end{aligned}$$

where, and also in what follows, all functions are complex-valued unless indicated otherwise. γ_t and γ_T are surjective bounded linear operators with bounded right inverses γ_t^{-1} and γ_T^{-1} (cf. [8, 41]). The space $H^{-1/2}(\text{div}_{\partial\Omega}, \partial\Omega)$ can be identified with the dual of $H^{-1/2}(\text{curl}_{\partial\Omega}, \partial\Omega)$, and for all $E, F \in H(\text{curl}, \Omega)$ we have the integration by parts formula

$$(2.1) \quad \int_{\Omega} (\nabla \times E) \cdot F \, dx - \int_{\Omega} E \cdot (\nabla \times F) \, dx = \int_{\partial\Omega} (\nu \times E|_{\partial\Omega}) \cdot (\nu \times (F|_{\partial\Omega} \times \nu)) \, dS$$

(cf. [8, 35]), where the dual pairing on $H^{-1/2}(\text{div}_{\partial\Omega}, \partial\Omega) \times H^{-1/2}(\text{curl}_{\partial\Omega}, \partial\Omega)$ is written as an integral for notational convenience.

The subspace of $H(\text{curl}, \Omega)$ -functions with vanishing tangential traces is denoted by

$$H_0(\text{curl}, \Omega) := \{E \in H(\text{curl}, \Omega) : \nu \times E|_{\partial\Omega} = 0\}.$$

$H_0(\text{curl}, \Omega)$ is a closed subspace of $H(\text{curl}, \Omega)$ and $C_0^\infty(\Omega)^3$ is dense in $H_0(\text{curl}, \Omega)$ (cf. [35]).

To treat partial boundary data on a relatively open subset $\Gamma \subseteq \partial\Omega$, we also introduce the space of functions on Γ that can be extended by zero to the trace of a $H(\text{curl}, \Omega)$ -function

$$(2.2) \quad H(\Gamma) := \text{closure of } C_c^\infty(\Gamma) \text{ in } H^{-1/2}(\text{div}_{\partial\Omega}, \partial\Omega).$$

For all $E \in H(\text{curl}, \Omega)$, we identify the restricted trace $\nu \times (E|_\Gamma \times \nu)$ with the quotient space element

$$\nu \times (E \times \nu)|_{\partial\Omega} + H(\Gamma)^\perp \in H^{-1/2}(\text{curl}_{\partial\Omega}, \partial\Omega)/H(\Gamma)^\perp = H(\Gamma)^*,$$

and thus define the restricted trace operator

$$\gamma_T^{(\Gamma)} : H(\text{curl}, \Omega) \rightarrow H(\Gamma)^*, \quad E \mapsto \gamma_T^{(\Gamma)} E := \nu \times (E|_\Gamma \times \nu).$$

2.2. Well-posedness of the anisotropic Maxwell system. Given anisotropic coefficients $\epsilon, \mu \in L^\infty(\Omega, \mathbb{R}^{3 \times 3})$ satisfying (1.1), $k > 0$, $J, K \in L^2(\Omega)^3$, and $f \in H^{-1/2}(\text{div}_{\partial\Omega}, \partial\Omega)$, we consider the Maxwell system for $(E, H) \in H(\text{curl}, \Omega) \times H(\text{curl}, \Omega)$ such that

$$(2.3) \quad \nabla \times E - ik\mu H = K \quad \text{in } \Omega,$$

$$(2.4) \quad \nabla \times H + ik\epsilon E = J \quad \text{in } \Omega,$$

$$(2.5) \quad \nu \times E|_{\partial\Omega} = f.$$

For the variational formulation of (2.3)–(2.4) we introduce the sesquilinear form

$$\begin{aligned} \mathcal{B} : H(\text{curl}, \Omega) \times H(\text{curl}, \Omega) &\rightarrow \mathbb{C}, \\ \mathcal{B}(E, F) &:= \int_\Omega (\mu^{-1} \nabla \times E) \cdot (\nabla \times \bar{F}) \, dx - \int_\Omega k^2 \epsilon E \cdot \bar{F} \, dx. \end{aligned}$$

Then we have the following variational formulation and well-posedness result.

THEOREM 2.1.

(a) $(E, H) \in H(\text{curl}, \Omega) \times H(\text{curl}, \Omega)$ solve (2.3)–(2.4) if and only if $E \in H(\text{curl}, \Omega)$ solves

$$\mathcal{B}(E, F) = \int_\Omega ikJ \cdot \bar{F} \, dx + \int_\Omega (\mu^{-1} K) \cdot (\nabla \times \bar{F}) \, dx \quad \text{for all } F \in H_0(\text{curl}, \Omega),$$

$$\text{and } H = -\frac{i}{k} \mu^{-1} (\nabla \times E - K).$$

(b) The set of $k > 0$ for which the homogeneous system (2.3)–(2.5) with $J = 0$, $K = 0$, and $f = 0$ possesses a nontrivial solution is discrete. We call these k resonance frequencies.

(c) If k is not a resonance frequency, then there exists a unique solution $(E, H) \in H(\text{curl}; \Omega) \times H(\text{curl}; \Omega)$ of (2.3)–(2.5), and the solution depends linearly and continuously on $J, K \in L^2(\Omega)^3$ and $f \in H^{-1/2}(\text{div}_{\partial\Omega}, \partial\Omega)$.

The proof of Theorem 2.1 follows from a standard argument. Since we couldn't find a convenient reference for precisely this setting, we supply a proof for the sake of completeness. We first derive the following lemma.

LEMMA 2.1. $(E, H) \in H(\text{curl}, \Omega) \times H(\text{curl}, \Omega)$ solves (2.3)–(2.4) if and only if $E \in H(\text{curl}, \Omega)$ solves

$$(2.6) \quad \mathcal{B}(E, F) = \int_{\Omega} ikJ \cdot \bar{F} \, dx + \int_{\Omega} (\mu^{-1}K) \cdot (\nabla \times \bar{F}) \, dx \quad \text{for all } F \in H_0(\text{curl}, \Omega),$$

and $H = -\frac{i}{k}\mu^{-1}(\nabla \times E - K)$.

Proof. Let $(E, H) \in H(\text{curl}, \Omega) \times H(\text{curl}, \Omega)$ solve (2.3)–(2.4). Then (2.3) implies that

$$E \in H(\text{curl}, \Omega) \quad \text{and} \quad H = -\frac{i}{k}\mu^{-1}(\nabla \times E - K),$$

and combining (2.3) and (2.4) we obtain

$$(2.7) \quad \nabla \times (\mu^{-1}(\nabla \times E - K)) - k^2\epsilon E = ikJ,$$

which also shows that $\mu^{-1}(\nabla \times E - K) \in H(\text{curl}; \Omega)$. Using (2.7) and the integration by parts formula (2.1), it follows that for all $F \in H_0(\text{curl}; \Omega)$

$$\begin{aligned} \int_{\Omega} ikJ \cdot \bar{F} \, dx &= \int_{\Omega} \left(\nabla \times \left(\frac{1}{\mu}(\nabla \times E - K) \right) \right) \cdot \bar{F} \, dx - \int_{\Omega} k^2\epsilon E \cdot \bar{F} \, dx \\ &= \int_{\Omega} (\mu^{-1}(\nabla \times E - K)) \cdot (\nabla \times \bar{F}) \, dx - \int_{\Omega} k^2\epsilon E \cdot \bar{F} \, dx, \end{aligned}$$

and thus (2.6) holds.

On the other hand, if $E \in H(\text{curl}, \Omega)$ fulfills (2.6) for all $F \in H_0(\text{curl}, \Omega)$, then this also holds for all $F \in C_0^\infty(\Omega)^3$, which (by the definition of distributional derivatives) shows that

$$\nabla \times (\mu^{-1}\nabla \times E) - k^2\epsilon E = ikJ + \nabla \times (\mu^{-1}K),$$

and thus

$$\frac{1}{ik}\nabla \times (\mu^{-1}(\nabla \times E - K)) + ik\epsilon E = J.$$

Defining $H := \frac{1}{ik}\mu^{-1}(\nabla \times E - K)$ it follows that $H \in H(\text{curl}; \Omega)$ and that E and H solve (2.3)–(2.4).

The proof is complete. □

If $H(\text{curl}, \Omega)$ was compactly embedded into $L^2(\Omega)$, then Theorem 2.1 would immediately follow from Lemma 2.1 by a Fredholm argument. However this is not the case, and we need to introduce an additional variational formulation on the space

$$\mathcal{H} := \{E \in L^2(\Omega)^3 : \nabla \times E \in L^2(\Omega)^3, \nabla \cdot (\epsilon E) = 0, \nu \times E|_{\partial\Omega} = 0\},$$

which is compactly embedded into $L^2(\Omega)^3$ (see, e.g., [26, Theorem 5.32]). We now first consider the Maxwell system with homogeneous boundary data and divergence free electric currents, so that the solution lies in \mathcal{H} . After that we shall show that the general Maxwell system can be transformed (or gauged) to fulfill this condition.

LEMMA 2.2. For $f = 0$ and $\nabla \cdot J = 0$, $(E, H) \in H(\text{curl}, \Omega) \times H(\text{curl}, \Omega)$ solves (2.3)–(2.5) if and only if $E \in \mathcal{H}$ solves

$$(2.8) \quad \mathcal{B}(E, F) = \int_{\Omega} ikJ \cdot \bar{F} \, dx + \int_{\Omega} (\mu^{-1}K) \cdot (\nabla \times \bar{F}) \, dx \quad \text{for all } F \in \mathcal{H},$$

$$\text{and } H = -\frac{i}{k}\mu^{-1}(\nabla \times E - K).$$

Proof. If $(E, H) \in H(\text{curl}, \Omega) \times H(\text{curl}, \Omega)$ fulfill (2.3)–(2.5), then clearly $E \in \mathcal{H}$ and Lemma 2.1 shows that (2.8) is fulfilled for all $F \in \mathcal{H} \subset H_0(\text{curl}, \Omega)$.

To prove the other direction, let $E \in \mathcal{H}$ fulfill (2.8) for all $F \in \mathcal{H}$. Given $\Phi \in H_0(\text{curl}, \Omega)$, there exists a solution $\varphi \in H_0^1(\Omega)$ of

$$\nabla \cdot (\epsilon \nabla \varphi) = -\nabla \cdot (\epsilon \Phi)$$

and thus $F := \Phi + \nabla \varphi \in \mathcal{H}$. Using (2.8) it follows that for

$$\begin{aligned} \mathcal{B}(E, \Phi) &= \mathcal{B}(E, F) - \mathcal{B}(E, \nabla \varphi) \\ &= \int_{\Omega} ikJ \cdot \bar{F} \, dx + \int_{\Omega} (\mu^{-1}K) \cdot (\nabla \times \bar{F}) \, dx + \int_{\Omega} k^2 \epsilon E \cdot \nabla \bar{\varphi} \, dx \\ &= \int_{\Omega} ikJ \cdot \bar{\Phi} \, dx + \int_{\Omega} (\mu^{-1}K) \cdot (\nabla \times \bar{\Phi}) \, dx, \end{aligned}$$

where we used $\nabla \times (\nabla \varphi) = 0$, $\nabla \cdot J = 0$, $\nabla \cdot (\epsilon E) = 0$, and $\varphi|_{\partial\Omega} = 0$.

The proof is complete. \square

We recall that we call $k > 0$ a resonant frequency, if the homogeneous Maxwell system (2.3)–(2.5) with $J = 0$, $K = 0$, and $f = 0$ admits a nontrivial solution.

LEMMA 2.3. If $k > 0$ is not a resonant frequency, then for every $J, K \in L^2(\Omega)^3$ with $\nabla \cdot J = 0$ and $f = 0$, there exists a unique solution $(E, H) \in H(\text{curl}, \Omega) \times H(\text{curl}, \Omega)$ of (2.3)–(2.5), and the solution depends continuously on $J, K \in L^2(\Omega)^3$. Moreover, the set of resonant frequencies is discrete.

Proof. Lemma 2.2 yields that $(E, H) \in H(\text{curl}, \Omega) \times H(\text{curl}, \Omega)$ solves (2.3)–(2.5) if and only if

$$(\mathcal{A} + \mathcal{K}(k))E = l,$$

where $\mathcal{A}, \mathcal{K}(k) : \mathcal{H} \rightarrow \mathcal{H}^*$ are defined by

$$\begin{aligned} \langle \mathcal{A}E, F \rangle &:= \int_{\Omega} (\mu^{-1}\nabla \times E) \cdot (\nabla \times F) \, dx + \int_{\Omega} E \cdot F \, dx \quad \text{for all } E, F \in \mathcal{H}, \\ \langle \mathcal{K}(k)E, F \rangle &:= -\int_{\Omega} (1 + k^2\epsilon)E \cdot F \, dx \quad \text{for all } E, F \in \mathcal{H}, \end{aligned}$$

and $l \in \mathcal{H}^*$ is defined by

$$\langle l, F \rangle = \int_{\Omega} ikJ \cdot F \, dx + \int_{\Omega} (\mu^{-1}K) \cdot (\nabla \times F) \, dx \quad \text{for all } F \in \mathcal{H},$$

where \mathcal{H}^* is the dual space of \mathcal{H} .

Then \mathcal{A} is a coercive linear bounded operator and thus continuously invertible due to the Lax–Milgram theorem. For every $k \in \mathbb{C}$, $\mathcal{K}(k) : \mathcal{H} \rightarrow \mathcal{H}^*$ is a linear compact operator due to the compact embedding of \mathcal{H} into $L^2(\Omega)^3$. $l \in \mathcal{H}^*$ depends linearly

and continuously on $J, K \in L^2(\Omega)^3$. It thus follows from the Fredholm alternative that $\mathcal{A} + \mathcal{K}(k)$ is continuously invertible if it is injective, i.e., if $k > 0$ is not a resonant frequency.

Moreover, $\mathcal{K}(k)$ depends analytically on k , and for $\widehat{k} := i$, $\mathcal{A} + \mathcal{K}(\widehat{k})$ is coercive and thus continuously invertible. Hence, it follows from the analytic Fredholm theorem that the set of resonances is discrete.

The proof is complete. □

Next we extend this result to nonhomogeneous boundary data f and non-divergence-free currents J and prove Theorem 2.1.

Proof of Theorem 2.1. Part (a) follows from Lemma 2.1. Part (b) and the uniqueness of the solution of the Maxwell system are proved in Lemma 2.3. To prove the existence of the solution, we let $J, K \in L^2(\Omega)^3$ and $f \in H^{-1/2}(\text{div}_{\partial\Omega}, \partial\Omega)$. Define $E_f = \gamma_t^{-1}f \in H(\text{curl}, \Omega)$, i.e., $\nu \times E_f|_{\partial\Omega} = f$ and E_f depends continuously and linearly on f . Moreover, we let $\psi \in H_0^1(\Omega)$ solve

$$\nabla \cdot (ik\epsilon\nabla\psi) = \nabla \cdot (J - ik\epsilon E_f),$$

which also depends continuously and linearly on J and E_f .

It follows from Lemma 2.3 that there exists a solution $(E_0, H) \in H(\text{curl}, \Omega) \times H(\text{curl}, \Omega)$ of the gauged system

$$\begin{cases} \nabla \times E_0 - ik\mu H = K - \nabla \times E_f & \text{in } \Omega, \\ \nabla \times H + ik\epsilon E_0 = J - ik\epsilon E_f - ik\epsilon\nabla\psi & \text{in } \Omega, \\ \nu \times E_0|_{\partial\Omega} = 0, \end{cases}$$

and E_0 and H depends linearly and continuously on $K - \nabla \times E_f \in L^2(\Omega)^3$ and $J - ik\epsilon E_f - ik\epsilon\nabla\psi \in L^2(\Omega)^3$. Hence, $E := E_0 + E_f + \nabla\psi$ and H solve (2.3)–(2.5) and depend linearly and continuously on $J, K \in L^2(\Omega)^3$ and $f \in H^{-1/2}(\text{div}_{\partial\Omega}, \partial\Omega)$.

The proof is complete. □

2.3. Unique continuation. The UCP is an important property to study the localized fields for differential equations. The UCP for the anisotropic Maxwell system was studied by [30, 36], which is of critical importance for our subsequent construction of the localized electromagnetic fields.

DEFINITION 2.1. *We say that (ϵ, μ) satisfies the UCP in Ω if any solution $(E, H) \in H(\text{curl}; \Omega) \times H(\text{curl}; \Omega)$ to the Maxwell system*

$$(2.9) \quad \begin{cases} \nabla \times E - ik\mu H = 0 & \text{in } \Omega, \\ \nabla \times H + ik\epsilon E = 0 & \text{in } \Omega \end{cases}$$

satisfies the property that if (E, H) vanishes in a nonempty open set D in Ω , then it must be identically vanishing in the whole domain Ω .

The UCP of the Maxwell system was proved by Leis [30] when the parameters ϵ, μ are C^2 scalar functions. When ϵ, μ are Lipschitz continuous anisotropic parameters, the UCP was proved by Nguyen and Wang [36]. In [32], Liu, Rondi, and Xiao have shown that the UCP holds for piecewise Lipschitz continuous matrix-valued functions ϵ and μ under some conditions, which we shall need for the subsequent study.

PROPOSITION 2.1 (UCP; Proposition 2.13 in [32]). *Given an open set $\widetilde{\Omega}$ in \mathbb{R}^3 , let $\epsilon, \mu \in L^\infty(\widetilde{\Omega}, \mathbb{R}^{3 \times 3})$ be matrix-valued functions in Ω satisfying (1.1) in $\widetilde{\Omega}$. Suppose the following:*

(1) There is a family $\{\Omega_i\}$ of pairwise disjoint domains with $\Omega_i \subset \tilde{\Omega}$ such that

$$\tilde{\Omega} \subset \cup_i \overline{\Omega_i}.$$

(2) The set $\Sigma_0 := \tilde{\Omega} \cap (\cup_i \partial\Omega_i)$ has Lebesgue measure zero (i.e., $|\Sigma_0| = 0$).

(3) The point $x \in \Sigma_0$ is called to be a partition point if there exists $\delta > 0$ such that

$$|B_\delta(x) \setminus (\Omega_i \cup \Omega_j)| = 0 \text{ for all } i \neq j$$

with $B_\delta(x) \cap \Omega_i$ and $B_\delta(x) \cap \Omega_j$ being nonempty sets. Consider the set

$$\mathcal{P}^c := \{x \in \Sigma_0 : x \text{ is not a partition point}\},$$

then we assume that $\tilde{\Omega} \setminus \mathcal{P}^c$ is connected.

(4) The functions $(\epsilon, \mu) = (\epsilon_i, \mu_i)$ in Ω_i , where (ϵ_i, μ_i) are locally Lipschitz matrix-valued functions in $\tilde{\Omega}$.

Then the UCP holds.

We have the following theorem.

THEOREM 2.2. Let Ω be a bounded Lipschitz domain and $\mathcal{F} \subset \Omega$ be a closed set in \mathbb{R}^3 such that $\Omega \setminus \mathcal{F}$ is connected to a relatively open boundary part $\Gamma \subseteq \partial\Omega$. Let $(E, H) \in H(\text{curl}; \Omega) \times H(\text{curl}; \Omega)$ solve

$$\begin{cases} \nabla \times E - ik\mu H = 0 & \text{in } \Omega \setminus \mathcal{F}, \\ \nabla \times H + ik\epsilon E = 0 & \text{in } \Omega \setminus \mathcal{F}. \end{cases}$$

If $\nu \times E|_\Gamma = \nu \times H|_\Gamma = 0$ on Γ , then $(E, H) = (0, 0)$ in $\Omega \setminus \mathcal{F}$.

Proof. Let \mathcal{O} be a nonempty open set in \mathbb{R}^3 such that $\mathcal{O} \cap \partial\Omega = \Gamma$ and $\mathcal{F} \subset \mathcal{O}$. In the open set $\tilde{\Omega} := \Omega \cup \mathcal{O}$, we define

$$\hat{\epsilon} := \begin{cases} \epsilon & \text{in } \Omega, \\ 1 & \text{in } \mathcal{O} \setminus \Omega, \end{cases} \text{ and } \hat{\mu} := \begin{cases} \mu & \text{in } \Omega, \\ 1 & \text{in } \mathcal{O} \setminus \Omega. \end{cases}$$

It can be seen that the parameters $\hat{\epsilon}$ and $\hat{\mu}$ satisfy the conditions (1)–(4) in Proposition 2.1 in the open set $\tilde{\Omega} = \Omega \cup \mathcal{O}$. Since $\nu \times E = \nu \times H = 0$ on Γ , we can extend (E, H) by $(0, 0)$ and define the extension functions

$$\hat{E} := \begin{cases} E & \text{in } \Omega, \\ 0 & \text{in } \mathcal{O} \setminus \Omega, \end{cases} \text{ and } \hat{H} := \begin{cases} H & \text{in } \Omega, \\ 0 & \text{in } \mathcal{O} \setminus \Omega. \end{cases}$$

First, we prove that $(\hat{E}, \hat{H}) \in H(\text{curl}; \Omega \cup \mathcal{O}) \times H(\text{curl}; \Omega \cup \mathcal{O})$. For any $\phi \in C_c^\infty(\Omega \cup \mathcal{O})$, we have

$$\begin{aligned} \int_{\Omega \cup \mathcal{O}} \hat{E} \cdot (\nabla \times \phi) dx &= \int_{\Omega} E \cdot (\nabla \times \phi) dx \\ &= \int_{\Omega} (\nabla \times E) \cdot \phi dx + \int_{\partial\Omega} E \cdot (\nu \times \phi) dS \\ &= \int_{\Omega \cup \mathcal{O}} ((\nabla \times E) \chi_\Omega) \cdot \phi dx, \end{aligned}$$

where we have used $E \cdot (\nu \times \phi) = -\phi \cdot (\nu \times E) = 0$ on Γ and $\phi = 0$ on $\partial\Omega \setminus \Gamma$. This shows that $\widehat{E} \in H(\text{curl}; \Omega \cup \mathcal{O})$ and that $\nabla \times \widehat{E}$ is the zero extension of $\nabla \times E$. The same holds for \widehat{H} , and thus it also follows that $(\widehat{E}, \widehat{H})$ is a solution of

$$\begin{cases} \nabla \times \widehat{E} - ik\widehat{\mu}\widehat{H} = 0 & \text{in } (\Omega \cup \mathcal{O}) \setminus \mathcal{F}, \\ \nabla \times \widehat{H} + ik\widehat{\epsilon}\widehat{E} = 0 & \text{in } (\Omega \cup \mathcal{O}) \setminus \mathcal{F}. \end{cases}$$

Notice that $\widehat{\epsilon}$ and $\widehat{\mu}$ are piecewise Lipschitz continuous functions fulfilling the ellipticity condition (1.1) and the conditions in Proposition 2.1. Recalling that $\widehat{E} = \widehat{H} = 0$ in $\mathcal{O} \setminus \overline{\Omega}$ (a nonempty open set), and by using Proposition 2.1, the UCP gives $\widehat{E} \equiv \widehat{H} \equiv 0$ in $(\Omega \cup \mathcal{O}) \setminus \mathcal{F}$.

The proof is complete. □

3. Localized electromagnetic fields. We are now in a position to present the main result on localizing and concentrating electromagnetic fields. We show that there exists boundary data (supported on an arbitrarily small boundary part) which can generate an electromagnetic field with an arbitrarily high energy on one part of the considered domain and an arbitrarily small energy on another part. This extends the related results in [12] for the conductivity equation and [20] for the Helmholtz equation to the more practical and challenging Maxwell system. In this section, we prove the existence of localized fields by using the functional analysis techniques from [12]. Recall our main result as follows.

THEOREM 3.1. *Let $\Omega \subset \mathbb{R}^3$ be a bounded Lipschitz domain and $\Gamma \subseteq \partial\Omega$ be a relatively open piece of the boundary. Let $\epsilon, \mu \in L^\infty(\Omega, \mathbb{R}^{3 \times 3})$ be real-valued, piecewise Lipschitz continuous functions satisfying (1.1) and $k \in \mathbb{R}_+$ be a nonresonant frequency. Let $D \Subset \Omega$ be a closed set with a connected complement $\Omega \setminus D$. For every open set $M \subseteq \Omega$ with $M \not\subseteq D$ (see Figure 1 for the schematic illustration), there exists a sequence $\{f^{(\ell)}\}_{\ell \in \mathbb{N}} \subset C_c^\infty(\Gamma)$ such that the electromagnetic fields fulfill*

$$(3.1) \quad \int_M (|E^{(\ell)}|^2 + |H^{(\ell)}|^2) \, dx \rightarrow \infty \quad \text{and} \quad \int_D (|E^{(\ell)}|^2 + |H^{(\ell)}|^2) \, dx \rightarrow 0 \quad \text{as } \ell \rightarrow \infty,$$

where, for $\ell \in \mathbb{N}$, $(E^{(\ell)}, H^{(\ell)}) \in H(\text{curl}, \Omega) \times H(\text{curl}, \Omega)$ is a solution of

$$(3.2) \quad \nabla \times E^{(\ell)} - ik\mu H^{(\ell)} = 0 \quad \text{in } \Omega,$$

$$(3.3) \quad \nabla \times H^{(\ell)} + ik\epsilon E^{(\ell)} = 0 \quad \text{in } \Omega$$

with the boundary data

$$(3.4) \quad \nu \times E^{(\ell)}|_{\partial\Omega} = \begin{cases} f^{(\ell)} & \text{on } \Gamma, \\ 0 & \text{otherwise.} \end{cases}$$

Proof of Theorem 3.1. We first note that it suffices to prove the theorem for an open subset of M . Hence, without loss of generality, we can assume that $M \Subset \Omega$ is open, $\overline{M} \cap D = \emptyset$, and $\Omega \setminus (\overline{M} \cup D)$ is connected. We follow the localized potentials strategy in [12, 20, 21] and first describe the energy terms in Theorem 3.1 as operator evaluations. Then we show that the ranges of the adjoints of these operators have trivial intersection. A functional analytic relation between the norm of an operator evaluation and the range of its adjoint then yields that the operator evaluations cannot be bounded by each other, which then shows that we can drive one energy term in Theorem 3.1 to infinity and the other one to zero.

For a measurable subset $O \subseteq \Omega$, we define

$$\mathcal{L}_O : H(\Gamma) \rightarrow L^2(O)^3 \times L^2(O)^3 \text{ by } f \mapsto (E|_O, H|_O),$$

where $H(\Gamma)$ is defined in (2.2), and $(E, H) \in L^2(\Omega)^3 \times L^2(\Omega)^3$ solves (3.2)–(3.3) with the boundary data $\nu \times E|_{\partial\Omega} = f$. Now we characterize the adjoint of this operator.

LEMMA 3.1. *The adjoint of \mathcal{L}_O is given by*

$$\mathcal{L}_O^* : L^2(O)^3 \times L^2(O)^3 \rightarrow H(\Gamma)^* \text{ by } (J, K) \rightarrow -\nu \times (\tilde{H} \times \nu)|_{\Gamma},$$

where $(\tilde{E}, \tilde{H}) \in H(\text{curl}, \Omega) \times H(\text{curl}, \Omega)$ solves the (adjoint) Maxwell system (cf. Theorem 2.1)

$$\begin{cases} \nabla \times \tilde{E} + ik\mu\tilde{H} = K\chi_O & \text{in } \Omega, \\ \nabla \times \tilde{H} - ik\epsilon\tilde{E} = J\chi_O & \text{in } \Omega, \\ \nu \times \tilde{E}|_{\partial\Omega} = 0, \end{cases}$$

and $K\chi_O$ and $J\chi_O$ denote the zero extensions of K and J to Ω , respectively.

Proof. Similar to section 2.1, we write the dual pairing on $H(\Gamma)^* \times H(\Gamma)$ as an integral for the sake of notational convenience. With this notation we have that

$$\begin{aligned} & \int_{\Gamma} \bar{f} \cdot \mathcal{L}_O^*(J, K) \, ds \\ &= \int_O (J, K) \cdot \overline{(E|_O, H|_O)} \, dx = \int_{\Omega} J\chi_O \cdot \bar{E} \, dx + \int_{\Omega} K\chi_O \cdot \bar{H} \, dx \\ &= \int_{\Omega} J\chi_O \cdot \bar{E} \, dx - \frac{1}{ik} \int_{\Omega} K\chi_O \cdot (\mu^{-1} \nabla \times \bar{E}) \, dx \\ &= \int_{\Omega} (\nabla \times \tilde{H} - ik\epsilon\tilde{E}) \cdot \bar{E} \, dx - \frac{1}{ik} \int_{\Omega} (\nabla \times \tilde{E} + ik\mu\tilde{H}) \cdot (\mu^{-1} \nabla \times \bar{E}) \, dx \\ &= \int_{\Omega} (\nabla \times \tilde{H}) \cdot \bar{E} \, dx - \int_{\Omega} \tilde{H} \cdot (\nabla \times \bar{E}) \, dx \\ &\quad - \frac{1}{ik} \left(\int_{\Omega} (\mu^{-1} \nabla \times \bar{E}) \cdot (\nabla \times \tilde{E}) \, dx - \int_{\Omega} k^2 \epsilon \bar{E} \cdot \tilde{E} \, dx \right) \\ &= - \int_{\partial\Omega} (\nu \times \bar{E}|_{\partial\Omega}) \cdot (\nu \times (\tilde{H}|_{\partial\Omega} \times \nu)) \, dS = - \int_{\Gamma} \bar{f} \cdot (\nu \times (\tilde{H}|_{\Gamma} \times \nu)) \, dS, \end{aligned}$$

where we make use of the fact that ϵ and μ are real-valued and symmetric. We also utilized the integration by parts formula (2.1) and that $\tilde{E} \in H_0(\text{curl}, \Omega)$ implies that

$$\int_{\Omega} (\mu^{-1} \nabla \times \bar{E}) \cdot (\nabla \times \tilde{E}) \, dx - \int_{\Omega} k^2 \epsilon \bar{E} \cdot \tilde{E} \, dx = 0$$

by Theorem 2.1(a). □

Next we show the following property for the ranges of the adjoint operators \mathcal{L}_M^* and \mathcal{L}_D^* .

LEMMA 3.2. *\mathcal{L}_M and \mathcal{L}_D^* are injective, the ranges $\mathcal{R}(\mathcal{L}_M^*)$ and $\mathcal{R}(\mathcal{L}_D^*)$ are both dense in $H(\Gamma)^*$, and*

$$(3.5) \quad \mathcal{R}(\mathcal{L}_M^*) \cap \mathcal{R}(\mathcal{L}_D^*) = \{0\}.$$

Proof. The proof follows from the UCP for the Maxwell system. By Proposition 2.1, one readily sees that \mathcal{L}_M and \mathcal{L}_D are injective, and therefore $\mathcal{R}(\mathcal{L}_M^*)$ and $\mathcal{R}(\mathcal{L}_D^*)$ both are dense in $H(\Gamma)^*$.

To prove (3.5), let $g \in \mathcal{R}(\mathcal{L}_M^*) \cap \mathcal{R}(\mathcal{L}_D^*)$, and then there exist $J_M, K_M \in L^2(M)$ and $J_D, K_D \in L^2(D)$ such that the solutions $(E_M, H_M), (E_D, H_D) \in H(\text{curl}, \Omega) \times H(\text{curl}, \Omega)$ of

$$\begin{cases} \nabla \times E_M + ik\mu H_M = K_M \chi_M & \text{in } \Omega, \\ \nabla \times H_M - ik\epsilon E_M = J_M \chi_M & \text{in } \Omega, \\ \nu \times E_M|_{\partial\Omega} = 0, & \text{on } \partial\Omega, \end{cases} \quad \text{and} \quad \begin{cases} \nabla \times E_D + ik\mu H_D = K_D \chi_D & \text{in } \Omega, \\ \nabla \times H_D - ik\epsilon E_D = J_D \chi_D & \text{in } \Omega, \\ \nu \times E_D|_{\partial\Omega} = 0 & \text{on } \partial\Omega \end{cases}$$

fulfill

$$\nu \times (H_M \times \nu)|_{\Gamma} = g = \nu \times (H_D \times \nu)|_{\Gamma}.$$

Since $\Omega \setminus (\overline{M} \cup D)$ is connected, we obtain by using Theorem 2.2 that

$$E_M = E_D \text{ in } \Omega \setminus (\overline{M} \cup D).$$

Hence, we can define

$$\mathbb{E} := \begin{cases} E_D & \text{in } M, \\ E_M & \text{in } D, \\ E_D = E_M & \text{in } \Omega \setminus (\overline{M} \cup D), \end{cases} \quad \text{and} \quad \mathbb{H} := \begin{cases} H_D & \text{in } M, \\ H_M & \text{in } D, \\ H_D = H_M & \text{in } \Omega \setminus (\overline{M} \cup D). \end{cases}$$

As in the proof of UCP, Theorem 2.2, it is easy to see that $(\mathbb{E}, \mathbb{H}) \in H(\text{curl}, \Omega) \times H(\text{curl}, \Omega)$ is a solution of

$$\begin{cases} \nabla \times \mathbb{E} + ik\mu \mathbb{H} = 0 & \text{in } \Omega, \\ \nabla \times \mathbb{H} - ik\epsilon \mathbb{E} = 0 & \text{in } \Omega, \\ \nu \times \mathbb{E}|_{\partial\Omega} = 0 & \text{on } \partial\Omega. \end{cases} \quad \square$$

Since k is nonresonant, it follows that $(\mathbb{E}, \mathbb{H}) = (0, 0)$, and hence $g = 0$. This completes the proof.

Now we can use the following tool from functional analysis.

LEMMA 3.3. *Let $X, Y_1,$ and Y_2 be Hilbert spaces, and $\mathcal{A}_1 : X \rightarrow Y_1$ and $\mathcal{A}_2 : X \rightarrow Y_2$ be linear bounded operators. Then*

$$\exists C > 0 : \|\mathcal{A}_1 x\| \leq C \|\mathcal{A}_2 x\| \quad \text{for all } x \in X \quad \text{if and only if} \quad \mathcal{R}(\mathcal{A}_1^*) \subseteq \mathcal{R}(\mathcal{A}_2^*).$$

Proof. This is proved for reflexive Banach spaces in [12, Lemma 2.5]. □

Proof of Theorem 1.1. From Lemma 3.2 it follows that $\mathcal{R}(\mathcal{L}_M^*) \not\subseteq \mathcal{R}(\mathcal{L}_D^*)$. Using Lemma 3.3 this shows that

$$\exists C > 0 : \|\mathcal{L}_M f\| \leq C \|\mathcal{L}_D f\| \quad \text{for all } f \in H(\Gamma),$$

and by continuity of \mathcal{L}_M and \mathcal{L}_D and density of $C_c^\infty(\Gamma) \subset H(\Gamma)$ this is equivalent to

$$(3.6) \quad \exists C > 0 : \|\mathcal{L}_M f\| \leq C \|\mathcal{L}_D f\| \quad \text{for all } f \in C_c^\infty(\Gamma).$$

Using (3.6) with $C := \ell^2$ for all $\ell \in \mathbb{N}$ we thus obtain a sequence $\{\tilde{f}^{(\ell)}\}_{\ell \in \mathbb{N}} \subset C_c^\infty(\Gamma)$ with

$$\|\mathcal{L}_M \tilde{f}^{(\ell)}\| > \ell^2 \|\mathcal{L}_D \tilde{f}^{(\ell)}\| \quad \text{for all } \ell \in \mathbb{N}.$$

By injectivity $\mathcal{L}_M \tilde{f}^{(\ell)} \neq 0$ implies $\tilde{f}^{(\ell)} \neq 0$ and $\mathcal{L}_D \tilde{f}^{(\ell)} \neq 0$, so that we can define

$$f^{(\ell)} := \frac{\tilde{f}^{(\ell)}}{\ell \|\mathcal{L}_D \tilde{f}^{(\ell)}\|} \in C_c^\infty(\Gamma)$$

and it follows that

$$\begin{aligned} \int_M \left(|E^{(\ell)}|^2 + |H^{(\ell)}|^2 \right) dx &= \|\mathcal{L}_M f^{(\ell)}\|^2 > \ell^2 \rightarrow \infty, \\ \int_D \left(|E^{(\ell)}|^2 + |H^{(\ell)}|^2 \right) dx &= \|\mathcal{L}_D f^{(\ell)}\|^2 = \frac{1}{\ell^2} \rightarrow 0. \end{aligned}$$

This completes the proof of Theorem 1.1. \square

Remark 3.1.

- (a) A constructive version of the existence proof for the localized fields can be obtained as in [12, Lemma 2.8].
- (b) With the same arguments as in [20, section 4.1] one can also show that for all spaces $W \subseteq H(\Gamma)$ with finite codimensions, one can find a sequence $\{f^{(\ell)}\}_{\ell \in \mathbb{N}} \subset W$ such that the corresponding electromagnetic fields fulfill (3.1). This might be useful for developing monotonicity-based reconstruction methods as in [20].

4. Runge approximation property for the partial data Maxwell system. In this section we derive an extension of the localization result in Theorem 1.1 and establish a *Runge approximation property* for the partial data Maxwell system, which is of mathematical interest for its own sake. We show that every solution of the Maxwell system on a subset of Ω with a Lipschitz boundary and a connected complement can be approximated arbitrarily well by a sequence of solutions on the whole domain Ω with partial boundary data. Since we can choose a solution that is zero on a part of Ω and nonzero on another part of Ω , this readily implies a fortiori the localization result in Theorem 1.1. We also refer to [20] for the connection between Runge approximation properties and localized solutions.

THEOREM 4.1. *Let $\Omega \subset \mathbb{R}^3$ be a bounded Lipschitz domain and $\Gamma \subseteq \partial\Omega$ be a relatively open piece of the boundary. Let $\epsilon, \mu \in L^\infty(\Omega, \mathbb{R}^{3 \times 3})$ be real-valued, piecewise Lipschitz continuous functions satisfying (1.1) and $k \in \mathbb{R}_+$ be a nonresonant frequency.*

Let $O \Subset \Omega$ be an open set with Lipschitz boundary and connected complement $\Omega \setminus \bar{O}$. For every solution $(e, h) \in H(\text{curl}, O) \times H(\text{curl}, O)$ of

$$(4.1) \quad \nabla \times e - ik\mu h = 0 \quad \text{in } O,$$

$$(4.2) \quad \nabla \times h + ik\epsilon e = 0 \quad \text{in } O,$$

there exists a sequence $\{f^{(\ell)}\}_{\ell \in \mathbb{N}} \subset C_c^\infty(\Gamma)$ such that the electromagnetic fields fulfill

$$\|E^{(\ell)} - e\|_{L^2(O)} \rightarrow 0 \quad \text{and} \quad \|H^{(\ell)} - h\|_{L^2(O)} \rightarrow 0 \quad \text{as } \ell \rightarrow \infty,$$

where $(E^{(\ell)}, H^{(\ell)}) \in H(\text{curl}, O) \times H(\text{curl}, O)$ solve (3.2)–(3.4) in O .

Proof. Let $(e, h) \in H(\text{curl}, O) \times H(\text{curl}, O)$ solve (4.1)–(4.2). With the operator \mathcal{L}_O introduced in section 3, we shall show that

$$(4.3) \quad (e, h) \in \overline{\mathcal{R}(\mathcal{L}_O)} = (\mathcal{R}(\mathcal{L}_O)^\perp)^\perp = \mathcal{N}(\mathcal{L}_O^*)^\perp,$$

where the closure and the orthogonal complement are understood with respect to the $L^2(O)^3 \times L^2(O)^3$ -scalar product. This shows that the assertion holds with a sequence $\{f^{(\ell)}\}_{\ell \in \mathbb{N}} \subset H(\Gamma)$, and it follows by density that the assertion holds with a sequence $\{f^{(\ell)}\}_{\ell \in \mathbb{N}} \subset C_c^\infty(\Gamma)$.

To prove (4.3), we let $(J, K) \in \mathcal{N}(\mathcal{L}_O^*) \subseteq L^2(O)^3 \times L^2(O)^3$. Then, by Lemma 3.1, there exists $(E, H) \in H(\text{curl}, O) \times H(\text{curl}, O)$ that solves

$$\begin{cases} \nabla \times E + ik\mu H = K\chi_O & \text{in } \Omega, \\ \nabla \times H - ik\epsilon E = J\chi_O & \text{in } \Omega, \\ \nu \times E|_{\partial\Omega} = 0 & \text{on } \partial\Omega \end{cases}$$

with $\nu \times (H \times \nu)|_\Gamma = \mathcal{L}_O^*(J, K) = 0$. The UCP in Theorem 2.2 implies that $(E, H) = (0, 0)$ on $\Omega \setminus \bar{O}$ and thus

$$\nu \times E|_{\partial O} = 0 = \nu \times H|_{\partial O}.$$

Hence, using the integration by parts formula (2.1), it follows that

$$\begin{aligned} & \int_O (e \cdot \bar{J} + h \cdot \bar{K}) \, dx \\ &= \int_O e \cdot (\nabla \times \bar{H} + ik\epsilon \bar{E}) \, dx + \int_O h \cdot (\nabla \times \bar{E} - ik\mu \bar{H}) \, dx \\ &= \int_O ((\nabla \times e) \cdot \bar{H} + ik\epsilon e \cdot \bar{E}) \, dx + \int_O ((\nabla \times h) \cdot \bar{E} - ik\mu h \cdot \bar{H}) \, dx \\ &= \int_O (\nabla \times e - ik\mu h) \cdot \bar{H} \, dx + \int_O (\nabla \times h + ik\epsilon e) \cdot \bar{E} \, dx = 0. \end{aligned}$$

This shows $(e, h) \perp (J, K)$ so that (4.3) holds, and thus the assertion is proved. \square

Remark 4.1. The Runge approximation property in Theorem 4.1 implies the localization property in Theorem 1.1 by the following argument. Let $D \Subset \Omega$ be a closed set with a connected complement $\Omega \setminus D$, and $M \subseteq \Omega$ be an open set with $M \not\subseteq D$ (see again Figure 1). By shrinking M and enlarging D , we can assume that M is an open set and D is a closed set with Lipschitz boundaries, $\bar{M} \cup D \Subset \Omega$, and $\bar{M} \cap D = \emptyset$ and that $\Omega \setminus (\bar{M} \cup D)$ is connected.

The UCP in Theorem 2.2 implies that a solution of the Maxwell system in Ω with nontrivial boundary data cannot vanish identically on M . This shows that there exists a nonzero solution of the Maxwell system on M . We extend this solution by zero on D and obtain a solution $(e, h) \in H(\text{curl}, O) \times H(\text{curl}, O)$ on $O := M \cup \text{int}D$ with $(e, h)|_{\text{int}D} \equiv (0, 0)$ and $(e, h)|_M \neq (0, 0)$. Then the Runge approximation sequence from Theorem 4.1 converges to zero on D but not on M and a simple scaling argument as in the proof of Theorem 1.1 in section 3 gives a sequence of electromagnetic fields such that

$$\int_M (|E^{(\ell)}|^2 + |H^{(\ell)}|^2) \, dx \rightarrow \infty \quad \text{and} \quad \int_D (|E^{(\ell)}|^2 + |H^{(\ell)}|^2) \, dx \rightarrow 0 \quad \text{as } \ell \rightarrow \infty,$$

and $\nu \times E^{(\ell)} \in C_c^\infty(\Gamma)$ for all $\ell \in \mathbb{N}$, which also proves Theorem 1.1.

5. Concluding remarks. We considered field localizing and concentrating for the electromagnetic waves governed by the time-harmonic Maxwell system in a bounded domain occupied by a given medium that is generic and could be anisotropic. It has been shown that through proper boundary inputs, one can generate electromagnetic fields with the corresponding energy concentrated in a given subregion while nearly vanishing in another given subregion. We would like to emphasize that the localizing results are known previously for several scalar models including the conductivity equation and the Helmholtz equation, but the extension to the Maxwell system with nonsmooth anisotropic coefficients requires considerable care and technical involvement. As pointed out in Remark 3.1, the result can be used to develop monotonicity-based reconstruction methods for inverse problems associated with the electromagnetic waves. In fact, with the localizing result established in the present article, this can be done by following a spirit similar to the one in [20] for the Helmholtz equation by one of the authors of this article. We also propose more applications of practical interest, including telecommunications and inductive charging. This new perspective poses interesting problems for future investigation. For example, for certain given boundary inputs, one may consider to construct a specific medium configuration for localizing and concentrating the electromagnetic fields in a desirable way.

Acknowledgment. The authors would like to thank the anonymous referees for many constructive comments and suggestions, which have led to significant improvement on the result and presentation of the paper.

REFERENCES

- [1] G. ALESSANDRINI, *Singular solutions of elliptic equations and the determination of conductivity by boundary measurements*, J. Differential Equations, 84 (1990), pp. 252–272.
- [2] H. AMMARI, G. CIRAULO, H. KANG, H. LEE, AND G. W. MILTON, *Spectral theory of a Neumann–Poincaré-type operator and analysis of cloaking due to anomalous localized resonance*, Arch. Ration. Mech. Anal., 208 (2013), pp. 667–692.
- [3] K. ANDO, H. KANG, AND H. LIU, *Plasmon resonance with finite frequencies: A validation of the quasi-static approximation for diametrically small inclusions*, SIAM J. Appl. Math., 76 (2016), pp. 731–749.
- [4] L. ARNOLD AND B. HARRACH, *Unique shape detection in transient eddy current problems*, Inverse Problems, 29 (2013), 095004.
- [5] A. BARTH, B. HARRACH, N. HYVÖNEN, AND L. MUSTONEN, *Detecting stochastic inclusions in electrical impedance tomography*, Inverse Problems, 33 (2017), 115012.
- [6] G. BOUCHITTÉ AND B. SCHWEIZER, *Cloaking of small objects by anomalous localized resonance*, Quart. J. Mech. Appl. Math., 63 (2010), pp. 437–463.
- [7] T. BRANDER, B. HARRACH, M. KAR, AND M. SALO, *Monotonicity and Enclosure Methods for the p -Laplace Equation*, SIAM J. Appl. Math., 78 (2018), pp. 742–758.
- [8] A. BUFFA, M. COSTABEL, AND D. SHEEN, *On traces for $H(\text{curl}, \Omega)$ in Lipschitz domains*, J. Math. Anal. Appl., 276 (2002), pp. 845–867.
- [9] H. GARDE, *Comparison of linear and non-linear monotonicity-based shape reconstruction using exact matrix characterizations*, Inverse Probl. Sci. Eng., 26 (2018), pp. 33–50.
- [10] H. GARDE AND S. STABOULIS, *Convergence and regularization for monotonicity-based shape reconstruction in electrical impedance tomography*, Numer. Math., 135 (2017), pp. 1221–1251.
- [11] H. GARDE AND S. STABOULIS, *The Regularized Monotonicity Method: Detecting Irregular Indefinite Inclusions*, <https://arxiv.org/abs/1705.07372>, 2017.
- [12] B. GEBAUER, *Localized potentials in electrical impedance tomography*, Inverse Probl. Imaging, 2 (2008), pp. 251–269.
- [13] R. GRIESMAIER AND B. HARRACH, *Monotonicity in Inverse Medium Scattering on Unbounded Domains*, SIAM J. Appl. Math., to appear.
- [14] B. HARRACH, *On uniqueness in diffuse optical tomography*, Inverse Problems, 25 (2009), 055010.

- [15] B. HARRACH, *Simultaneous determination of the diffusion and absorption coefficient from boundary data*, Inverse Probl. Imaging, 6 (2012), pp. 663–679.
- [16] B. HARRACH, E. LEE, AND M. ULLRICH, *Combining frequency-difference and ultrasound modulated electrical impedance tomography*, Inverse Problems, 31 (2015), 095003.
- [17] B. HARRACH AND Y.-H. LIN, *Monotonicity-based Inversion of the Fractional Schrödinger Equation*, <https://arxiv.org/abs/1711.05641>, 2017.
- [18] B. HARRACH AND M. N. MINH, *Enhancing residual-based techniques with shape reconstruction features in electrical impedance tomography*, Inverse Problems, 32 (2016), 125002.
- [19] B. HARRACH AND M. N. MINH, *Monotonicity-based regularization for phantom experiment data in electrical impedance tomography*, in New Trends in Parameter Identification for Mathematical Models, B. Hofmann, A. Leitao, and J. P. Zubelli, eds., Trends Math., Springer, New York, 2018, pp. 107–120.
- [20] B. HARRACH, V. POHJOLA, AND M. SALO, *Monotonicity and Local Uniqueness for the Helmholtz Equation*, <https://arxiv.org/abs/1709.08756>, 2017.
- [21] B. HARRACH AND J. K. SEO, *Exact shape-reconstruction by one-step linearization in electrical impedance tomography*, SIAM J. Math. Anal., 42 (2010), pp. 1505–1518.
- [22] B. HARRACH AND M. ULLRICH, *Monotonicity-based shape reconstruction in electrical impedance tomography*, SIAM J. Math. Anal., 45 (2013), pp. 3382–3403.
- [23] B. HARRACH AND M. ULLRICH, *Resolution guarantees in electrical impedance tomography*, IEEE Trans. Med. Imaging, 34 (2015), pp. 1513–1521.
- [24] B. HARRACH AND M. ULLRICH, *Local uniqueness for an inverse boundary value problem with partial data*, Proc. Amer. Math. Soc., 145 (2017), pp. 1087–1095.
- [25] V. ISAKOV, *On uniqueness of recovery of a discontinuous conductivity coefficient*, Comm. Pure Appl. Math., 41 (1988), pp. 865–877.
- [26] A. KIRSCH AND F. HETTLICH, *Mathematical Theory of Time-harmonic Maxwell's Equations*, Springer, New York, 2016.
- [27] R. KOHN AND M. VOGELIUS, *Determining conductivity by boundary measurements*, Comm. Pure Appl. Math., 37 (1984), pp. 289–298.
- [28] R. V. KOHN AND M. VOGELIUS, *Determining conductivity by boundary measurements II. Interior results*, Comm. Pure Appl. Math., 38 (1985), pp. 643–667.
- [29] P. D. LAX, *A stability theorem for solutions of abstract differential equations, and its application to the study of the local behavior of solutions of elliptic equations*, Comm. Pure Appl. Math., 9 (1956), pp. 747–766.
- [30] R. LEIS, *Initial Boundary Value Problems in Mathematical Physics*, Teubner, Stuttgart, 1986.
- [31] H. LI, J. LI, AND H. LIU, *On quasi-static cloaking due to anomalous localized resonance in \mathbb{R}^3* , SIAM J. Appl. Math., 75 (2015), pp. 1245–1260.
- [32] H. LIU, L. RONDI, AND J. XIAO, *Mosco Convergence for $H(\text{curl})$ Spaces, Higher Integrability for Maxwell's Equations, and Stability in Direct and Inverse EM Scattering Problems*, <https://arxiv.org/abs/1603.07555>, 2016.
- [33] A. MAFFUCCI, A. VENTO, S. VENTRE, AND A. TAMBURRINO, *A novel technique for evaluating the effective permittivity of inhomogeneous interconnects based on the monotonicity property*, IEEE Trans. Compon. Packag. Manuf. Tech., 6 (2016), pp. 1417–1427.
- [34] G. W. MILTON AND N.-A. P. NICOROVICI, *On the cloaking effects associated with anomalous localized resonance*, in Proceedings of the Royal Society of London A: Mathematical, Physical and Engineering Sciences, Vol. 462, The Royal Society, London, 2006, pp. 3027–3059.
- [35] P. MONK, *Finite Element Methods for Maxwell's Equations*, Numer. Math. Sci. Comput., Oxford University Press, Oxford, 2003, <https://doi.org/10.1093/acprof:oso/9780198508885.001.0001>.
- [36] T. NGUYEN AND J.-N. WANG, *Quantitative uniqueness estimate for the Maxwell system with Lipschitz anisotropic media*, Proc. Amer. Math. Soc., 140 (2012), pp. 595–605.
- [37] N.-A. P. NICOROVICI, G. W. MILTON, R. C. MCPHEDRAN, AND L. C. BOTTEN, *Quasistatic cloaking of two-dimensional polarizable discrete systems by anomalous resonance*, Opt. Express, 15 (2007), pp. 6314–6323.
- [38] Z. SU, L. UDPA, G. GIOVINCO, S. VENTRE, AND A. TAMBURRINO, *Monotonicity principle in pulsed eddy current testing and its application to defect sizing*, in Proceedings of the International Applied Computational Electromagnetics Society Symposium (ACES), 2017, IEEE, 2017, pp. 1–2.
- [39] A. TAMBURRINO AND G. RUBINACCI, *A new non-iterative inversion method for electrical resistance tomography*, Inverse Problems, 18 (2002), p. 1809.
- [40] A. TAMBURRINO, Z. SUA, S. VENTRE, L. UDPA, AND S. S. UDPA, *Monotonicity based imaging method in time domain eddy current testing*, in Electromagnetic Nondestructive Evaluation (XIX), Stud. Appl. Electromagn. Mech. 41, 2016, pp. 1–8.

- [41] L. TARTAR, *On the characterization of traces of a Sobolev space used for Maxwell's equation*, in Proceedings of a Meeting held in Bordeaux, 1997.
- [42] S. VENTRE, A. MAFFUCCI, F. CAIRE, N. LE LOSTEC, A. PERROTTA, G. RUBINACCI, B. SARTRE, A. VENTO, AND A. TAMBURRINO, *Design of a real-time eddy current tomography system*, IEEE Trans. Magnetics, 53 (2017), pp. 1–8.
- [43] WIKIPEDIA, *Inductive Charging*, https://en.wikipedia.org/wiki/Inductive_charging.
- [44] WIKIPEDIA, *Telecommunication*, <https://en.wikipedia.org/wiki/Telecommunication>.
- [45] L. ZHOU, B. HARRACH, AND J. K. SEO, *Monotonicity-based electrical impedance tomography for lung imaging*, Inverse Problems, 34 (2018), 045005.