

## Spatial point pattern analysis by using Voronoi diagrams and Delaunay tessellations - A comparative study

CHIU, Sung Nok

*Published in:*  
Biometrical Journal

*DOI:*  
[10.1002/bimj.200390018](https://doi.org/10.1002/bimj.200390018)

Published: 01/01/2003

[Link to publication](#)

*Citation for published version (APA):*

CHIU, S. N. (2003). Spatial point pattern analysis by using Voronoi diagrams and Delaunay tessellations - A comparative study. *Biometrical Journal*, 45(3), 367-376. <https://doi.org/10.1002/bimj.200390018>

### General rights

Copyright and intellectual property rights for the publications made accessible in HKBU Scholars are retained by the authors and/or other copyright owners. In addition to the restrictions prescribed by the Copyright Ordinance of Hong Kong, all users and readers must also observe the following terms of use:

- Users may download and print one copy of any publication from HKBU Scholars for the purpose of private study or research
- Users cannot further distribute the material or use it for any profit-making activity or commercial gain
- To share publications in HKBU Scholars with others, users are welcome to freely distribute the permanent publication URLs

# Spatial Point Pattern Analysis by using Voronoi Diagrams and Delaunay Tessellations — A Comparative Study

S. N. Chiu

*Hong Kong Baptist University*

SUMMARY. Given a spatial point pattern, we use various characteristics of its Voronoi diagram and Delaunay tessellation to extract information of the dependence between points. In particular, we use the characteristics to construct statistics for testing complete spatial randomness. It is shown that the minimum angle of a typical Delaunay triangle is sensitive to both regularity and clustering alternatives, whilst the triangle's area or perimeter is more sensitive to clustering than regularity. These statistics are also sensitive to the Baddeley-Silverman cell process.

KEY WORDS: Complete spatial randomness, Delaunay tessellation, goodness of fit, spatial point pattern, Voronoi diagram.

RUNNING TITLE: Spatial Point Pattern Analysis

## 1. Introduction

An important task of the analysis of spatial point patterns is to examine the dependence between points. Many summary functions, for example, the nearest neighbour distance distribution, Ripley's  $K$ -function, the empty space function and the periodogram, have been proposed to extract such information. One way to assess how good a summary function can capture the dependence structure of a given point pattern is to see how sensitive it will be when it is used to test the complete spatial randomness (CSR) hypothesis against various alternatives.

However, the distribution theory for most summary functions has not yet been well developed and so simulation-based inference (Møller and Waagepetersen, 2002) has usually been adopted. Comparative studies can be found in the literature (Diggle, 1979; Grabarnik and Chiu, 2002; Mugglestone and Renshaw, 2001; Ripley, 1979; Thönnies and van Lieshout, 1999), and the conclusion from these studies is that there is no uniformly best test.

This paper discusses test statistics constructed from the Voronoi diagram and the Delaunay tessellation generated by the observed point pattern. Since the Voronoi diagram and the Delaunay tessellation are uniquely determined by the generating points and vice versa, they can be used to construct summary functions to test the CSR hypothesis.

The idea of course is not new. Geographers have used Voronoi diagrams and Delaunay tessellations to study point patterns for a long time, e.g. Brown (1965) and Evans (1967), but the idea was not widely developed until later (see Okabe *et al.*, 2000, Chapter 8, for a review). A typical difficulty in this approach is that the distributions of most characteristics of the Voronoi diagram and the Delaunay tessellation generated by a stationary Poisson

process are unknown and can only be approximated by simulation. Using approximated distributions may lead to lower power tests.

Since the distributions of the interior angles and edge lengths of Poisson Delaunay triangles are known, Vincent *et al.* (1977) assessed goodness of fit of an empirical pattern to a Poisson process by using Pearson's  $\chi^2$  statistics for the interior angles and the edge lengths of Delaunay triangles, as well as the numbers of edges of Voronoi polygons; the distribution of the latter was approximated by ten realisations with one thousand Poisson points; Boots (1975, 1986) considered the Kolmogorov-Smirnov statistic, the Cramér-von Mises statistic, the Watson statistic and the Anderson-Darling statistic for the interior angles, the minimum angles and the maximum angles of Delaunay triangles. However, the correlation between Delaunay triangles and between Voronoi polygons in a realisation had not been taken into consideration in these studies.

Since numerically tractable expressions for the distributions of some characteristics were recently derived (see Okabe *et al.*, 2000, Sections 5.5.4 and 5.11), this short paper on one hand re-does the goodness-of-fit analyses by Boots (1975, 1986) and Vincent *et al.* (1977) in a more appropriate way and on the other hand compares the powers of tests based on more different characteristics. The Kolmogorov-Smirnov statistic  $\sup_x |\sqrt{n}\{F(x) - F_n(x)\}|$  and the Cramér-von Mises statistic  $\int_{-\infty}^{\infty} n\{F(x) - F_n(x)\}^2 dF(x)$ , where  $F$  and  $F_n$  respectively denote the distribution function under the null hypothesis and the empirical distribution function of a given sample of size  $n$ , will be used but their asymptotic critical values will not because we have neither independent nor a large number of observations; Monte Carlo tests will be used instead.

Since the statistics based on the characteristics of Voronoi diagrams and Delaunay tessellations are space-domain techniques, we compared the powers of these statistics with the statistic based on the most popular space-domain summary function, namely Ripley's  $K$ -function. We restrict ourselves to stationary alternatives. For a comparison between various space-domain tests and spectral tests against regular, clustered and non-stationary alternatives, see Mugglestone and Renshaw (2001).

## 2. Voronoi diagram and Delaunay tessellation

Given a realisation  $\{x_i\}$  of a point process on the plane, we associate all locations on the plane with the closest member(s) of the realisation, i.e.  $x_i$  is associated with the polygon  $V_i = \{x : \|x - x_i\| \leq \|x - x_j\| \text{ for } j \neq i\}$ . The collection  $\{V_i\}$  forms the Voronoi diagram of the given realisation and the points of the process are called generators. The polygon  $V_i$  is called the Voronoi polygon of  $x_i$ . The Delaunay tessellation of  $\{x_i\}$  is obtained by joining all pairs of generators whose Voronoi polygons share a common edge. If the point process is simple and stationary, then almost surely the Delaunay tessellation is a collection of triangles, which are called Delaunay triangles.

The following nine characteristics will be examined:

- the length  $L_V$  of an edge in a Voronoi diagram,
- the distance  $R$  between a generating point and a vertex of the Voronoi polygon of the generating point (the radius of the circle circumscribing a Delaunay triangle)
- the area  $A_D$  of a Delaunay triangle,
- the perimeter  $P_D$  of a Delaunay triangle,

- the length  $L_D$  of an edge in a Delaunay tessellation,
- an interior angle  $\alpha$  of a Delaunay triangle,
- the minimum angle  $\alpha_{\min}$  of a Delaunay triangle,
- the middle angle  $\alpha_{\text{mid}}$  of a Delaunay triangle,
- the maximum angle  $\alpha_{\max}$  of a Delaunay triangle.

Numerically tractable expressions for their distributions under the CSR hypothesis are known (Okabe *et al.*, 2000, Sections 5.5.4 and 5.11).

Other characteristics and their combinations, such as the roundness factor  $4\pi A_V/P_V^2$  (Marcelpoil and Usson, 1992), where  $A_V$  and  $P_V$  are the area and the perimeter of a Voronoi polygon, may be equally or even more capable of detecting deviations from CSR; however, their distribution functions can only be approximated by simulation. The powers of such combinations in testing CSR will be left for future endeavours. This paper focuses on characteristics with known distributions.

### 3. Simulation

In reality a point pattern is observed via a bounded observation window. Thus, when we simulate Voronoi polygons, no matter how many points there are, allowance must be made for edge effects introduced by the polygons close to the boundary of the observation window. One practical approach is to exclude from consideration any polygons for which a circle, centred at any vertex of the polygon and passing through the three points in patterns which are equidistant from the vertex, intersects the boundary of the observation window. If the

window is a rectangle, another approach is to convert the window into a torus, which is usually referred to as a periodic boundary condition. The latter approach is adopted in this paper.

The alternative models used here generate regular or clustered point patterns with 100 points in a unit square with periodic boundary condition. The simple sequential inhibition process, which puts sequentially 100 non-overlapping discs of radius  $r$  uniformly and independently on the torus and then places points at the centres of these discs, was used to generate regular patterns. To get clustered patterns, we simulate a version of the Matérn cluster process by putting a cluster centre uniformly distributed in the torus and then placing a Poisson number, with mean  $\rho$ , of points one by one uniformly and independently within the disc of radius  $r$  centred at the cluster centre; this procedure is repeated until 100 points are generated.

For each simulated pattern, 999 realisations of a Poisson process with the same number of points were generated. If under the CSR hypothesis the value of the Kolmogorov-Smirnov statistic or the Cramér-von Mises statistic for the distribution of a characteristic from the pattern, when pooled together with the values from the simulated realisations of the Poisson process, is one of the largest fifty, then the CSR hypothesis is rejected. Tables 1 and 2, respectively, give estimated powers against regular alternatives with hard core radius  $r = 0.05, 0.2$  and  $0.3$ , and clustered alternatives with cluster radius  $r = 0.1$  and  $0.2$  and mean cluster size  $\rho = 5$  and  $10$ . The powers were estimated from 100 realisations of the alternative models.

Table 1 about here
--------------------

Table 2 about here

We can see from the estimated powers that neither the Kolmogorov–Smirnov statistics nor the Cramér–von Mises statistics are always superior to the other. Both  $\alpha_{\max}$  and  $\alpha_{\min}$  perform better than  $\alpha$ , and  $\alpha_{\min}$  has a higher power than  $\alpha_{\max}$ . This agrees with the conclusion in Boots (1986), even though asymptotic critical values were inappropriately used there. For alternatives that exhibit moderate regularity,  $\alpha_{\min}$  yields the most powerful statistic among all these characteristics;  $A_D$  the second, and the rest performed poorly. However, for alternatives that exhibit moderate clustering, say cluster radius  $r = 0.2$  and mean cluster size  $\rho = 5$ ,  $\alpha_{\min}$ , although still better than  $\alpha_{\max}$ ,  $\alpha_{\text{mid}}$  and  $\alpha$ , performs worse than  $L_V$ ,  $R$ ,  $A_D$ ,  $P_D$  and  $L_D$ . Thus, we may conclude that these five characteristics are more able to detect clustering, while  $\alpha_{\min}$  is more able to detect regularity.

Consider the probability density functions  $f$  and  $g$ , respectively, of  $L_D$  and  $\alpha_{\min}$  in the Delaunay tessellation generated by a stationary Poisson process with intensity  $\lambda$  (Muche, 1996; Mardia *et al.*, 1977):

$$f(l) = \frac{\lambda\pi l}{3} \left\{ \lambda^{1/2} l \exp\left(\frac{-\lambda\pi l^2}{4}\right) + \frac{2}{\sqrt{\pi}} \int_{(\lambda\pi)^{1/2} l/2}^{\infty} \exp(-x^2) dx \right\}, \quad l \geq 0,$$

$$g(\alpha) = \frac{2}{\pi} \{(\pi - 3\alpha) \sin 2\alpha + \cos 2\alpha - \cos 4\alpha\}, \quad 0 \leq \alpha \leq \pi/3.$$

The reason for the difference in their sensitivities can be seen from Figures 1 and 2. For clustered patterns, the distributions of  $L_D$  are somewhat bimodal, including short edges within clusters and long edges between clusters. The distributions of  $L_D$  in regular patterns are unimodal. As a result  $L_D$ , as well as  $L_V$ ,  $R$ ,  $A_D$  and  $P_D$ , are much more sensitive to clustered alternatives than to regular alternatives. However, the distribution of  $\alpha_{\min}$



is unimodal except in cases with strong clustering. Nevertheless, it does not mean that the empirical frequency distributions resemble the function  $g$ . Actually, as can be seen from Figure 2, the empirical frequency distributions of  $\alpha_{\min}$  from either clustered or regular patterns show deviations from  $g$ , meaning that  $\alpha_{\min}$  is sensitive to both alternatives.

From the distributions given in Figures 1 and 2, we can see that the shape of the empirical distributions of these characteristics may be used as a guideline for choosing alternative models if the CSR hypothesis is rejected. If the shape of the empirical distribution of a characteristic in a point pattern resembles a mixture of two distributions, one of which has a small mean and small dispersion and the other has a much larger mean and greater dispersion, then likely the point pattern is a realisation of a process that produces clustered patterns. The two peaks of the distributions give information on the cluster radii and inter-cluster distances. On the other hand, if the empirical distribution has a very narrow range or concentrates only on a limited number of values, then likely the point pattern exhibits regularity.

Figure 1 about here

Figure 2 about here

As a comparison, the powers of the statistic  $\sup_t |\sqrt{\hat{K}(t)/\pi} - r|$ , where  $\hat{K}$  is the empirical Ripley's  $K$ -function with periodic boundary condition, are also given in Tables 1 and 2. We can see that the  $K$ -function was superior to the above characteristics. It is not surprising because it is well-known that the  $K$ -function is rather successful in this aspect, and so this superiority reinforces its practical value. The  $K$ -function, however, will fail if the given point pattern is a realisation of the cell process (Baddeley and Silverman, 1984). The cell

process simultaneously exhibits regularity and clustering but has the same  $K$ -function as the stationary Poisson process, meaning that the  $K$ -function could not capture the dependence structure of such a point process.

Table 3 gives the estimated power in testing CSR by using these characteristics against the cell process. The simulation results here show that  $A_D$  and  $\alpha_{\min}$  are rather able to detect the departure of the cell process from CSR. In order to see if the clustering or regularity nature can be revealed, we compare the empirical frequency distribution of  $L_D$  and  $\alpha_{\min}$  of the cell process given in Figure 3 with those given in Figures 1 and 2. The empirical distribution of  $\alpha_{\min}$  of the cell process resembles those of regular patterns but that of  $L_D$  does not particularly indicate clustering.

Table 3 about here

Figure 3 about here

#### 4. Examples

The empirical frequency distributions of  $L_D$  and  $\alpha_{\min}$  under periodic boundary condition of the three standard examples given in Diggle (1983, pp. 1-2) are given in Figure 4. The  $p$ -values for the null model being CSR are given in Table 4. The locations of Japanese black pine saplings do not show any significant clustering or regularity. The bimodal nature of  $L_D$  and  $\alpha_{\min}$  of redwood seedlings shows significant clustering, whereas the empirical distributions of  $L_D$  and  $\alpha_{\min}$  of the locations of biological cells clearly exhibit regularity.

Figure 4 about here

Table 4 about here

## Acknowledgements

Research supported by the Research Grants Council of the Hong Kong Special Administrative Region, China (Project numbers HKBU 2075/98P and HKBU 2048/02P) and an FRG grant of the Hong Kong Baptist University. I thank Maggie Yu for writing simulation programmes, Fred Hickernell and Regina Hong for their advices on numerical integration of the densities and the referees for their helpful comments and suggestions.

## References

- Baddeley, A. J. and Silverman, B. W., 1984: A cautionary example on the use of second-order methods for analyzing point patterns. *Biometrics* **40**, 1089-1093.
- Boots, B. N., 1975: Patterns of urban settlements revisited. *The Professional Geographer* **27**, 426-431.
- Boots, B. N., 1986: Using angular properties of Delaunay triangles to evaluate point patterns. *Geographical Analysis* **18**, 250-260.
- Brown, G. S., 1965: Point density in stems per acre. *New Zealand Forestry Service Research Notes* **38**, 1-11.
- Diggle, P. J., 1979: On parameter estimation and goodness-of-fit testing for spatial point patterns. *Biometrics* **35**, 87-101.
- Diggle, P. J., 1983: *Statistical Analysis of Spatial Point Patterns*. Academic Press, New York.
- Evans, I. S., 1967: The properties of patterns of points, measured by space filling and angular relationships. *Geographical Articles (Cambridge)* **8**, 63-77.

- Grabarnik, P. and Chiu, S. N., 2002: Goodness-of-fit test for complete spatial randomness against mixtures of regular and clustered spatial point processes. *Biometrika* **89**, 411-421.
- Marcelpoil, R. and Usson, Y., 1992: Methods for the study of cellular sociology: Voronoi diagrams and parametrization of the spatial relationships. *Journal of Theoretical Biology* **154**, 359-369.
- Mardia, K. V., Edwards, R. and Puri, M. L., 1977: Analysis of central place theory. *Bulletin of the International Statistical Institute* **47**, 93-110.
- Møller, J. and Waagepetersen, R. P., 2002: Simulation based inference for spatial point processes. In: M. B. Hansen and J. Møller (eds): *Spatial Statistics and Computational Methods*, Springer, New York, to appear.
- Muche, L., 1996: Distributional properties of the three-dimensional Poisson Delaunay cell. *Journal of Statistical Physics* **84**, 147-167.
- Mugglestone, M. A. and Renshaw, E., 2001: Spectral tests of randomness for spatial point patterns. *Environmental and Ecological Statistics* **8**, 237-251.
- Okabe, A., Boots, B., Sugihara, K. and Chiu, S. N., 2000: *Spatial Tessellations. Concepts and Applications of Voronoi Diagrams*, 2nd edition. Wiley, Chichester.
- Ripley, B. D., 1979: Tests of 'randomness' for spatial point patterns. *Journal of the Royal Statistical Society. Series B* **41**, 368-374.
- Thönnies E. and van Lieshout, M. C., 1999: A comparative study on the power of van Lieshout and Baddeley's  $J$ -function. *Biometrical Journal* **41**, 721-734.
- Vincent, P. J., Howarth, J., Griffiths, J. and Collins, B., 1977: Urban settlement patterns

and the properties of the simplicial graph. *The Professional Geographer* **29**, 21-25.

*Postal address:* S. N. Chiu

*Department of Mathematics*

*Hong Kong Baptist University*

*Kowloon Tong*

*Hong Kong*

*E-mail:* [snchiu@math.hkbu.edu.hk](mailto:snchiu@math.hkbu.edu.hk)

Table 1: Estimated powers (in percent) against simple sequential inhibition alternatives with radius  $r$  (K-S = Kolmogorov-Smirnov statistic; C-vM = Cramér-von Mises statistic)

Characteristic	$r = 0.05$		$r = 0.2$		$r = 0.3$	
	K-S	C-vM	K-S	C-vM	K-S	C-vM
$L_V$	7	5	11	8	10	7
$R$	8	7	18	14	98	84
$A_D$	4	5	74	65	100	100
$P_D$	2	6	9	12	100	99
$L_D$	5	4	3	9	100	99
$\alpha$	8	7	6	7	76	76
$\alpha_{\min}$	6	5	82	84	100	100
$\alpha_{\text{mid}}$	4	5	11	10	45	40
$\alpha_{\max}$	5	3	45	42	96	97
$K$ -function	19		100		100	

Table 2: Estimated powers (in percent) against Matérn cluster alternatives with cluster radius  $r$  and mean cluster size  $\rho$  (K-S = Kolmogorov-Smirnov statistic; C-vM = Cramér-von Mises statistic)

Characteristic	$r = 0.1, \rho = 5$		$r = 0.2, \rho = 5$		$r = 0.2, \rho = 10$	
	K-S	C-vM	K-S	C-vM	K-S	C-vM
$L_V$	98	92	74	71	98	93
$R$	100	100	100	94	100	100
$A_D$	100	100	100	100	100	100
$P_D$	100	100	100	100	100	100
$L_D$	100	100	99	99	100	100
$\alpha$	70	78	22	24	29	31
$\alpha_{\min}$	100	100	73	79	88	90
$\alpha_{\text{mid}}$	67	66	36	30	59	56
$\alpha_{\max}$	100	100	61	68	87	89
$K$ -function	100		100		100	


Table 3: Estimated powers (in percent) against the Baddeley-Silver cell process (K-S = Kolmogorov-Smirnov statistic; C-vM = Cramér-von Mises statistic)

	$L_V$	$R$	$A_D$	$P_D$	$L_D$	$\alpha$	$\alpha_{\min}$	$\alpha_{\text{mid}}$	$\alpha_{\max}$
K-S	16	65	63	45	39	8	68	9	67
C-vM	15	36	58	48	40	8	73	8	62

Table 4: Estimated  $p$ -values (K-S = Kolmogorov-Smirnov statistic; C-vM = Cramér-von Mises statistic) for the CSR of the locations of 65 Japanese black pine saplings, 62 redwood seedlings and 42 biological cell centres (Diggle, 1983, pp. 1-2)

	$L_V$	$R$	$A_D$	$P_D$	$L_D$	$\alpha$	$\alpha_{\min}$	$\alpha_{\text{mid}}$	$\alpha_{\max}$
Japanese black pine saplings									
K-S	0.078	0.235	0.967	0.958	0.755	0.176	0.622	0.798	0.803
C-vM	0.094	0.154	0.966	0.990	0.818	0.229	0.706	0.612	0.798
redwood seedlings									
K-S	0.207	0.023	0.000	0.000	0.000	0.073	0.000	0.000	0.000
C-vM	0.136	0.024	0.000	0.000	0.000	0.095	0.000	0.009	0.000
biological cell centres									
K-S	0.062	0.001	0.000	0.000	0.001	0.022	0.000	0.125	0.000
C-vM	0.048	0.006	0.000	0.000	0.000	0.008	0.000	0.045	0.000






Matérn cluster

Simple sequential inhibition

Figure 1: Empirical frequency distribution of  $L_D$  in Matérn cluster processes with cluster radii (from top to bottom) 0.01, 0.1 and 0.2 and mean cluster size  $\rho$  and simple sequential inhibition processes with radii (from top to bottom) 0.05, 0.2 and 0.3. The solid curves represent the corresponding frequency distribution for a stationary Poisson process.



Matérn cluster

Simple sequential inhibition

Figure 2: Empirical frequency distribution of  $\alpha_{\min}$  in Matérn cluster processes with cluster radii (from top to bottom) 0.01, 0.1 and 0.2 and mean cluster size  $\rho$  and simple sequential inhibition processes with radii (from top to bottom) 0.05, 0.2 and 0.3. The solid curves represent the corresponding frequency distribution for a stationary Poisson process.





Figure 3: Empirical frequency distribution of  $L_D$  and  $\alpha_{\min}$  of 100 realisations of the cell process.



Japanese black pine saplings

Redwood seedlings

Biological cells

Figure 4: Empirical frequency distribution of  $L_D$  (upper row) and  $\alpha_{\min}$  (lower row) of the locations of 65 Japanese black pine saplings, 62 redwood seedlings and 42 biological cell centres (Diggle, 1983, pp. 1-2). The solid curves represent the corresponding frequency distribution for a stationary Poisson process.