

MASTER'S THESIS

Synthesis and Biomedical Studies of Lanthanide-based Probes as Biomarkers

MENG, Ziyuan

Date of Award:
2017

[Link to publication](#)

General rights

Copyright and intellectual property rights for the publications made accessible in HKBU Scholars are retained by the authors and/or other copyright owners. In addition to the restrictions prescribed by the Copyright Ordinance of Hong Kong, all users and readers must also observe the following terms of use:

- Users may download and print one copy of any publication from HKBU Scholars for the purpose of private study or research
- Users cannot further distribute the material or use it for any profit-making activity or commercial gain
- To share publications in HKBU Scholars with others, users are welcome to freely distribute the permanent URL assigned to the publication

Abstract

Primary cilia are organelles protruding from the cell membrane and can act as antennae to get information from the surroundings. Dysfunction of these organelles may result in serious liver and kidney diseases. Thus directly visualizing of primary cilia is of great value for diagnosing ciliopathies and other biomedical research. Normally, visualizing primary needs antibodies labelled with chromophores which are expensive. Here, a europium(III) based cyclen type non-toxic probe HGEu001 is developed for visualizing primary cilia. This non-toxic complex has good quantum yield (around 10%) with high stability in aqueous solution. Co-staining experiments of HGEu001 and its motif complex HGEu002 (control complex) with commercially available primary cilium markers acetylated tubulin or ARL13B in ciliated NIH3T3 cells with a two-photon confocal microscope suggest HGEu001 can specifically bind to primary cilia while HGEu002 do not have such property. Thus HGEu001 can be used for directly visualizing primary cilia with confocal microscopy.

Contents

Chapter 1 Research background	1
Introduction.....	1
1.1 Primary cilia.....	1
1.1.1 The structure of primary cilia	2
1.1.2 Function of primary cilia and diseases caused by its disorders.....	7
1.1.3 Visualizing technique for primary cilia.....	16
1.2 Fluorescence of lanthanide and its bio-imaging application.....	24
1.2.1 Fluorescence, phosphorescence and lanthanide ion luminescence.....	24
1.2.2 Lanthanide-based complexes for bioimaging applications	35
1.3 Research scope	37
Chapter 2. A luminescent lanthanide approach towards direct visualization of primary cilia.....	39
2.1 Synthesis, characterization of lanthanide-based probes for primary cilia	39
2.1.1 Introduction.....	39
2.1.2 Results and discussion	40
2.1.3 Experimental	41
2.1.4 Conclusion	49
Appendix.....	50
2.2. Photophysical properties studies of Lanthanide-based probes for primary cilia.	59

2.2.1 Introduction.....	59
2.2.2 Results and discussion	59
2.2.3 Experimental.....	62
2.2.4 Conclusion	64
2.3. cytotoxicity assays of Lanthanide-based probes for primary cilia.	66
2.3.1 Introduction.....	66
2.3.2 Results and discussion	66
2.3.3 Experimental	67
2.3.4 Conclusion	68
2.4. Cell imaging of Lanthanide-based probes for primary cilia.....	69
2.4.1 Introduction.....	69
2.4.2 Results and discussion	69
2.4.3 Experimental	75
2.4.4 Conclusion	80
Chapter 3. Conclusion and future work.....	81
References.....	83

List of schemes, tables and figures

Scheme 1	The structure of primary cilia.
Scheme 2	Intraocular pathway of ciliogenesis.
Scheme 3	Mechanism of fluorescence illustrated by Bohr model.
Scheme 4	Mechanism of fluorescence and phosphorescence illustrated by Jabłoński diagram
Scheme 5	Mechanism of lanthanide luminescence.
scheme 2.1.1	the synthetic route for HGEu001 and HGEu002.
Table 1	Classification of cilia.
Table 2	Feature and genes related to ciliopathies.
Table 3	Antibodies commercially available for primary cilia staining in fluorescent microscopy.
Table 4	Ground state, excited state, and emitting regions of lanthanide ions.
Table 2.1.1	Solvent gradient of HPLC for the characterization of HGEu001 and HGEu002.
Table 2.2.1	Photophysical properties of the europium complexes HGEu001 and HGEu002.
Table 2.3.1	IC ₅₀ of HGEu001 and HGEu002 against different cell lines after 24-hour treatment. (IC ₅₀ /μM)
Figure 1	Primary cilia under transmission electron microscopy.
Figure 2	Primary cilia under scanning electron microscopy.

- Figure 3 Primary cilia under fluorescent microscopy stained with different methods.
- Figure 4 Some examples of the sensitizing aromatic antennae
- Figure 5 Antenna covalently bind to the cyclen framework.
- Figure 6 Chromosomal DNA specific cyclen based probe
- Figure 7 A series of probe with different amplicities mitochondrial localizing Eu-based probe.
- Figure 8. High molar absorption coefficients antenna based on pyridylalkynylaryl group.
- Figure 9 Cyclin A targeted peptide modified Eu(III) based probes.
- Figure 10 A mitochondria specific probe with TACN framework.
- Figure11 Helicate based Eu(III) probe.
- Figure. 2.1.2 HPLC characterization of HGEu001 and HGEu002.
- Figure. 2.1.3 HRMS(+ESI) spectrum of the complex HGEu001.
- Figure. 2.1.4 HRMS(+ESI) spectrum of the complex HGEu002.
- Figure 2.1.5 ^1H NMR spectrum of compound 2. (400MHz, DMSO- d_6)
- Figure 2.1.6 ^{13}C NMR spectrum of compound 2. (100MHz, DMSO- d_6)
- Figure 2.1.7 ^1H NMR spectrum of compound 3. (400MHz, DMSO- d_6)
- Figure 2.1.8 ^{13}C NMR spectrum of compound 3. (100MHz, DMSO- d_6)
- Figure 2.1.9 ^1H NMR spectrum of compound 4. (400MHz, DMSO- d_6)
- Figure 2.1.10 ^{13}C NMR spectrum of compound 4. (100MHz, DMSO- d_6)
- Figure 2.1.11 ^1H NMR spectrum of compound 5. (400MHz, CDCl_3)

- Figure 2.1.12 ^{13}C NMR spectrum of compound 5. (100MHz, CDCl_3)
- Figure 2.1.13 ^1H NMR spectrum of HGL001. (400MHz, DMSO-d_6)
- Figure 2.1.14 ^{13}C NMR spectrum of HGL001. (100MHz, DMSO-d_6)
- Figure 2.1.15 ^1H NMR spectrum of HGL002. (400MHz, DMSO-d_6)
- Figure 2.1.16 ^{13}C NMR spectrum of HGL002. (100MHz, DMSO-d_6)
- Figure 2.2.1 The absorption spectra of HGEu001 and HGEu002 in aqueous solution (10 μM)
- Figure 2.2.2 The chemical structures and emission spectra of HGEu001 and HGEu002 in aqueous solution on Edinburgh instrument FLS920 spectrophotometer.
- Figure 2.2.3 The emission spectra of HGEu001 and HGEu002 aqueous solution on Horiba Fluorolog-3 spectrophotometer.
- Figure 2.2.4 The two-photon induced emission spectra of HGEu001 and HGEu002 in aqueous solution.
- Figure 2.2.5 The emission decay curve of the complexes HGEu001 and HGEu002 in D_2O and H_2O .
- Figure 2.3.1 Raw data of cytotoxicity of HGEu001 and HGEu002.
- Fig. 2.4.1 The two-photon living cell imaging of HGEu001 and HGEu002 in HeLa cells
- Figure 2.4.2 Linear fluorescence microscopy co-localization images of HGEu001 with Golgi tracker, lysosome tracker, mitochondria tracker.

Figure 2.4.3 Two-photon confocal in vitro images of HGEu001 and HGEu002 with co-localization of green GFP-ARL13B/GFP-IFT88.

Figure 2.4.4 The co-staining experiments of HGEu001 and HGEu002 with primary cilium markers acetylated tubulin or ARL13B in ciliated NIH3T3 cells with a two-photon confocal microscope.