

## MASTER'S THESIS

# Real Space Renormalization Group Transformation from Machine Learning

CHAN, Chak Ming

*Date of Award:*  
2021

[Link to publication](#)

### General rights

Copyright and intellectual property rights for the publications made accessible in HKBU Scholars are retained by the authors and/or other copyright owners. In addition to the restrictions prescribed by the Copyright Ordinance of Hong Kong, all users and readers must also observe the following terms of use:

- Users may download and print one copy of any publication from HKBU Scholars for the purpose of private study or research
- Users cannot further distribute the material or use it for any profit-making activity or commercial gain
- To share publications in HKBU Scholars with others, users are welcome to freely distribute the permanent URL assigned to the publication

# Abstract

Machine learning (ML) is a powerful tool for extracting important features from high dimensional system and performing data compression. The process of capturing relevant features while eliminating irrelevant ones in ML algorithms, such as the neural networks, has a close connection to the renormalization group (RG). Motivated by this, we study the phase transition of the  $q$ -state Potts model using real-space RG transformation obtained from two types of machine learning: supervised and unsupervised learning. We also investigate the RG transformation with vacancy for the first-order phase transition. For supervised learning, we showed that the convolutional neural network (CNN) is capable of classifying phases of the Potts model at different  $q$ . Then, by modifying the structure of the CNN, we are able to obtain RG transformation by training the CNN to classify ordered and disordered Potts configurations. For unsupervised learning, we obtain RG transformation from the restricted Boltzmann machine (RBM) by training it on Potts configurations at transition temperature. Given the RG transformation, we can obtain the critical exponents and estimate the renormalized Hamiltonian using Monte Carlo renormalization group (MCRG) and pseudolikelihood estimation respectively.

# Table of Contents

DECLARATION	i
Abstract	ii
Acknowledgements	iii
Table of Contents	iv
List of Tables	vi
List of Figures	vii
List of abbreviations	x
<b>1 Introduction</b>	<b>1</b>
<b>2 Theory</b>	<b>8</b>
2.1 Renormalization group transformation . . . . .	10
2.2 Fixed point . . . . .	13
2.3 Finite-size scaling . . . . .	16
2.4 First-order phase transition . . . . .	16
<b>3 Methodology and implementation</b>	<b>19</b>
3.1 Convolutional neural network . . . . .	19

3.1.1	Implementation . . . . .	21
3.2	Restricted Boltzmann Machines . . . . .	25
3.3	Monte Carlo Renormalization Group . . . . .	28
3.4	Pseudolikelihood estimation . . . . .	30
<b>4</b>	<b>Results and Discussion</b>	<b>32</b>
4.1	Phase classification with CNN . . . . .	32
4.2	RG transformation from CNN . . . . .	36
4.2.1	Second-order phase transition . . . . .	36
4.2.2	First-order phase transition . . . . .	42
4.2.3	Limitation . . . . .	47
4.3	RG transformation from RBM . . . . .	47
4.3.1	Ising model . . . . .	48
4.3.2	First-order phase transition . . . . .	50
<b>5</b>	<b>Summary</b>	<b>54</b>
	<b>Appendix</b>	<b>60</b>
	<b>List of References</b>	<b>61</b>
	<b>CURRICULUM VITAE</b>	<b>65</b>

# List of Tables

4.1	Training temperatures in unit of $T_c(\infty)$ . . . . .	33
4.2	MCRG result for the Ising model, where $N_r$ is the number of RG iterations performed on the initial system. . . . .	40
4.3	MCRG result for CNN RG transformation with filter size of 4. . .	40
4.4	Probability distribution of $\sigma'$ given initial box configuration $\{\sigma_i\}$ for $2 \times 2$ CNN transformation at $q = 10$ , where $A, B, C$ and $D$ are different number. . . . .	42
4.5	Part of the probability distribution of $\sigma'$ given initial box configuration $\{\sigma_i\}$ for $2 \times 2$ CNN transformation at $q = 10$ , where $A \neq B \neq C$ and not vacant state, $V =$ vacant state, and $E$ denotes any other state . . . . .	43
4.6	Thermal critical exponent for $2 \times 2$ CNN transformation at $q = 10$ and majority-rule transformation. . . . .	44
4.7	Probability distribution of $\sigma'$ given initial box configuration $\{\sigma_i\}$ for $2 \times 2$ CNN transformation at $q = 5$ , where $A \neq B \neq C$ . . . . .	44
4.8	Thermal critical exponent for $2 \times 2$ CNNs and majority-rule transformation at $q = 5$ . . . . .	46
4.9	Average probability of the hidden variable in the vacant state given randomly initialized initial block $v$ . . . . .	52
4.10	Thermal critical exponent from RBM RG transformation. . . . .	53

# List of Figures

1.1	Schematic diagram of an encoder neural network. . . . .	3
1.2	Averaged output of the CNN as a function of $T/J$ for square lattice Ising model. Adapted from “Juan Carrasquilla and Roger G Melko. Machine learning phases of matter. Nature Physics, 13(5):431–434, 2017”. . . . .	5
2.1	Schematic plot of (a) phases at zero external field $h$ and changing temperature $T$ and (b) magnetization $m$ against temperature for $q \leq 4$ Potts model. . . . .	10
2.2	Kadanoff block spin in 2D square lattice. . . . .	12
2.3	Schematic diagram of RG flow. The arrow indicates the direction of flow given by the recursion relation from equation (2.14). The dotted line indicates system with $\frac{K_1}{K_2}$ . . . . .	14
3.1	Conceptual model of CNN. . . . .	20
3.2	An example of the convolution operation. . . . .	20
3.3	Schematic diagram of the CNN used to perform RG transformation. . . . .	23
3.4	An example of convolution for the 4 by 4 Ising model input configuration with vacancy. Each filter is composed of 4 elements of the same value. The stride of the filters is 2, which reduces the linear size of the Ising system by 2. . . . .	24

3.5	Schematic diagram of the RBM. . . . .	25
3.6	Schematic diagram of the partition of the visible configuration and hidden configuration. . . . .	27
3.7	Estimated parameters of the Ising model at different temperatures from pseudolikelihood estimation. . . . .	31
3.8	The first six neighbor interaction. . . . .	31
4.1	Training loss at each epoch for the 10-state Potts system at $L = 32$ . . . . .	33
4.2	Averaged output at different temperature for the 10-state Potts model. Vertical line indicates $T_c$ . . . . .	34
4.3	Data collapse of averaged $P(\text{ordered})$ against $tL^{1/\nu}$ for (a) the Ising model and (b) the 10-state Potts model. . . . .	34
4.4	Averaged output against averaged magnetization at different temperature for (a) the Ising model and (b) 10-state Potts model. . . . .	35
4.5	Value of a hidden unit against $m$ in the fully connected layer of a trained CNN for the 10-state Potts model. . . . .	35
4.6	Coarse-grained configurations from (a) majority-rule transformation and (b) CNN RG transformation with $2 \times 2$ filter. The left most figure is the initial configuration and the subsequent figures are configurations at 1 <sup>st</sup> , 2 <sup>nd</sup> and 3 <sup>rd</sup> coarse-graining iteration respectively. . . . .	37
4.7	(a) $2 \times 2$ convolution filters and (b) $6 \times 6$ convolution filters from CNN on the Ising model. . . . .	38
4.8	Estimated couplings of the renormalized Hamiltonian of the Ising model after one RG transformation. . . . .	39
4.9	Projection of RG flow of the $3 \times 3$ CNN RG transformation onto $K_1$ and $K_2$ . . . . .	40
4.10	Odd terms considered in the MCRG calculation for the Ising model. . . . .	40

4.11	Finite-size scaling of CNN output for (a) $q = 2$ , (b) $q = 3$ and (c) $q = 4$ . . . . .	41
4.12	Finite-size scaling of CNN output for $q = 10$ (a) without vacancy and (b) with vacancy. . . . .	43
4.13	One site entropy $S$ as a function of the temperature. The horizontal bar indicates the maximum possible entropy $S = \ln 10$ . . . . .	44
4.14	Example of coarse grained configurations for (a) CNN without vacancy and (b) CNN with vacancy, where the value 10 represent the vacant site and the configurations on the lower row shows the vacant site in yellow. . . . .	45
4.15	Thermal exponent $y_t$ as a function of the filter parameter $W_1$ , where the horizontal bar indicates the exact $y_t$ . . . . .	46
4.16	Convolution filters from RBM for the Ising model. . . . .	49
4.17	Couplings of the renormalized system of the Ising model at $T_c$ . . . . .	50
4.18	Convolution filters from RBM for the 10-state Potts model without vacancy. To make the filters more interpretable, we performed a permutation on the columns of the filters. . . . .	51
4.19	Convolution filters from RBM for the 10-state Potts model with vacancy. . . . .	52
4.20	Sample configuration at each RG iteration for (a) $q = 5$ with vacancy, and (b) $q = 10$ with vacancy, where $\mu = q$ denote the vacant state. For $q = 10$ , the lower row shows the vacant states for better visibility. . . . .	53