

Effects of 8-week high-intensity interval training and moderate-intensity continuous training on bone metabolism in sedentary young females

Lu, Mingyue; Li, Mingxing; Yi, Longyan; LI, Feifei; Feng, Lin; Ji, Tianyi; Zang, Yanpeng; Qiu, Junqiang

Published in:
Journal of Exercise Science & Fitness

DOI:
[10.1016/j.jesf.2022.01.001](https://doi.org/10.1016/j.jesf.2022.01.001)

Published: 01/04/2022

Document Version:
Publisher's PDF, also known as Version of record


[Link to publication](#)

Citation for published version (APA):
Lu, M., Li, M., Yi, L., LI, F., Feng, L., Ji, T., Zang, Y., & Qiu, J. (2022). Effects of 8-week high-intensity interval training and moderate-intensity continuous training on bone metabolism in sedentary young females. *Journal of Exercise Science & Fitness*, 20(2), 77-83. <https://doi.org/10.1016/j.jesf.2022.01.001>

General rights

Copyright and intellectual property rights for the publications made accessible in HKBU Scholars are retained by the authors and/or other copyright owners. In addition to the restrictions prescribed by the Copyright Ordinance of Hong Kong, all users and readers must also observe the following terms of use:

- Users may download and print one copy of any publication from HKBU Scholars for the purpose of private study or research
- Users cannot further distribute the material or use it for any profit-making activity or commercial gain
- To share publications in HKBU Scholars with others, users are welcome to freely distribute the permanent publication URLs



Contents lists available at ScienceDirect

Journal of Exercise Science & Fitness

journal homepage: www.elsevier.com/locate/jesf

Effects of 8-week High-Intensity Interval Training and Moderate-Intensity Continuous Training on Bone Metabolism in Sedentary Young Females

Mingyue Lu ^{a,1}, Mingxing Li ^{a,1}, Longyan Yi ^b, Feifei Li ^{c,d}, Lin Feng ^a, Tianyi Ji ^a, Yanpeng Zang ^c, Junqiang Qiu ^{a,*}

^a College of Sport Science School, Beijing Sport University, Beijing, China

^b Institute of Sport and Health Sciences, Beijing Sport University, Beijing, China

^c Department of Sport, Physical Education and Health, Hong Kong Baptist University, Hong Kong, China

^d Centre for Health and Exercise Science Research, Hong Kong Baptist University, Hong Kong, China

ARTICLE INFO

Article history:

Received 28 September 2021

Received in revised form

23 December 2021

Accepted 1 January 2022

Available online 5 January 2022

Keywords:

Sedentary

High-intensity interval training

Bone mineral density

Bone metabolism

ABSTRACT

Objective: High-intensity interval training (HIIT) and moderate-intensity continuous training (MICT) have been reported as effective exercise modes on bone metabolism. However, very few studies focused on young women with sedentary behavior. The purpose of this study was to investigate the effects of 8-week HIIT on bone metabolism in sedentary young women.

Methods: 26 healthy, sedentary female participants were randomly assigned to either the HIIT ($n = 13$, age 23.2 ± 2.9 yr, weight 59.2 ± 7.2 kg, height 162.9 ± 3.3 cm, body mass index 22.3 ± 2.7 kg/m²) or MICT ($n = 13$, age 21.9 ± 1.7 yr, weight 59.3 ± 6.6 kg, height 160.9 ± 4.4 cm, body mass index 21.6 ± 2.4 kg/m²) group. Both groups completed 8 weeks (3 sessions/week) of training on the treadmill, where the HIIT group were asked to complete 6×3 -min bouts of running at the intensity of 80–90% maximum oxygen uptake (VO_{2max}) separated by 2-min active recovery at 30–40% VO_{2max} and the MICT group completed 30-min continuous running at the intensity of 60–70% VO_{2max} . The body composition, bone mineral density (BMD), calcaneus quantitative ultrasound, bone turnover markers, and lower limb muscle strength were measured pre and post interventions.

Results: After 8-week interventions, 1) The total body BMD (HIIT, +8.5%; MICT, +5.5%) significantly increased ($p < 0.05$) without difference between the two groups ($p > 0.05$). The calcaneus broadband ultrasound attenuation (CUBA) (HIIT, +16.0%; MICT, +4.6%) and calcaneus stiffness index (CSI) (HIIT, +16.7%; MICT, +2.5%) significantly increased in HIIT group ($p < 0.05$), but not in MICT group ($p > 0.05$). 2) The 1,25-dihydroxyvitamin D₃ ($1,25(OH)_2D_3$) (HIIT, +42.8%; MICT, +24.9%) level increased in both groups with significantly higher changes in HIIT ($p < 0.05$). 3) The score of standing long jump (HIIT, +10.3%; MICT, +3.8%) and vertical jump (HIIT, +5.3%; MICT, +2.0%) increased in both groups with significantly higher changes in HIIT ($p < 0.05$).

Conclusions: It suggested that 8-week HIIT and MICT interventions could improve bone metabolism. Compared with a similar workload of MICT, HIIT elicited superior benefits on bone metabolism.

© 2022 The Society of Chinese Scholars on Exercise Physiology and Fitness. Published by Elsevier (Singapore) Pte Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Osteoporosis is an osteometabolic disease characterized by substantial loss of bone mass and mineral content, microarchitecture

deterioration of bone tissue, changes of bone shape alternations and geometry, affecting bone quality and strength, and increasing fracture risk.^{1–3} It develops slowly over several years until a fall with a sudden bone fracture before clinical diagnosis.¹ Women are at more

* Corresponding author.

E-mail address: qiuqunqiang@bsu.edu.cn (J. Qiu).

¹ Co-first authors.

risk of osteoporosis, particularly under early menopause before 45 years old or ovaries removed.⁴ Young women, however, may also develop osteoporosis when their bone mass density (BMD) falls below the Z-score for a given age and are unable to reach the level of peak bone mass at the corresponding age. Therefore, maximizing BMD at a young age appears to be an effective protective strategy in the prevention of osteoporotic fracture later in life.^{5,6}

People who are less physically active throughout life are more likely to develop osteoporosis.⁷ With advances in technology, screen time, including watching television, using a computer, and playing video games, is becoming a central component of daily life and the most common sedentary behavior.⁸ Accumulating evidence of the link between sedentary behavior and adverse health indicators has perpetuated this interest.^{8–10} Sedentary behavior is defined as activities that consume energy expenditures less than 1.5 metabolic equivalents (METs) while in a sitting or reclining posture.¹¹ Sedentary behavior leads to a lack of systemic muscle activity and a decrease in muscle strength; there is a low load to stimulate bone accretion, which accelerates the loss of BMD and damages bone health.^{12–14}

There is abundant evidence that physical exercise plays an essential role in maintaining or increasing BMD by improving bone metabolism.^{2,15} However, reports have shown that muscle exercise without physical load is ineffective in preventing bone loss.^{16,17} Exercises that stimulate osteogenic formation are usually characterized by dynamic exercise with high intensity and short stimulus duration.^{18,19} High-intensity interval training (HIIT) has attracted increased attention as a possible time-effective alternative to traditional aerobic exercise.²⁰ HIIT is forecasted as the most popular trend in fitness, according to the American College of Sports Medicine (ACSM) Annual Fitness Trend Forecast.²¹ HIIT refers to exercise characterized by relatively short bursts of vigorous activity at an intensity close to that which elicits maximum oxygen uptake (VO_{2max}) (based on an intensity greater than 80% VO_{2max}), interspersed by periods of rest or low-intensity exercise for recovery.^{22–24} In contrast, moderate-intensity continuous training (MICT) is considered traditional aerobic exercise. It is performed as a continuous bout of moderate-intensity aerobic activity at a steady state for a set duration (typically between 20 and 60 min).²⁰ Studies have pointed out that repeated high-intensity exercises can stimulate the bone many times to reach the threshold of bone formation, and intermittent loading cycles determine a greater increase in bone formation than a single, even prolonged cycles; however, continuous stimulation desensitizes osteocytes.^{13,17,25} Hence, HIIT could be a more efficient way to increase BMD and bone metabolism.

There are many indices for reflecting bone metabolism. Dual X-ray absorptiometry (DXA) is currently the most widely used technique for estimating BMD.²⁶ A meta-analysis indicated that quantitative ultrasound also has excellent sensitivity to assess exercise-induced changes in bone status, especially turnover rates in trabecular bone.²⁷ Calcaneus, essentially a trabecular structure, is more metabolically active than cortical bone and highly responsive to metabolic and mechanical stimuli for bone remodeling. It has been recommended as an appropriate skeletal site for assessing the impact of integrated physical activity on bone.²⁸ The calcaneus speed of sound (CSOS) was used to assess the elastic resistance of the bone correlated with the mineral and protein contents of the calcaneus. Calcaneus broadband ultrasound attenuation (CBAU) measures the loss of ultrasound energy due to absorption or dispersion, reflecting the spatial orientation of bone trabeculae. The calcaneus stiffness index (CSI) is a linear combination of CSOS and CBAU.^{28,29} In addition, bone turnover markers reflect a generally dynamic process, including both resorption and formation of bones.^{30,31} Studies have shown that serum osteocalcin (OC), as a

bone turnover marker,³⁰ was positively correlated with BMD. The catabolite pyridinoline (PYD) of ossein is a marker of bone resorption.³² Furthermore, as a stimulant hormone, 1,25-dihydroxyvitamin D₃ (1,25 (OH)₂D₃) can promote the enhancement of osteoblast activity.^{33,34}

Although it was suggested to use a 4–6 month intervention to observe bone remodeling, many studies with exercise interventions shorter than 4 months have increased bone mineral density and plasma bone turnover markers.^{35–37} Additionally, very few studies have focused on young women with sedentary behavior, representing the most critical osteoporosis prevention population. Physical activity at this age can not only maximize peak bone mass but also carry over into later adulthood to lessen the age-associated decline in bone mass.³⁸ Therefore, the purpose of this study was to investigate the effects of an 8-week HIIT program on bone metabolism, including BMD, quantitative ultrasound of the calcaneus, and bone turnover markers, in sedentary young women. Assessing multiple indicators to understand bone metabolism better and discern the effects of HIIT compared with the MICT program was proposed. It was hypothesized that HIIT would have a better effect than MICT on bone metabolism in sedentary young women.

2. Materials and methods

2.1. Participants

Participants were recruited via posters and distribution of flyers on the campus of a local university. The inclusion criteria were as follows: 1) female age of 20–30 years, 2) no regular physical activity habits of moderate intensity (equal to 3.0 to 5.9 METs, such as brisk walking, as a rule of thumb, a person performing moderate-intensity aerobic activity can talk, but not sing, during the activity³⁹) (30 min or more per day at least 3 days per week) during the 3-month period prior to the onset of this experimental protocol, 3) sedentary behavior with energy expenditure less than 1.5 METs with sitting or reclining posture for more than 6 h per day evaluated by acceleration sensor,¹¹ 4) negative responses to all questions on the Physical Activity Readiness Questionnaire, 5) no history of smoking, 6) medical examinations in the past 6 months revealed no abnormalities related to bone and mineral metabolism, no presence of any disease that affects bone metabolism and leads to secondary osteoporosis (e.g., hyperthyroidism and hyperparathyroidism), 7) no hormonal, orthopedic, or cardiovascular diseases, diabetes, hypertension, hyperlipidemia, and polycystic ovary syndrome, 8) no use of prescribed medication may alter bone metabolism (e.g., corticosteroids, estrogen, and thiazide diuretics), no hormonal contraceptives, no calcium, phosphorus, vitamin D, or K supplements. The present study was performed in accordance with the Helsinki Declaration and approved by the Ethical Committee for the Use of Human and Animal Participants in Research of the local university. Fully informed about this study's purpose, procedures, and potential risks, participants provided their written informed consent. During the interventions, no participants quit, all training sessions were completed by all participants, and the participants did not experience any adverse effects (CONSORT flow diagram as shown in Fig. 1).

2.2. Study design and procedures

This study was a randomized, controlled trial with two major procedures, experimental tests and 8-week interventions. Before the experimental tests, a three-axis acceleration sensor was used to screen participants' sedentary behavior for 7 days. Then, participants paid two visits to the laboratory. On the first visit, initial assessments, including body composition, BMD, and lower limb




Fig. 1. Consolidated Standards of Reporting Trials (CONSORT) diagram showing the flow diagram of participants.

muscle strength, were measured, and venous blood was drawn. Free exercise sessions were performed to accustom to the equipment, exercise intensity, and environment. On the second visit, participants performed graded exercise testing (GXT) on a treadmill to evaluate VO_{2max} . The exercise intensities for the interventions were determined accordingly. Each participant completed 8 weeks of HIIT and MICT protocols on the treadmill, 1 session per day, and 3 non-consecutive days per week. After 8 weeks of interventions, assessments, blood sampling, and GXT were performed again to estimate the effects of interventions.

All tests and interventions were conducted at the local sports science research center from April to July, with the air-conditioning temperature controlled at 20 °C and humidity controlled at 50%. Participants were asked to avoid strenuous exercise and alcohol the day before the laboratory visit and refrained from consuming carbonated drinks, caffeine, or other substances that could affect the GXT and blood test results within 2 h before. During the interventions, participants were also asked to avoid any other professional training regimen and nutritional supplement (e.g., calcium, phosphorus, vitamin D, and K) and to maintain their usual dietary intake, activities, and lifestyles.

2.2.1. Sedentary behavior assessment

The three-axis acceleration sensor (ActiGraph wGT3X-BT, ActiGraph, USA) was used to evaluate participants' daily energy expenditure. Data were collected at 60 Hz sampling rate and the time interval set to 1 epoch. All participants wore the accelerometer on their waists for 7 days, except for bathing and sleeping.⁴⁰

2.2.2. VO_{2max}

All participants performed GXT on the treadmill (H/p/cosmos Pulsar 4.0, H/p/cosmos Sports and Medical gmbh, Nussdorf-Traunstein, Germany) to assess VO_{2max} . Participants completed a 5-min warm-up at 6 km/h and then initially ran at 8 km/h. The velocity was increased by 1 km/h every 3 min until the velocity

reached 12 km/h. Then, the gradient was increased by 1% every 3 min until termination.^{41–43} Participants were strongly encouraged verbally to ensure maximum efforts were achieved throughout the test. Oxygen uptake (VO_2) was measured breath-by-breath using a gas analyzer (MAX-II, Physio-Dyne Instrument, New York, USA) and subsequently averaged over 15-sec intervals throughout the test. Heart rate (HR, Polar T34, Polar, Finland) was monitored during the test with an HR transmitter paired with the gas analyzer. The criteria for determinant VO_{2max} included the following: 1) HR reached within $\pm 10 \text{ min}^{-1}$ of the predicted maximal HR ($220 - \text{age}$)⁴⁴ or entered the plateau phase with increasing load; 2) respiratory exchange ratio reached or remained 1.1; 3) the variation range of VO_2 did not exceed 5% or 150 ml/min or 2 ml/min/kg; 4) VO_2 did not increase with the increase of load, and the participants voluntarily stopped the treadmill when they were exhausted. VO_{2max} was determined when at least 3 of the above criteria were met. The value of VO_{2max} was calculated as the highest 15-s value. During the test, if participants had shortness of breath and consciousness, they stopped immediately.

Following the VO_{2max} test, the corresponding velocities in the HIIT and MICT groups were estimated from the linear relationship of steady-state oxygen consumption versus running velocity.

2.2.3. Interventions

All training sessions were conducted on the treadmill. The standardized 5-min warm-up and 10-min final relaxation and cool-down were identical in both groups. For the HIIT group, 6 × 3-min bouts of running at the intensity of 80–90% VO_{2max} (corresponding running velocity: 9.2 ± 0.3 to 10.0 ± 0.4 km/h) separated by 2-min active recovery at 30–40% VO_{2max} (5.5 ± 0.8 to 6.2 ± 0.6 km/h).^{45–47} For MICT group, 30-min continuous running at the intensity of 60–70% VO_{2max} (8.0 ± 0.6 to 8.6 ± 0.5 km/h). The participants trained at the lower intensity border for the first 2 weeks before increasing towards the upper border.

2.3. Measurements

2.3.1. Body composition

Body composition was measured using the bioelectrical impedance method (Inbody 230, Biospace Company, Seoul, Korea). During the measurement, participants were required to fast and to avoid carrying any metal objects. Participants were required to extend their arms and remain still at an angle of 30° with the torso, put their fingers on the electrodes, and step on the metal electrodes with bare feet. The percentages of body fat and lean muscle mass were recorded.

2.3.2. Total, lower body and calcaneus BMD

BMD was measured at the total and lower body by one registered technician using DXA (XR-46, Norland, Wisconsin, USA) with a standardized procedure. Between pre- and post-intervention measurements, the calibration of the densitometer was performed by standard calibration blocks daily. The instrument should be preheated and calibrated before every use. During the test, participants were required to fast and avoid carrying any metal objects and electronic equipment.

Calcaneus BMD, CSOS, CBUA, and CSI were measured using a calcaneus ultrasound densitometer (Achilles EXP II, GE Healthcare, USA) with a standardized procedure. The machine should be preheated and calibrated before every use. Participants were required to be in the sitting position, then put their left foot (without shoes and socks) in the apparatus with coupling agent on the ankle and probe part, and kept motionless.

2.3.3. Bone turnover markers

A venous blood sample (5 ml) was extracted from the antecubital vein after participants fasted for one night (>10 h). Blood samples were coagulated and balanced at room temperature and then centrifuged at 3000 g for 15 min. The obtained serum sample was stored at -80 °C for subsequent analysis. The biomarkers of bone formation and resorption OC were measured with commercial assay kits (Cloud-Clone Corp, Houston, USA), as well as PYD and 1,25(OH)₂D₃ (Shanghai EK-Bioscience Biotechnology Co., Shanghai, China) in a microplate reader (SYNERGY H1, BioTek Company, Vermont, USA).

2.3.4. Lower limb muscle strength

The standing long jump test was measured on a special gym mat. The participants were instructed on the correct technique prior to the test, and they performed several jumps after a warm-up. The participants were required to jump forward vigorously as far as possible from a standing position during the test. Each of them completed 3 effective trials interspersed by a 1-min rest period, and the best performance was included in the analysis. The performance was measured with 1 cm precision.

The vertical jump test was measured in a special vertical jump height instrument. The participants were instructed on the correct technique prior to the test, and they performed several jumps after a warm-up. During the test, the participants were asked to stand barefoot on the mat, naturally separate their feet, bend their legs and squat, and jump up vigorously as high as possible from a standing position while touching the test strip. Each of them completed 3 effective trials interspersed by a 1-min rest period, and the best performance was included in the analysis. The performance was measured with 1 cm precision.

2.4. Statistical analysis

Power analysis was conducted by G*Power version 3.1.9.2 (Universitat Kiel, Germany) to estimate the target sample size.

Using an ANOVA, repeated measures within-between interaction design, 12 participants per group was required as the smallest sample size for an effect size (ES) of 0.53⁴⁸ based on the main outcome (osteocalcin), alpha of 0.05, and 80% power. Finally, 26 participants were included in the initial assessment and then divided into two groups (HIIT or MICT) using the computer-generated random numbers by IBM SPSS Statistics for Windows (v21.0; Armonk, NY).

Data are presented as the means ± standard deviation (SD). Log-transformation was applied prior to further analysis.⁴⁹ An independent sample T test was used to assess the difference in age, height, and METs pre-intervention between the HIIT and MICT groups. Two-way repeated-measure ANOVA was performed to assess the interaction between time (pre- and post-intervention) and group (HIIT and MICT) with post hoc Bonferroni tests when appropriate. Partial eta square (η^2) group × time interaction ES were calculated and interpreted as follows: < 0.06 as small, < 0.14 as moderate, and ≥ 0.14 as large.⁵⁰ The absolute value of each test result was used to calculate the ES for comparisons, represented as Cohen's d. It was interpreted according to the following thresholds: < 0.2 as trivial, 0.2–0.6 as small, 0.6–1.2 as moderate, 1.2–2.0 as large, and > 2.0 as very large.⁵¹ The level of significance was set at $p < 0.05$ for all tests. Statistical analysis was performed by IBM SPSS Statistics for Windows (v21.0; Armonk, NY).

3. Results

All 26 participants completed every training session of 8-week intervention. There were no significant differences in age (HIIT vs MICT: 23.2 ± 2.9 vs 21.9 ± 1.7 years, $p > 0.05$), height (162.9 ± 3.3 vs 160.9 ± 4.4 cm, $p > 0.05$), averaged energy expenditure (1.43 ± 0.03 vs 1.41 ± 0.05 METs, $p > 0.05$), and other assessments pre-intervention listed in [Table 1](#).

After 8-week intervention, VO_{2max} (HIIT, +18.8%; MICT, +17.8%) significantly increased in both groups without between-group difference. The total body BMD (HIIT, +8.5%; MICT, +5.5%) significantly increased in both groups without between-group difference. The CBUA (HIIT, +16.0%; MICT, +4.6%) and CSI (HIIT, +16.7%; MICT, +2.5%) significantly increased in HIIT group, but not in MICT ([Table 1](#)).

Changes in bone turnover markers are summarized in [Table 1](#). The serum level of 1,25(OH)₂D₃ (HIIT, +42.8%; MICT, +24.9%) increased in both groups with significantly greater changes in HIIT ([Table 1](#)).

Changes in indicators of lower limb muscle strength are summarized in [Table 1](#). After 8-week interventions, the score of standing long jump (HIIT, +10.3%; MICT, +3.8%) and vertical jump (HIIT, +5.3%; MICT, +2.0%) increased in both groups with significantly greater changes in HIIT.

4. Discussion

Our findings suggested that both supervised 8-week HIIT and MICT could improve BMD, bone turnover markers, and lower limb muscle strength in sedentary young women; moreover, HIIT did better than MICT in improving bone metabolism.

Available human data show that the magnitude of benefit on bone from exercise is inconsistent and is often influenced by safety concerns (such as “high impact/ground reaction” imposed by HIIT), leading to conservatively prescribed training loads.^{52–54} However, in our study, all 26 participants completed every training session, and no injury or adverse events were reported. It may be that our study adopted a progressive approach (i.e., from lower intensity border to upper border) which might help minimize the risk of

Table 1
All obtained indexes data in the pre- and post-intervention.

	HIIT				MICT				Interaction effect			
	pre	post	Changes (%)	p	ES	pre	post	Changes (%)	p	ES	p	Partial η^2
Weight (kg)	59.22 ± 7.24	57.37 ± 6.64	-3.0 ± 3.1	0.002	0.27	59.27 ± 6.64	57.70 ± 6.38	-2.6 ± 2.7	0.005	0.24	0.76	0.00
BMI (kg/m ²)	22.33 ± 2.73	22.99 ± 3.23	-0.9 ± 0.0	0.001	0.22	21.63 ± 2.42	22.37 ± 3.07	-0.9 ± 0.0	0.006	0.27	0.70	0.00
Body fat (%)	31.43 ± 5.65	28.85 ± 5.52	-8.2 ± 6.0	<0.001	0.46	33.15 ± 4.48	31.48 ± 4.73	-5.0 ± 6.8	<0.001	0.36	0.17	0.08
Lean Mass (kg)	40.31 ± 3.17	40.55 ± 2.27	0.6 ± 3.1	0.585	0.09	39.37 ± 2.27	39.30 ± 2.57	-0.3 ± 3.6	0.876	0.03	0.62	0.01
VO _{2max} (ml/min/kg)	37.45 ± 3.69	44.35 ± 4.72	18.8 ± 9.6	<0.001	1.12	35.56 ± 3.91	42.84 ± 4.49	17.8 ± 11.9	<0.001	1.73	0.78	0.00
HR _{max} (min ⁻¹)	184.00 ± 16.44	173.85 ± 8.82	-5.1 ± 6.3	0.002	0.77	189.23 ± 9.35	179.23 ± 8.82	-5.2 ± 4.0	0.002	1.10	1.00	0.00
Total body BMD (g/cm ²)	0.936 ± 0.050	1.014 ± 0.052*	8.5 ± 3.7	<0.001	1.53	0.947 ± 0.046	1.000 ± 0.053*	5.5 ± 3.3	<0.001	1.07	0.03	0.18
Lower limb BMD (g/cm ²)	1.010 ± 0.079	1.058 ± 0.063	4.8 ± 4.1	<0.001	0.67	0.992 ± 0.077	1.039 ± 0.096	4.4 ± 3.8	<0.001	0.54	0.92	0.00
Calcaneus BMD (g/cm ²)	0.962 ± 0.067	0.971 ± 0.079	0.8 ± 2.3	0.144	0.13	0.968 ± 0.054	0.973 ± 0.054	0.3 ± 1.3	0.552	0.09	0.53	0.02
CSOS (m/s)	1601.42 ± 35.15	1598.88 ± 34.39	-0.2 ± 1.1	0.876	0.07	1593.13 ± 37.19	1599.01 ± 41.62	0.4 ± 2.2	0.533	0.15	0.58	0.01
CBUA (dB/MHz)	121.70 ± 9.47	140.40 ± 9.17*#	16.0 ± 11	<0.001	2.01	112.27 ± 20.94	116.55 ± 17.92	4.6 ± 7.7	0.093	0.22	0.01	0.30
CSI (%)	108.77 ± 10.47	126.38 ± 11.92*#	16.7 ± 10.2	<0.001	1.57	99.46 ± 21.24	101.54 ± 19.89	2.5 ± 3.2	0.224	0.10	0.01	0.50
OC (ug/L)	6.82 ± 1.25	10.01 ± 3.37	50.8 ± 53.7	0.005	1.26	5.74 ± 2.02	7.36 ± 1.40	46.5 ± 63.6	0.019	0.93	0.70	0.01
PYD (nmol/L)	29.93 ± 7.00	25.42 ± 6.79	-15.4 ± 7.4	<0.001	0.65	31.64 ± 6.22	25.13 ± 5.39	-20.4 ± 8.7	<0.001	1.12	0.16	0.08
1,25(OH) ₂ D ₃ (ug/L)	12.10 ± 2.59	17.19 ± 3.44*#	42.8 ± 13.3	<0.001	1.67	12.08 ± 2.11	14.80 ± 2.01*	24.9 ± 22.9	<0.001	1.32	0.01	0.23
Standing long jump (m)	1.61 ± 0.78	1.77 ± 0.83*#	10.3 ± 2.2	<0.001	0.20	1.59 ± 0.13	1.65 ± 0.14*	3.8 ± 2.1	<0.001	0.44	0.01	0.75
Vertical jump (m)	2.30 ± 0.05	2.42 ± 0.05*#	5.3 ± 1.8	<0.001	2.40	2.33 ± 0.04	2.37 ± 0.04*	2.0 ± 1.5	<0.001	1.00	0.01	0.47

Values were summarized as mean ± standard deviation. HIIT = high-intensity interval training, MICT = moderate-intensity continuous training, BMI = body mass index, VO_{2max} = maximum oxygen uptake, HR_{max} = maximum heart rate, BMD = bone mineral density, CSOS = calcaneus speed of sound, CBUA = calcaneus broadband ultrasound attenuation, CSI = calcaneus stiffness index; OC = osteocalcin, PYD = pyridinoline, 1,25(OH)₂D₃ = 1,25-dihydroxyvitamin D₃.

*indicates a significant difference from pre-to post-intervention (*p* < 0.05).

#indicates a significant difference (*p* < 0.05) vs. MICT-group.

adverse events. Thus, our HIIT program may be safe for sedentary female participants.

In our study, the total body BMD value increased over 8 weeks in both groups. This is because physical activity forces can be exerted on bones through ground reaction forces and the contractile activity of muscles,⁵⁵ resulting in a gain of BMD in both groups. Our findings were similar to those observed in other studies. Basat et al.³⁰ reported that 6-month high-impact exercise increased BMD in the lumbar spine and femoral neck regions. There have been few studies on young women, but a study pointed out that high-impact football has a higher BMD among pubertal girls than other sports.⁵⁶ In addition, young female international football players had a 23% higher total body BMD than their untrained and age-matched counterparts.⁵⁷ Furthermore, although our intervention lasted for only 8 weeks, the optimal single exercise duration should be longer than 6 months before significant changes in BMD occur.⁵² It might be that our participants had been sedentary, bones became more sensitive to mechanical loading for sedentary individuals. When their lifestyles were changed, BMD increased more rapidly and significantly during exercise than in the rest of the population. The study showed that people who exercise regularly had greater total body BMD than sedentary individuals.⁵⁶ Additionally, individuals with a low BMD baseline would benefit more from any bone-stimulating activities. Although the average baseline total body BMD level of participants was within the normal range, it was less than 1 g/cm.² Ravnholt et al.³⁵ demonstrated that 7 weeks of the high-intensity intermittent running of untrained adults was effective in increasing BMD. However, after 12 weeks of intensive training, the subjects' total bone mass remained unchanged in the high-intensity training,⁵⁸ which may be related to the small amount of exercise (40 min per week). Regrettably, there was no difference between both the HIIT and MICT groups. It might be that the BMD represents the static state of bone minerals,⁵⁹ and our intervention was only 8 weeks and was unable to make a significant difference between the two groups. However, the time threshold for the significant difference between the two groups of BMD was not known, and future research can focus on this aspect.

Although there was no difference between the two groups in total body BMD, the CBUA and CSI closely significantly increased in

the HIIT group but not in the MICT group. The CBUA and CSI could accurately reflect bone mineral status by abating the influence of soft tissue thickness variations.²⁸ The study showed that CUBA and CSI were excellent sensitives for assessing exercise-induced changes in bone status.^{27,29,60} The study also highlighted that a high-impact exercise regimen could significantly increase CBUA, while no significant changes were observed in the control group.⁶¹ Therefore, HIIT was more effective in improving bone status.

In this study, 1,25(OH)₂D₃, a bone formation marker, increased significantly in the HIIT groups, and bone metabolism improved after 8 weeks of intervention. Notably, the 1,25(OH)₂D₃ concentrations in HIIT were significantly higher than those in the MICT group. Impact force is a relevant element in the stimulation of bone metabolism.³⁰ Under mechanical loading, the bones remodel themselves to repair the microdamage and increase density to prevent any future damage to the same loading sites. In other words, bone increases the threshold of stress tolerance when encountered with high impact mechanical loading.⁶² High-impact physical activities involving ground reaction forces and joint reaction forces have greater loading stimulus on bone, thus promoting bone metabolism and increasing BMD to a greater extent.⁶³ However, 1,25(OH)₂D₃, a dynamic indicator of bone metabolism, can evaluate the dynamic fluctuations in bone turnover and better reflect the dynamic changes of bones.⁵⁹ One previous study pointed out that 7 and 11 weeks of treadmill running exercise increased the OC and 1,25(OH)₂D₃ levels,⁶⁴ although no significant changes in OC were found in our study. Basat et al.³⁰ studied 42 postmenopausal women and divided them into three groups (strengthening exercise; high-impact exercise; no exercise); after 6 months, OC increased only in the high-impact group, and the bone resorption marker significantly decreased. Similarly, Ravnholt et al.³⁵ demonstrated that a 7-week high-intensity interval training elevated the bone formation marker OC. Altogether, high-intensity exercise could improve bone formation marker levels.³⁵ Thus, our findings suggested that HIIT was more osteogenic than MICT. It is known that exercise stimulates bone formation and suppresses bone resorption, resulting in an increased demand for minerals that are satisfied by an increase in serum 1,25(OH)₂D₃ levels and increased intestinal absorption of calcium. HIIT had a greater

impact on bones and muscles than MICT, consistent with the increases in 1,25(OH)₂D₃ observed in the present study.

Efficient muscle strength also affects bone mass.⁶⁵ Sports with high ground reaction forces produce high muscle forces. Exercise activities that combine the mechanical stimulus from ground reaction forces and the tension produced by intense muscle contractions are considered better for bone stimulation.²⁹ Studies have confirmed that muscle strength, significantly lower limb muscle strength, has a significant positive effect on BMD in young women and has become a predictor of BMD.^{66,67} Improving lower limb strength in young women could help increase BMD and prevent osteoporosis later in life.⁶⁷ This study suggested that both types of training could significantly improve lower limb muscle strength, and the HIIT group was superior to the MICT group. Therefore, HIIT elicited superior benefits on bone metabolism. Increased muscle strength also protects the bones and reduces the risk of falls and fractures.²

This study had several limitations. First, although participants were required to avoid other nutritional supplements during the study, we did not record all food and fluid intake during the study period. Therefore, it was recommended that future studies perform a full assessment of participants' diets. Second, the period of bone remodeling required at least 4–6 months, and the intervention period of our study was 8 weeks, which had already had an impact on bone turnover markers or BMD. It might be that the effect of sedentary on bone health is different from the mechanism of bone remodeling. This study was encouraged to repeatedly observe the effect of different intervention periods on bone metabolism and further explore the mechanism of bone metabolism. In addition, our study used a relatively small sample of only sedentary female participants, which limits the generalizability to other populations and the ability to draw some conclusions about the relative potency of HIIT versus that of MICT. Therefore, caution should be taken when generalizing to other populations; simultaneously, we would encourage duplication of this study's protocol with other more populations in future research. Finally, many indicators can be used to assess bone metabolism, such as parathyroid hormone, insulin, OC, 1,25(OH)₂D₃, PYD, cortisol, procollagen type 1 amino-terminal propeptide, carboxy-terminal cross-linking telopeptide of type I collagen, and N-telopeptides of type I collagen. However, we only selected three of them, and there were limitations in the selection of indicators. Future research should focus on more indicators and comprehensively analyze bone metabolism.

5. Conclusions

Our data suggested that 8-week HIIT and MICT interventions could improve bone metabolism. Compared with similar workload MICT, HIIT elicited superior benefits on bone metabolism.

Funding

This work was supported by the National Key R&D Program of China [grant number 2018YFC2000600].

Author statement

MLu, MLi, and FLi wrote the manuscript. MLu, MLi, LY, TJ and JQ conceived and designed the study. MLu, MLi, FLi and TJ was responsible for data collection and data interpretation. MLu, MLi, LY, FLi, LF, TJ, YZ and JQ assisted in revision of the manuscript.

Declaration of competing interest

The authors declare no conflict of interest.

References

- Cosman F, de Beur SJ, LeBoff M, et al. Clinician's guide to prevention and treatment of osteoporosis. *Osteoporos Int*. 2014;25(10):2359–2381. <https://doi.org/10.1007/s00198-014-2794-2>.
- Moreira LD, Oliveira ML, Lirani-Galvão AP, Marin-Mio RV, Santos RN, Lazaretti-Castro M. Physical exercise and osteoporosis: effects of different types of exercises on bone and physical function of postmenopausal women. *Arq Bras Endocrinol Metabol*. 2014;58(5):514–522. <https://doi.org/10.1590/0004-2730000003374>.
- Trajanoska K, Rivadeneira F. The genetic architecture of osteoporosis and fracture risk. *Bone*. 2019;126:2–10. <https://doi.org/10.1016/j.bone.2019.04.005>.
- Rodriguez M, Shoupe D. Surgical menopause. *Endocrinol Metab Clin North Am*. 2015;44(3):531–542. <https://doi.org/10.1016/j.ecl.2015.05.003>.
- Armas LA, Recker RR. Pathophysiology of osteoporosis: new mechanistic insights. *Endocrinol Metab Clin North Am*. 2012;41(3):475–486. <https://doi.org/10.1016/j.ecl.2012.04.006>.
- Habibzadeh N. Effects of two-month walking exercise on bone mass density in young, thin women. *Biomed Hum Kinet*. 2010;2:5–8. <https://doi.org/10.2478/v10101-010-0002-1>.
- Xu J, Lombardi G, Jiao W, Banfi G. Effects of exercise on bone status in female subjects, from young girls to postmenopausal women: an overview of systematic reviews and meta-analyses. *Sports Med*. 2016;46(8):1165–1182. <https://doi.org/10.1007/s40279-016-0494-0>.
- Wang X, Li Y, Fan H. The associations between screen time-based sedentary behavior and depression: a systematic review and meta-analysis. *BMC Pub Health*. 2019;19(1):1524. <https://doi.org/10.1186/s12889-019-7904-9>.
- Bertolini GN, Santos Vrd, Alves MJ, et al. Relation between high leisure-time sedentary behavior and low functionality in older adults. Relação entre alto comportamento sedentário no lazer e baixa funcionalidade de idosos. *Revista Brasileira de Cineantropometria Desempenho Humano*. 2016;18(6):713–721. <https://doi.org/10.5007/1980-0037.2016v18n6p713>, 2016-12.
- Townsend LK, Islam H, Dunn E, Eys M, Robertson-Wilson J, Hazell TJ. Modified sprint interval training protocols. Part II. Psychological responses. *Appl Physiol Nutr Metabol*. 2017;42(4):347–353. <https://doi.org/10.1139/apnm-2016-0479>.
- Bames J, Behrens TK, Benden ME, et al. Letter to the Editor: standardized use of the terms "sedentary" and "sedentary behaviours". *Appl Physiol Nutr Metabol*. 2012;37(2012):540–542. <https://doi.org/10.1139/H2012-024>.
- Koedijk J, van Rijswijk J, Oranje W, et al. Sedentary behaviour and bone health in children, adolescents and young adults: a systematic review. *Osteoporos Int*. 2017;28(9):2507–2519. <https://doi.org/10.1007/s00198-017-4076-2>.
- Aboarrage Junior AM, Teixeira CVLS, dos Santos RN, et al. A high-intensity jump-based aquatic exercise program improves bone mineral density and functional fitness in postmenopausal women. *Rejuvenation Res*. 2018;21(6):535–540. <https://doi.org/10.1089/rej.2018.2069>.
- Villareal DT, Aguirre L, Gurney AB, et al. Aerobic or resistance exercise, or both, in dieting obese older adults. *N Engl J Med*. 2017;376(20):1943–1955. <https://doi.org/10.1056/NEJMoa1616338>.
- Hammami A, Chamari K, Slimani M, Shephard RJ, Bouhler E. Effects of Recreational soccer on physical fitness and health indices in sedentary healthy and unhealthy subjects. *Biol Sport*. 2016;33(2):127–137. <https://doi.org/10.5604/20831862.1198209>.
- Giangregorio L, El-Kotob R. Exercise, muscle, and the applied load-bone strength balance. *Osteoporos Int*. 2017;28(1):21–33. <https://doi.org/10.1007/s00198-016-3780-7>.
- Lombardi G, Sanchis-Gomar F, Perego S, Sansoni V, Banfi G. Implications of exercise-induced adipo-myokines in bone metabolism. *Endocrine*. 2016;54(2):284–305. <https://doi.org/10.1007/s12020-015-0834-0>.
- Mohr M, Helge EW, Petersen LF, et al. Effects of soccer vs swim training on bone formation in sedentary middle-aged women. *Eur J Appl Physiol*. 2015;115(12):2671–2679. <https://doi.org/10.1007/s00421-015-3231-8>.
- Vieira S, Lemes B, Silva Jr JA, Ds B, Fs S. Different land-based exercise training programs to improve bone health in postmenopausal women. *Med Sci Tech*. 2013;54(12):158–163. <https://doi.org/10.12659/mst.889899>.
- Way KL, Sultana RN, Sabag A, Baker MK, Johnson NA. The effect of high intensity interval training versus moderate intensity continuous training on arterial stiffness and 24h blood pressure responses: a systematic review and meta-analysis. *J Sci Med Sport*. 2019;22(4):385–391. <https://doi.org/10.1016/j.jsams.2018.09.228>.
- Thompson WR. Worldwide survey of fitness trends for 2018: the CREP Edition. *ACSM's Health & Fit J*. 2017;21(6):10–19. <https://doi.org/10.1249/FIT.0000000000000341>.
- Andreato LV. High-intensity interval training: methodological considerations for interpreting results and conducting research. *Trends Endocrinol Metab*. 2020;31(11):812–817. <https://doi.org/10.1016/j.tem.2020.08.003>.
- Gibala MJ, McGee SL. Metabolic adaptations to short-term high-intensity interval training: a little pain for a lot of gain? *Exerc Sport Sci Rev*. 2008;36(2):58–63. <https://doi.org/10.1097/JES.0b013e318168ec1f>.
- Gibala MJ, Jones AM. Physiological and performance adaptations to high-intensity interval training. *Nestle Natl Ins workshop ser*. 2013;76:51–60. <https://doi.org/10.1159/000350256>.
- Turner CH, Robling AG. Mechanisms by which exercise improves bone strength. *J Bone Miner Metabol*. 2005;23(suppl):16–22. <https://doi.org/>

- 10.1007/bf03026318.
26. Nogueira RC, Weeks BK, Beck BR. Targeting bone and fat with novel exercise for peripubertal boys: the CAPO kids trial. *Pediatr Exerc Sci*. 2015;27(1):128–139. <https://doi.org/10.1123/pes.2014-0069>.
 27. Babatunde OO, Forsyth JJ. Quantitative Ultrasound and bone's response to exercise: a meta analysis. *Bone*. 2013;53(1):311–318. <https://doi.org/10.1016/j.bone.2012.12.011>.
 28. Zulfarina MS, Sharif R, Syarifah-Noratiqah SB, et al. Modifiable factors associated with bone health in Malaysian adolescents utilising calcaneus quantitative ultrasound. *PLoS One*. 2018;13(8):202–321. <https://doi.org/10.1371/journal.pone.0202321>.
 29. Lara B, Salinero JJ, Gutiérrez J, et al. Influence of endurance running on calcaneal bone stiffness in male and female runners. *Eur J Appl Physiol*. 2016;116(2):327–333. <https://doi.org/10.1007/s00421-015-3285-7>.
 30. Basat H, Esmaeilzadeh S, Eskiyurt N. The effects of strengthening and high-impact exercises on bone metabolism and quality of life in postmenopausal women: a randomized controlled trial. *J Back Musculoskelet Rehabil*. 2013;26(4):427–435. <https://doi.org/10.3233/BMR-130402>.
 31. Moreira LDF, Fronza FCAO, Santos RN, et al. The benefits of a high-intensity aquatic exercise program (HydROS) for bone metabolism and bone mass of postmenopausal women. *J Bone Miner Metabol*. 2014;32(4):411–419. <https://doi.org/10.1007/s00774-013-0509-y>.
 32. Banfi G, Lombardi G, Colombini A, Lippi G. Bone metabolism markers in sports medicine. *Sports Med*. 2010;40(8):697–714. <https://doi.org/10.2165/11533090-000000000-00000>.
 33. Penna G, Amuchastegui S, Giarratana N, et al. 1,25-Dihydroxyvitamin D3 selectively modulates tolerogenic properties in myeloid but not plasmacytoid dendritic cells. *J Immunol*. 2007;178(1):145–153. <https://doi.org/10.4049/jimmunol.178.1.145>.
 34. Siddiqui JA, Partridge NC. Physiological bone remodeling: systemic regulation and growth factor involvement. *Physiology (Bethesda)*. 2016;31(3):233–245. <https://doi.org/10.1152/physiol.00061.2014>.
 35. Ravnholt T, Tybirk J, Jørgensen NR, Bangsbo J. High-intensity intermittent "5-10-15" running reduces body fat, and increases lean body mass, bone mineral density, and performance in untrained subjects. *Eur J Appl Physiol*. 2018;118(6):1–10. <https://doi.org/10.1007/s00421-018-3851-x>.
 36. Hosny IA, Elghawabi HS, Younan WB, Sabbour AA, Gorbial MA. Beneficial impact of aerobic exercises on bone mineral density in obese premenopausal women under caloric restriction. *Skeletal radiology*. 2012;41(4):423–427. <https://doi.org/10.1007/s00256-011-1196-1>.
 37. Hinton PS, Rector RS, Thomas TR. Weight-bearing, aerobic exercise increases markers of bone formation during short-term weight loss in overweight and obese men and women. *Metab Clin Exp*. 2006;55(12):1616–1618. <https://doi.org/10.1016/j.metabol.2006.07.023>.
 38. Wetter AC, Economos CD. Relationship between quantitative ultrasound, anthropometry and sports participation in college aged adults. *Osteoporos Int*. 2004;15(10):799–806. <https://doi.org/10.1007/s00198-004-1607-4>.
 39. US Department of Health and Human Services. *Supporting and Sustaining the 2008 Physical Activity Guidelines for Americans*. Association of State and Territorial Health Officials; 2008.
 40. Vaara JP, Vasankari T, Wyss T, et al. Device-based measures of sedentary time and physical activity are associated with physical fitness and body fat content. *Front spor active liv*. 2020;2(18):1–13. <https://doi.org/10.3389/fspor.2020.587789>.
 41. Bentley DJ, Newell J, Bishop D. Incremental exercise test design and analysis: implications for performance diagnostics in endurance athletes. *Sports Med*. 2007;37(7):575–586. <https://doi.org/10.2165/00007256-200737070-00002>.
 42. Poon ET, Little JP, Sit CH, Wong SH. The effect of low-volume high-intensity interval training on cardiometabolic health and psychological responses in overweight/obese middle-aged men. *J Sports Sci*. 2020;38(17):1997–2004. <https://doi.org/10.1080/02640414.2020.1766178>.
 43. Randell RK, Rollo I, Roberts TJ, Dalrymple KJ, Jeukendrup AE, Carter JM. Maximal fat oxidation rates in an athletic population. *Med Sci Sports Exerc*. 2017;49(1):133–140. <https://doi.org/10.1249/mss.0000000000001084>.
 44. Howley E. Criteria for maximal oxygen uptake : review and commentary. *Med Sci Sports Exerc*. 1995;27(9):1292–1301. <https://doi.org/10.1249/00005768-199509000-00009>.
 45. Weng TP, Huang SC, Chuang YF, Wang JS. Effects of interval and continuous exercise training on CD4 lymphocyte apoptotic and autophagic responses to hypoxic stress in sedentary men. *PLoS One*. 2013;8(11):e80248. <https://doi.org/10.1371/journal.pone.0080248>.
 46. de Souza DC, Matos VAF, Dos Santos VOA, et al. Effects of high-intensity interval and moderate-intensity continuous exercise on inflammatory, leptin, IgA, and lipid peroxidation responses in obese males. *Front Physiol*. 2018;9:1–9. <https://doi.org/10.3389/fphys.2018.00567>.
 47. Santos VOA, Browne RAV, Souza DC, et al. Effects of high-intensity interval and moderate-intensity continuous exercise on physical activity and sedentary behavior levels in inactive obese males: a crossover trial. *J Sports Sci Med*. 2019;18(3):390–398.
 48. Basat H, Esmaeilzadeh S, Eskiyurt N. The effects of strengthening and high-impact exercises on bone metabolism and quality of life in postmenopausal women: a randomized controlled trial. *J Back Musculoskelet Rehabil*. 2013;26(4):427–435. <https://doi.org/10.3233/bmr-130402>.
 49. Li F, Yi L, Yan H, et al. High-sensitivity cardiac troponin T release after a single bout of high-intensity interval exercise in experienced marathon runners. *J Exerc Sci Fit*. 2017;15(2):49–54. <https://doi.org/10.1016/j.jesf.2017.08.001>.
 50. Cohen J. *Statistical Power Analysis for the Behavioral Sciences*. second ed. Statistical power analysis for the behavioral sciences; 1988.
 51. Hopkins WG, Marshall SW, Batterham AM, Hanin J. Progressive statistics for studies in sports medicine and exercise science. *Med Sci Sports Exerc*. 2009;41(1):3–13. <https://doi.org/10.1249/MSS.0b013e31818cb278>.
 52. Ma D, Wu L, He Z. Effects of walking on the preservation of bone mineral density in perimenopausal and postmenopausal women: a systematic review and meta-analysis. *Menopause*. 2013;20(11):1216–1226. <https://doi.org/10.1097/GME.000000000000100>.
 53. Zhao R, Zhao M, Xu Z. The effects of differing resistance training modes on the preservation of bone mineral density in postmenopausal women: a meta-analysis. *Osteoporos Int*. 2015;26(5):1605–1618. <https://doi.org/10.1007/s00198-015-3034-0>.
 54. Martyn-St James M, Carroll S. High-intensity resistance training and postmenopausal bone loss: a meta-analysis. *Osteoporos Int*. 2006;17(8):1225–1240. <https://doi.org/10.1007/s00198-006-0083-4>.
 55. Fernandes MLD, de OMnL, L-GoA Paula, Villa M-MR, dos SRN, Marise L-C. Physical exercise and osteoporosis: effects of different types of exercises on bone and physical function of postmenopausal women. *Arq Bras Endocrinol*. 2014;58(5):514–522. <https://doi.org/10.1590/0004-2730000003374>.
 56. Ubago-Guisado E, Gómez-Cabello A, Sánchez-Sánchez J, García-Unanue J, Gallardo L. Influence of different sports on bone mass in growing girls. *J Sports Sci*. 2015;33(16):1710–1718. <https://doi.org/10.1080/02640414.2015.1004639>.
 57. Jackman SR, Scott S, Randers MB, et al. Musculoskeletal health profile for elite female footballers versus untrained young women before and after 16 weeks of football training. *J Sports Sci*. 2013;31(13):1468–1474. <https://doi.org/10.1080/02640414.2013.796066>.
 58. Nybo L, Sundstrup E, Jakobsen MD, et al. High-intensity training versus traditional exercise interventions for promoting health. *Med Sci Sports Exerc*. 2010;42(10):1951–1958. <https://doi.org/10.1249/MSS.0b013e3181d99203>.
 59. Smith C, Tacey A, Mesinovic J, et al. The effects of acute exercise on bone turnover markers in middle-aged and older adults: a systematic review. *Bone*. 2021;143:1–16. <https://doi.org/10.1016/j.bone.2020.115766>.
 60. Lehtonen-Veromaa M, Möttönen T, Nuotio I, Heinonen OJ, Viikari J. Influence of physical activity on ultrasound and dual-energy X-ray absorptiometry bone measurements in peripubertal girls: a cross-sectional study. *Calcif Tissue Int*. 2000;66(4):248–254. <https://doi.org/10.1007/s002230010050>.
 61. Vainionpää A, Korpelainen R, Leppälouto J, Jämsä T. Effects of high-impact exercise on bone mineral density: a randomized controlled trial in premenopausal women. *Osteoporos Int*. 2005;16(2):191–197. <https://doi.org/10.1007/s00198-004-1659-5>.
 62. Zagdsuren B. *Effects Of Jump Training On Bone Mineral Density In Young Adult Females*. [master's Thesis]. Western Kentucky University; 2014.
 63. Watson S, Weeks B, Weis L, Harding A, Horan S, Beck B. High-intensity resistance and impact training improves bone mineral density and physical function in postmenopausal women with osteopenia and osteoporosis: the LIFTMOR randomized controlled trial. *J Bone Miner Res*. 2019;34(3):572. <https://doi.org/10.1002/jbmr.3659>.
 64. Iwamoto J, Shimamura C, Takeda T, et al. Effects of treadmill exercise on bone mass, bone metabolism, and calciotropic hormones in young growing rats. *J Bone Miner Metabol*. 2004;22(1):26–31. <https://doi.org/10.1007/s00774-003-0443-5>.
 65. Takahata Y, Wang DH, Anai T, Ogino K. Relation of prenatal and postnatal status to calcaneus quantitative ultrasound in adolescents. *Acta Med Okayama*. 2012;66(3):221–229. <https://doi.org/10.18926/amo/48561>.
 66. Ribom E, Ljunggren O, Piehl-Aulin K, et al. Muscle strength correlates with total body bone mineral density in young women but not in men. *Scand J Med Sci Sports*. 2004;14(1):24–29. <https://doi.org/10.1111/j.1600-0838.2003.00346.x>.
 67. Guimarães BR, Pimenta LD, Massini DA, et al. Muscle strength and regional lean body mass influence on mineral bone health in young male adults. *PLoS One*. 2018;13(1):1–13. <https://doi.org/10.1371/journal.pone.0191769>.