

# The Convergence of Collocation Solutions in Continuous Piecewise Polynomial Spaces for Weakly Singular Volterra Integral Equations

Liang, Hui; Brunner, Hermann

*Published in:*  
SIAM Journal on Numerical Analysis

*DOI:*  
[10.1137/19M1245062](https://doi.org/10.1137/19M1245062)

Published: 30/07/2019

*Document Version:*  
Publisher's PDF, also known as Version of record

[Link to publication](#)

*Citation for published version (APA):*  
Liang, H., & Brunner, H. (2019). The Convergence of Collocation Solutions in Continuous Piecewise Polynomial Spaces for Weakly Singular Volterra Integral Equations. *SIAM Journal on Numerical Analysis*, 57(4), 1875-1896. <https://doi.org/10.1137/19M1245062>

## General rights

Copyright and intellectual property rights for the publications made accessible in HKBU Scholars are retained by the authors and/or other copyright owners. In addition to the restrictions prescribed by the Copyright Ordinance of Hong Kong, all users and readers must also observe the following terms of use:

- Users may download and print one copy of any publication from HKBU Scholars for the purpose of private study or research
- Users cannot further distribute the material or use it for any profit-making activity or commercial gain
- To share publications in HKBU Scholars with others, users are welcome to freely distribute the permanent publication URLs

# THE CONVERGENCE OF COLLOCATION SOLUTIONS IN CONTINUOUS PIECEWISE POLYNOMIAL SPACES FOR WEAKLY SINGULAR VOLTERRA INTEGRAL EQUATIONS\*

HUI LIANG<sup>†</sup> AND HERMANN BRUNNER<sup>‡</sup>

**Abstract.** Collocation solutions by globally continuous piecewise polynomials to second-kind Volterra integral equations (VIEs) with smooth kernels are uniformly convergent only for certain sets of collocation points. In this paper we establish the analogous convergence theory for VIEs with weakly singular kernels, for both uniform and graded meshes.

**Key words.** Volterra integral equations, weakly singular kernels, collocation solutions, globally continuous piecewise polynomials, uniform convergence

**AMS subject classifications.** 65R20, 65L20, 65L60

**DOI.** 10.1137/19M1245062

**1. Introduction.** Collocation solutions for second-kind Volterra integral equations (VIEs) of the form

$$(1.1) \quad u(t) = f(t) + \int_0^t (t-s)^{-\alpha} K(t,s)u(s) ds, \quad t \in I := [0, T] \quad (0 \leq \alpha < 1),$$

are usually sought either in the space

$$(1.2) \quad S_{m-1}^{(-1)}(I_h) := \{v : v|_{\sigma_n} \in P_{m-1}(\sigma_n) \quad (0 \leq n \leq N-1)\}$$

of *discontinuous* piecewise polynomials of degree  $m-1 \geq 0$ , or in the space of *globally continuous* piecewise polynomials of degree  $m \geq 1$ ,

$$(1.3) \quad S_m^{(0)}(I_h) := \{v \in C(I) : v|_{\sigma_n} \in P_m(\sigma_n) \quad (0 \leq n \leq N-1)\}.$$

Here,  $I_h := \{t_n : 0 = t_0 < t_1 < \dots < t_N = T\}$  denotes a (uniform or graded) mesh for  $I$  with  $h_n := t_{n+1} - t_n$  and mesh diameter  $h := \max_{(n)} \{h_n\}$ , while  $P_k = P_k(\sigma_n)$  is the linear space of (real) polynomials on  $\sigma_n := [t_n, t_{n+1}]$  of degree not exceeding  $k$ . We observe that the dimensions of these linear spaces are given, respectively, by

$$\dim S_{m-1}^{(-1)}(I_h) = Nm \quad \text{and} \quad \dim S_m^{(0)}(I_h) = Nm + 1.$$

For a given mesh  $I_h$  the set

$$(1.4) \quad X_h := \{t_n + c_i h_n : 0 < c_1 < \dots < c_m \leq 1 \quad (0 \leq n \leq N-1)\}$$

\*Received by the editors February 14, 2019; accepted for publication (in revised form) May 28, 2019; published electronically July 30, 2019.

<https://doi.org/10.1137/19M1245062>

**Funding:** The work of the first author was supported by the National Natural Science Foundation of China grants 11771128 and 11101130, by the China Scholarship Council grant 201708230219, and by the University Nursing Program for Young Scholars with Creative Talents in Heilongjiang Province grant UNPYSCT-2016020. The work of the second author was supported by the Hong Kong Research Grants Council GRF grants HKBU 200113 and 12300014.

<sup>†</sup>School of Science, Harbin Institute of Technology, Shenzhen 518055, China (wise2peak@126.com).

<sup>‡</sup>Department of Mathematics, Hong Kong Baptist University, Hong Kong SAR, China, and Department of Mathematics and Statistics, Memorial University of Newfoundland, St. John's, NL A1C 5S7, Canada (hbrunner@math.hkbu.edu.hk).

will denote the collocation points corresponding to prescribed collocation parameters  $\{c_i\}$ . We note that the cardinality of  $X_h$  is  $|X_h| = Nm$ .

For VIEs (1.1) with  $\alpha = 0$ ,

$$(1.5) \quad u(t) = f(t) + \int_0^t K(t, s)u(s) ds, \quad t \in I,$$

and smooth functions  $K$  and  $f$ , collocation solutions  $u_h \in S_{m-1}^{(-1)}(I_h)$  converge uniformly on  $I$  for any choice of the collocation parameters  $\{c_i\}$  (cf. Brunner [3] or [2, Chap. 2]). This is no longer true for collocation solutions  $u_h \in S_m^{(0)}(I_h)$ : in this case, a necessary and sufficient condition for the uniform convergence of  $u_h$  is given by

$$(1.6) \quad -1 \leq \rho_m := (-1)^m \prod_{i=1}^m \frac{1-c_i}{c_i} \leq 1$$

(cf. Liang and Brunner [7]).

The convergence theory of collocation solutions in the space  $S_{m-1}^{(-1)}(I_h)$  for weakly singular VIEs (1.1) with  $0 < \alpha < 1$  is completely understood (see Brunner, Pedaş, and Vainikko [4]). The analogous convergence analysis for collocation solutions in the space  $S_m^{(0)}(I_h)$  of *globally continuous* piecewise polynomials has remained an open question. It is the aim of the present paper to close this gap, for both uniform and graded meshes.

The outline of the paper is as follows. In section 2, we describe the collocation scheme in continuous piecewise polynomial spaces. In section 3, we state the convergence results for uniform meshes, for both  $m = 1$  and  $m > 1$ , and the convergence results for graded meshes are shown in section 4. The detailed proofs are given in section 5. Section 6 illustrates the theoretical results by means of some numerical examples, and section 7 offers a concluding remark.

**2. Collocation solutions in  $S_m^{(0)}(I_h)$ .** For a given integer  $N \geq 2$  and a real number  $r \geq 1$  we define the mesh

$$(2.1) \quad I_h^r := \left\{ t_n := \left( \frac{n}{N} \right)^r T : n = 0, 1, \dots, N \right\}.$$

If  $r > 1$ ,  $I_h^r$  is a graded mesh on  $[0, T]$  with grading exponent  $r$  (cf. [1]). The mesh is a uniform one when  $r = 1$ , and we then use the notation  $I_h := I_h^1$ . Set  $\sigma_n := [t_n, t_{n+1}]$ ,  $h_n := t_{n+1} - t_n$ , and  $h := \max_{(n)} \{h_n\} = h_{N-1}$ . The solution  $u$  of (1.1) will be approximated by an element  $u_h$  of the continuous piecewise polynomial space  $S_m^{(0)}(I_h^r)$  defined by (1.3), by using the collocation points given by (1.4). Thus,  $u_h$  is defined by the collocation equation

$$(2.2) \quad u_h(t) = f(t) + \int_0^t (t-s)^{-\alpha} K(t, s)u_h(s) ds, \quad t \in X_h,$$

with  $u_h(0) = f(0)$ .

On the subinterval  $\sigma_n$  the local representation of the collocation solution  $u_h$  can be written as

$$(2.3) \quad u_h(t_n + sh_n) = \sum_{j=0}^m L_j(s)U_{n,j}, \quad s \in [0, 1]$$

(cf. [6]), where  $U_{n,0} := u_h(t_n)$ ,  $U_{n,j} := u_h(t_{n,j})$ ,  $t_{n,j} := t_n + c_j h_n$ . The polynomials

$$L_0(s) := (-1)^m \prod_{k=1}^m \frac{s - c_k}{c_k}, \quad L_j(s) := \frac{s}{c_j} \prod_{k=1, k \neq j}^m \frac{s - c_k}{c_j - c_k} \quad (j = 1, \dots, m; s \in [0, 1])$$

denote the (local) Lagrange basis functions with respect to the distinct points  $\{0\} \cup \{c_i\}$ .

Setting  $t = t_{n,i}$  ( $i = 1, \dots, m$ ) in (2.2) we obtain the local collocation equations on  $\sigma_n$ , namely

$$\begin{aligned} U_{n,i} &= f(t_{n,i}) + \int_0^{t_{n,i}} (t_{n,i} - s)^{-\alpha} K(t_{n,i}, s) u_h(s) ds \\ (2.4) \quad &= f(t_{n,i}) + \sum_{l=0}^{n-1} h_l^{1-\alpha} \int_0^1 \left( \frac{t_{n,i} - t_l}{h_l} - s \right)^{-\alpha} K(t_{n,i}, t_l + sh_l) \sum_{j=0}^m L_j(s) U_{l,j} ds \\ &\quad + h_n^{1-\alpha} \int_0^{c_i} (c_i - s)^{-\alpha} K(t_{n,i}, t_n + sh_n) \sum_{j=0}^m L_j(s) U_{n,j} ds \quad (i = 1, \dots, m). \end{aligned}$$

In order to write (2.4) in concise form we set

$$\begin{aligned} \mathbf{e} &:= (1, \dots, 1)^T, \quad \mathbf{F}_n := (f(t_{n,1}), \dots, f(t_{n,m}))^T, \quad \mathbf{U}_n := (U_{n,1}, \dots, U_{n,m})^T, \\ \mathbf{B}_n^{(l)}(\alpha) &:= \begin{pmatrix} \int_0^1 \left( \frac{t_{n,i} - t_l}{h_l} - s \right)^{-\alpha} K(t_{n,i}, t_l + sh_l) L_j(s) ds \\ (i, j = 1, \dots, m) \end{pmatrix} \quad (0 \leq l \leq N-1), \\ \mathbf{B}_n(\alpha) &:= \begin{pmatrix} \int_0^{c_i} (c_i - s)^{-\alpha} K(t_{n,i}, t_n + sh_n) L_j(s) ds \\ (i, j = 1, \dots, m) \end{pmatrix}, \\ \mathbf{C}_n^{(l)}(\alpha) &:= \text{diag} \begin{pmatrix} \int_0^1 \left( \frac{t_{n,i} - t_l}{h_l} - s \right)^{-\alpha} K(t_{n,i}, t_l + sh_l) L_0(s) ds \\ (i = 1, \dots, m) \end{pmatrix} \quad (0 \leq l \leq N-1), \\ \text{and} \\ \mathbf{C}_n(\alpha) &:= \text{diag} \begin{pmatrix} \int_0^{c_i} (c_i - s)^{-\alpha} K(t_{n,i}, t_n + sh_n) L_0(s) ds \\ (i = 1, \dots, m) \end{pmatrix}. \end{aligned}$$

The collocation equation (2.4) then assumes the form

$$(2.5) \quad [\mathbf{I}_m - h_n^{1-\alpha} \mathbf{B}_n(\alpha)] \mathbf{U}_n = \mathbf{F}_n + h_n^{1-\alpha} \mathbf{C}_n(\alpha) \mathbf{e} u_h(t_n) + \sum_{l=0}^{n-1} h_l^{1-\alpha} [\mathbf{C}_n^{(l)}(\alpha) \mathbf{e} u_h(t_l) + \mathbf{B}_n^{(l)}(\alpha) \mathbf{U}_l],$$

where  $\mathbf{I}_m$  denotes the identity matrix in  $\mathbb{R}^{m \times m}$ .

If  $f \in C(I)$  and  $K \in C(D)$  ( $D := \{(t, s) : 0 \leq s \leq t \leq T\}$ ), then standard arguments show that there exists an  $\bar{h} > 0$  so that for any mesh  $I_h^r$  with mesh diameter  $h \in (0, \bar{h})$ , the linear algebraic system (2.5) has a unique solution  $\mathbf{U}_n$  for all  $n = 0, 1, \dots, N-1$ . Hence for all sufficiently small  $h > 0$ , (2.2) determines a unique collocation solution  $u_h \in S_m^{(0)}(I_h^r)$  for (1.1) whose local representation on  $\sigma_n = [t_n, t_{n+1}]$  is given by (2.3).

### 3. The convergence results for uniform meshes.

#### 3.1. Continuous piecewise linear collocation solutions.

**THEOREM 3.1.** *Assume that in (1.1) we have  $f \in C^2(I)$  and  $K \in C^2(D)$ . Let  $u$  and  $u_h \in S_1^{(0)}(I_h^1)$  be the exact solution and the collocation solution on uniform meshes defined by the collocation equation (2.2) (with  $m = 1$ ) whose underlying meshes have mesh diameters  $h < \bar{h}$ . Then for any  $\alpha \in (0, 1)$  in (1.1) there holds*

$$\lim_{h \rightarrow 0} \|u - u_h\|_\infty = 0$$

if and only if the collocation parameter  $\{c_1\}$  satisfies the condition

$$-1 \leq L_0(1) = \frac{1 - c_1}{-c_1} \leq 1 \quad \left( \text{i.e., } c_1 \geq \frac{1}{2} \right).$$

The corresponding attainable order of convergence (with respect to a uniform mesh  $I_h^1$ ) is given by

$$\|u - u_h\|_\infty := \max_{t \in I} |u(t) - u_h(t)| \leq Ch^{1-\alpha}$$

and

$$\|u - u_h\|_{h,\infty} := \max_{t \in I_h^1} |u(t) - u_h(t)| \leq C \begin{cases} h^{2(1-\alpha)} & \text{if } c_1 = 1, \\ h^{1-\alpha} & \text{if } \frac{1}{2} \leq c_1 < 1. \end{cases}$$

Here (and in what follows) the constant  $C$  occurring in the error estimate is a generic constant that is independent of  $h$  and  $N$  (but will in general depend on  $\alpha$ , the grading exponent  $r \geq 1$  in (2.1), and certain derivatives of the exact solution  $u$  of the weakly singular VIE (1.1)).

*Remark 3.2.* Theorem 3.1 reveals that for second-kind VIEs with weakly singular kernels ( $0 < \alpha < 1$  in (1.1)), the sufficient and necessary condition for the convergence of the globally continuous collocation solution  $u_h \in S_1^{(0)}(I_h^1)$  coincides with the one for second-kind VIEs with smooth kernels (see [7]); that is, it does not depend on the value of  $\alpha$ .

#### 3.2. Collocation solutions in $S_m^{(0)}(I_h^1)$ with $m > 1$ .

**THEOREM 3.3.** *Assume that  $f \in C^{m+1}(I)$  and  $K \in C^{m+1}(D)$ , and let  $u$  and  $u_h \in S_m^{(0)}(I_h^1)$  denote, respectively, the exact solution of (1.1) and the corresponding collocation solution defined by (2.2). Then on uniform meshes  $I_h := I_h^1$  with mesh diameters  $h < \bar{h}$  we have, for any  $\alpha \in (0, 1)$ ,*

$$\lim_{h \rightarrow 0} \|u - u_h\|_\infty = 0$$

if and only if the collocation parameters  $\{c_i\}$  are such that

$$-1 \leq L_0(1) = (-1)^m \prod_{i=1}^m \frac{1 - c_i}{-c_i} \leq 1.$$

The corresponding attainable order of convergence is given by

$$\|u - u_h\|_\infty := \max_{t \in I} |u(t) - u_h(t)| \leq Ch^{1-\alpha}$$

and

$$\|u - u_h\|_{h,\infty} := \max_{t \in I_h^1} |u(t) - u_h(t)| \leq C \begin{cases} h^{2(1-\alpha)} & \text{if } c_m = 1, \\ h^{1-\alpha} & \text{if } c_m < 1, \end{cases}$$

regardless of the value of  $m$ .

*Remark 3.4.* We observe that (owing to the low regularity of the solution  $u = u(t)$  at  $t = 0$ ) the convergence order on uniform meshes cannot be improved by increasing the value of  $m$ . This is similar to the situation for collocation in  $S_{m-1}^{(-1)}(I_h^1)$  (cf. Brunner [2, Section 6.2.5]).

#### 4. The convergence results for graded meshes.

**THEOREM 4.1.** *Assume that  $f \in C^{m+2}(I)$ ,  $K \in C^{m+2}(D)$ , and let  $u$  be the corresponding exact solution of (1.1) with  $0 < \alpha < 1$ . If  $u_h \in S_m^{(0)}(I_h^r)$  is the collocation solution determined by the collocation equation (2.2) with underlying graded mesh  $I_h^r$  defined by (2.1) and  $r > 1$ , then for all sufficiently small mesh diameters  $h < \bar{h}$  there holds*

$$\lim_{h \rightarrow 0} \|u - u_h\|_\infty = 0,$$

if and only if the collocation parameters  $\{c_i\}$  satisfy the condition

$$-1 \leq L_0(1) = (-1)^m \prod_{i=1}^m \frac{1 - c_i}{-c_i} \leq 1.$$

The corresponding attainable order of convergence is then given by

$$\|u - u_h\|_\infty := \max_{t \in I} |u(t) - u_h(t)| \leq C \begin{cases} h^{\min\{r(1-\alpha), m+1\}} & \text{if } -1 \leq L_0(1) < 1, \\ h^{\min\{r(1-\alpha), m\}} & \text{if } L_0(1) = 1, \end{cases}$$

and

$$\|u - u_h\|_{h,\infty} := \max_{t \in I_h^r} |u(t) - u_h(t)| \leq Ch^{\min\{2r(1-\alpha), m+1\}} \quad \text{if } c_m = 1.$$

**COROLLARY 4.2.** *Under the conditions of Theorem 4.1, if  $L_0(1) = 0$ , e.g., the collocation parameters  $\{c_i\}$  are taken as the Radau II points, we obtain that*

$$\|u - u_h\|_{h,\infty} := \max_{t \in I_h^r} |u(t) - u_h(t)| \leq Ch^{\min\{2r(1-\alpha), m+1\}}.$$

*Remark 4.3.* Theorem 4.1 shows that for VIEs with weakly singular kernels, the condition for the convergence of the continuous collocation method on graded meshes again only depends on the collocation parameters, but is independent of  $\alpha$  and  $r$ .

*Remark 4.4.* It follows from Theorem 4.1 that for the graded meshes  $I_h^r$  with  $r \geq \frac{m+1}{1-\alpha}$  and  $-1 \leq L_0(1) < 1$ , the highest obtainable order is  $m+1$ . For  $\alpha = 0$ , if the collocation points are taken as Gauss points and  $m$  is even, then an order reduction can be observed (see [7]); for  $0 < \alpha < 1$ , by Theorem 4.1, we also observe a similar order reduction.

**5. Proofs.** We only prove Theorem 4.1 for  $r \geq 1$ , since Theorems 3.1 and 3.3 are direct consequences of Theorem 4.1 with  $r = 1$ .

On the first subinterval  $[t_0, t_1] = [0, h_0]$  it follows, by [5] (see also [2, Theorem 6.1.6]), that the exact solution of (1.1) can be expressed in the form

$$u(t_0 + vh_0) = \sum_{(j,k)_\alpha} \gamma_{j,k}(t_0 + vh_0)^{j+k(1-\alpha)} + h_0^{m+1} Y_{m+1,0}(v; \alpha), \quad v \in [0, 1],$$

where

$$(j, k)_\alpha := \{(j, k) : j, k \in \mathbb{N}_0, j + k(1 - \alpha) < m + 1\},$$

and with obvious adaptation of the meaning of the definition of  $Y_{m+1,0}(v; \alpha)$ . We rewrite this representation in the form

$$u(t_0 + vh_0) = \sum_{(j,k)'_\alpha} \gamma_{j,k}(\alpha) h_0^{j+k(1-\alpha)} v^{j+k(1-\alpha)} + \sum_{(j,k)''_\alpha} \gamma_{j,k}(\alpha) h_0^{j+k(1-\alpha)} v^{j+k(1-\alpha)} + h_0^{m+1} Y_{m+1,0}(v; \alpha), \quad v \in [0, 1],$$

where

$$(j, k)'_\alpha := \{(j, k) : j + k(1 - \alpha) \in \mathbb{N}_0; j + k(1 - \alpha) < m + 1\}$$

and

$$(j, k)''_\alpha := \{(j, k) : j + k(1 - \alpha) \notin \mathbb{N}_0; j + k(1 - \alpha) < m + 1\}.$$

With self-explanatory meaning of the coefficients  $c_{j,k}(\alpha)$  we thus obtain the local representation

$$u(t_0 + vh_0) = \sum_{j=0}^m c_{j,0}(\alpha) v^j + h_0^{1-\alpha} \Phi_{m+1,0}(v; \alpha) + h_0^{m+1} Y_{m+1,0}(v; \alpha), \quad v \in [0, 1],$$

with  $\Phi_{m+1,0}(v; \alpha) := \sum_{(j,k)'_\alpha} c_{j,k}(\alpha) v^{j+k(1-\alpha)}$ . Since  $u_h \in S_m^{(0)}(I_h^r)$ , we also have

$$u_h(t_0 + vh_0) = \sum_{j=0}^m d_{j,0} v^j, \quad v \in [0, 1],$$

where the  $d_{j,0}$  may depend on  $h_0$  but are bounded as  $h \rightarrow 0$ . Set  $e_h := u - u_h$  and  $\beta_{j,0} := c_{j,0} - d_{j,0}$  for each  $j$ . Then

$$(5.1) \quad e_h(t_0 + vh_0) = \sum_{j=0}^m \beta_{j,0} v^j + h_0^{1-\alpha} \Phi_{m+1,0}(v; \alpha) + h_0^{m+1} Y_{m+1,0}(v; \alpha), \quad v \in [0, 1].$$

At the collocation points  $t_{0,i} = t_0 + c_i h_0 = c_i h_0$  in  $[t_0, t_1]$ , (5.1), (1.1), and (2.2) imply that

$$\begin{aligned} e_h(t_0 + c_i h_0) &= \int_0^{c_i h_0} (c_i h_0 - s)^{-\alpha} K(c_i h_0, s) e_h(s) ds \\ &= h_0^{1-\alpha} \int_0^{c_i} (c_i - s)^{-\alpha} K(c_i h_0, s h_0) e_h(s h_0) ds \\ &= h_0^{1-\alpha} \int_0^{c_i} (c_i - s)^{-\alpha} K(c_i h_0, s h_0) \\ &\quad \times \left[ \sum_{j=0}^m \beta_{j,0} s^j + h_0^{1-\alpha} \Phi_{m+1,0}(s; \alpha) + h_0^{m+1} Y_{m+1,0}(s; \alpha) \right] ds, \end{aligned}$$

where we have used (5.1). But from (5.1) one also has

$$e_h(t_0 + c_i h_0) = \sum_{j=0}^m \beta_{j,0} c_i^j + h_0^{1-\alpha} \Phi_{m+1,0}(c_i; \alpha) + h_0^{m+1} Y_{m+1,0}(c_i; \alpha).$$

By the definition of  $\Phi_{m+1,0}(v; \alpha)$ , we have  $\Phi_{m+1,0}(0; \alpha) = 0$ . Since  $e_h(0) = 0$ , taking  $v = 0$  in (5.1), we have

$$\beta_{0,0} = -h_0^{m+1} Y_{m+1,0}(0; \alpha);$$

thus

$$\begin{aligned} & -h_0^{m+1} Y_{m+1,0}(0; \alpha) + \sum_{j=1}^m \beta_{j,0} c_i^j + h_0^{1-\alpha} \Phi_{m+1,0}(c_i; \alpha) + h_0^{m+1} Y_{m+1,0}(c_i; \alpha) \\ & = h_0^{1-\alpha} \int_0^{c_i} (c_i - s)^{-\alpha} K(c_i h_0, s h_0) \left[ -h_0^{m+1} Y_{m+1,0}(0; \alpha) \right. \\ (5.2) \quad & \left. + \sum_{j=1}^m \beta_{j,0} s^j + h_0^{1-\alpha} \Phi_{m+1,0}(s; \alpha) + h_0^{m+1} Y_{m+1,0}(s; \alpha) \right] ds. \end{aligned}$$

Defining the  $m \times m$  matrices

$$V_m := \begin{pmatrix} c_i^j \\ (i, j = 1, \dots, m) \end{pmatrix}, \quad \tilde{B}^{(0)} := \begin{pmatrix} \int_0^{c_i} (c_i - s)^{-\alpha} K(c_i h_0, s h_0) s^j ds \\ (i, j = 1, \dots, m) \end{pmatrix},$$

and setting

$$\beta_0 := (\beta_{1,0}, \dots, \beta_{m,0})^T,$$

(5.2) can be written as

$$\left[ V_m - h_0^{1-\alpha} \tilde{B}^{(0)} \right] \beta_0 = h_0^{1-\alpha} Q_0 - h_0^{m+1} \tilde{Y}_{m+1,0},$$

with obvious meanings of the  $m$ -vectors  $Q_0$  and  $\tilde{Y}_{m+1,0}$ . Hence, following the argument used in [2, p. 380], we find that

$$(5.3) \quad \beta_0 = O(h_0^{1-\alpha}).$$

Since

$$\begin{aligned} h & = h_{N-1} = t_N - t_{N-1} = \left( \frac{N}{N} \right)^r T - \left( \frac{N-1}{N} \right)^r T \\ & = TN^{-r} [N^r - (N-1)^r] = TN^{-r} r (N - \xi)^{r-1}, \end{aligned}$$

where  $\xi \in (0, 1)$ , here we have used the mean-value theorem. So

$$h \leq TN^{-r} r N^{r-1} = Tr N^{-1}$$

and

$$h \geq TN^{-r} r (N-1)^{r-1} \geq Tr N^{-r} \left( N - \frac{N}{2} \right)^{r-1} = Tr 2^{1-r} N^{-1}$$

due to  $N \geq 2$ . Therefore, the quantities  $h$  and  $N^{-1}$  are interchangeable in our error estimates. Thus,

$$h_0 = \left( \frac{1}{N} \right)^r T = O \left( \left( \frac{1}{N} \right)^r \right) = O(h^r).$$



Equation (5.1) implies that the error on  $[t_0, t_1]$  satisfies

$$(5.4) \quad e_h(t_0 + vh_0) = O(h_0^{1-\alpha}) = O(h^{r(1-\alpha)}) \quad \text{for } 0 \leq v \leq 1.$$

It follows from the collocation equation that

$$\begin{aligned} e_h(t_0 + c_i h_0) &= \int_0^{c_i h_0} (c_i h_0 - s)^{-\alpha} K(c_i h_0, s) e_h(s) ds \\ &= h_0^{1-\alpha} \int_0^{c_i} (c_i - s)^{-\alpha} K(c_i h_0, s h_0) e_h(s h_0) ds, \end{aligned}$$

so we obtain a further order  $1 - \alpha$  at the collocation points.

For  $1 \leq n \leq N - 1$ , the collocation error on  $[t_n, t_{n+1}]$  has the local Lagrange representation

$$(5.5) \quad e_h(t_n + v h_n) = L_0(v) e_h(t_n) + \sum_{j=1}^m L_j(v) \varepsilon_{n,j} + h_n^{m+1} R_{m+1,n}(v),$$

where  $\varepsilon_{n,j} := e_h(t_{n,j})$  and

$$R_{m+1,n}(v) := \frac{u^{(m+1)}(\xi_{m,n}(v))}{(m+1)!} v \prod_{i=1}^m (v - c_i), \quad t_n < \xi_{m,n}(v) < t_{n+1}.$$

By [5] (see also [2, Theorem 6.1.6]),

$$(5.6) \quad |R_{m+1,n}(v)| = \left| \frac{u^{(m+1)}(\xi_{m,n}(v))}{(m+1)!} v \prod_{i=1}^m (v - c_i) \right| \leq C t_n^{-\alpha-m}.$$

By (1.1) and (2.2), for  $i = 1, \dots, m$  we have

$$\begin{aligned} (5.7) \quad \varepsilon_{n,i} &= h_n^{1-\alpha} \int_0^{c_i} (c_i - s)^{-\alpha} K(t_{n,i}, t_n + s h_n) L_0(s) ds e_h(t_n) \\ &\quad + h_n^{1-\alpha} \sum_{j=1}^m \int_0^{c_i} (c_i - s)^{-\alpha} K(t_{n,i}, t_n + s h_n) L_j(s) ds \varepsilon_{n,j} \\ &\quad + \sum_{l=1}^{n-1} h_l^{1-\alpha} \int_0^1 \left( \frac{t_{n,i} - t_l}{h_l} - s \right)^{-\alpha} K(t_{n,i}, t_l + s h_l) L_0(s) ds e_h(t_l) \\ &\quad + \sum_{l=1}^{n-1} h_l^{1-\alpha} \sum_{j=1}^m \int_0^1 \left( \frac{t_{n,i} - t_l}{h_l} - s \right)^{-\alpha} K(t_{n,i}, t_l + s h_l) L_j(s) ds \varepsilon_{l,j} \\ &\quad + r_{m+1,n}(\alpha; c_i), \end{aligned}$$

where

$$\begin{aligned} r_{m+1,n}(\alpha; c_i) &:= h_0^{1-\alpha} \int_0^1 \left( \frac{t_{n,i} - t_0}{h_0} - s \right)^{-\alpha} K(t_{n,i}, t_0 + s h_0) e_h(t_0 + s h_0) ds \\ &\quad + h_n^{m+2-\alpha} \int_0^{c_i} (c_i - s)^{-\alpha} K(t_{n,i}, t_n + s h_n) R_{m+1,n}(s) ds \end{aligned}$$

$$+ \sum_{l=1}^{n-1} h_l^{m+2-\alpha} \int_0^1 \left( \frac{t_{n,i} - t_l}{h_l} - s \right)^{-\alpha} K(t_{n,i}, t_l + sh_l) R_{m+1,l}(s) ds.$$

By (5.4), [2, Lemma 6.2.10], and (5.6), we have

$$(5.8) \quad |r_{m+1,n}(\alpha; c_i)| \leq C \left[ h_0^{2(1-\alpha)} n^{-\alpha} + h_n^{m+2-\alpha} t_n^{-m-\alpha} + \sum_{l=1}^{n-1} h_l^{m+2-\alpha} (n-l)^{-\alpha} t_l^{-m-\alpha} \right].$$

We need the following two lemmas. Their proofs use well-known standard arguments.

LEMMA 5.1. For  $1 \leq n \leq N - 1$  and any  $\beta \geq \alpha$ ,

$$h_n^{m+\beta} t_n^{-m-\alpha} \leq (r2^{r-1})^{m+\beta} T^{\beta-\alpha} \frac{n^{r(\beta-\alpha)-(m+\beta)}}{N^{r(\beta-\alpha)}}.$$

In particular,

$$(5.9) \quad h_n^{m+2-\alpha} t_n^{-m-\alpha} \leq (r2^{r-1})^{m+2-\alpha} T^{2(1-\alpha)} \frac{n^{2r(1-\alpha)-(m+2-\alpha)}}{N^{2r(1-\alpha)}},$$

$$(5.10) \quad h_n^{m+1} t_n^{-m-\alpha} \leq (r2^{r-1})^{m+1} T^{1-\alpha} \frac{n^{r(1-\alpha)-(m+1)}}{N^{r(1-\alpha)}}.$$

LEMMA 5.2. For  $1 \leq n \leq N$ ,

$$\sum_{l=1}^n (n-l)^{-\alpha} l^{-(1-\alpha)} \leq \frac{1}{\alpha} + \frac{1}{1-\alpha}.$$

By (5.6) and (5.10) of Lemma 5.1, we have

$$(5.11) \quad |h_n^{m+1} R_{m+1,n}(1)| \leq C h_n^{m+1} t_n^{-\alpha-m} \leq C \frac{n^{r(1-\alpha)-(m+1)}}{N^{r(1-\alpha)}} \leq C h^{\min\{r(1-\alpha), m+1\}}.$$

For (5.8) we obtain, by Lemmas 5.1 and 5.2,

$$(5.12) \quad |r_{m+1,n}(\alpha; c_i)| \leq C \left[ h_0^{2(1-\alpha)} n^{-\alpha} + \frac{n^{2r(1-\alpha)-(m+2-\alpha)}}{N^{2r(1-\alpha)}} + \sum_{l=1}^{n-1} (n-l)^{-\alpha} \frac{l^{2r(1-\alpha)-(m+1)-(1-\alpha)}}{N^{2r(1-\alpha)}} \right] \leq C h^{\min\{2r(1-\alpha), m+1\}}.$$

Case I:  $c_m = 1$ . Here, we have  $e_h(t_n) = \varepsilon_{n-1,m}$ . Setting

$$\varepsilon_n := (\varepsilon_{n,1}, \dots, \varepsilon_{n,m})^T, \quad \mathbf{r}_{m+1,n}(\alpha) := (r_{m+1,n}(\alpha; c_1), \dots, r_{m+1,n}(\alpha; c_m))^T,$$

(5.7) can be written as

$$[\mathbf{I}_m - h_n^{1-\alpha} \mathbf{B}_n(\alpha)] \varepsilon_n = h_n^{1-\alpha} \mathbf{C}_n(\alpha) \mathbf{e} \mathbf{e}_m^T \varepsilon_{n-1} + h_1^{1-\alpha} \mathbf{C}_n^{(1)}(\alpha) \mathbf{e} e_h(t_1)$$

$$(5.13) \quad \begin{aligned} & + \sum_{l=2}^{n-1} h_l^{1-\alpha} \mathbf{C}_n^{(l)}(\alpha) \mathbf{e} \mathbf{e}_m^T \boldsymbol{\varepsilon}_{l-1} + \sum_{l=1}^{n-1} h_l^{1-\alpha} \mathbf{B}_n^{(l)}(\alpha) \boldsymbol{\varepsilon}_l \\ & + \mathbf{r}_{m+1,n}(\alpha). \end{aligned}$$

By [2, Lemma 6.2.10] we obtain the estimate

$$(5.14) \quad \left\| \mathbf{C}_n^{(l)}(\alpha) \right\|_1 \leq C(n-l)^{-\alpha}, \quad \left\| \mathbf{B}_n^{(l)}(\alpha) \right\|_1 \leq C(n-l)^{-\alpha}.$$

LEMMA 5.3. For  $1 \leq l \leq N-1, 1 \leq k \leq N-l-1$ , and any  $\beta \geq 0$ ,

$$\begin{aligned} 1 & \leq \frac{h_{l+k}}{h_l} \leq \left(1 + \frac{k+1}{l}\right)^{r-1}, \\ h_{l+k}^\beta - h_l^\beta & \leq (r-1)\beta \max \left\{ \left(1 + \frac{k+1}{l}\right)^{(r-1)\beta-1}, 1 \right\} \frac{k+1}{l} h_l^\beta. \end{aligned}$$

It thus follows that

$$1 \leq \frac{h_{l+1}}{h_l} \leq \left(1 + \frac{2}{l}\right)^{r-1} \leq 3^{r-1},$$

and hence, by (5.13), (5.14), and (5.12), for  $h < \bar{h}$ ,

$$\|\boldsymbol{\varepsilon}_n\|_1 \leq C \sum_{l=1}^{n-1} h_l^{1-\alpha} (n-l)^{-\alpha} \|\boldsymbol{\varepsilon}_l\|_1 + Ch^{\min\{2r(1-\alpha), m+1\}}.$$

Since  $\|\boldsymbol{\varepsilon}_1\|_1 = O(h^{2r(1-\alpha)})$ , it follows from the discrete Gronwall lemma (see [2, Theorem 6.1.19]) that

$$\|\boldsymbol{\varepsilon}_n\|_1 \leq Ch^{\min\{2r(1-\alpha), m+1\}}.$$

The proof is completed by recalling (5.5), (5.11), and (5.4).

Case II:  $c_m < 1$ . We further divide into the following three cases.

Case (1):  $-1 < L_0(1) < 1$ . By (5.5) and (5.7), we have

$$\begin{aligned} & \begin{pmatrix} 1 & \mathbf{0}_{1 \times m} \\ -h_n^{1-\alpha} \mathbf{C}_n(\alpha) \mathbf{e} & \mathbf{I}_m - h_n^{1-\alpha} \mathbf{B}_n(\alpha) \end{pmatrix} \begin{pmatrix} e_h(t_n) \\ \boldsymbol{\varepsilon}_n \end{pmatrix} \\ & = \begin{pmatrix} L_0(1) & L_1(1) & \dots & L_m(1) \\ \mathbf{0}_{m \times 1} & \mathbf{0}_{m \times 1} & \dots & \mathbf{0}_{m \times 1} \end{pmatrix} \begin{pmatrix} e_h(t_{n-1}) \\ \boldsymbol{\varepsilon}_{n-1} \end{pmatrix} \\ & + \sum_{l=1}^{n-1} h_l^{1-\alpha} \begin{pmatrix} 0 & \mathbf{0}_{1 \times m} \\ \mathbf{C}_n^{(l)}(\alpha) \mathbf{e} & \mathbf{B}_n^{(l)}(\alpha) \end{pmatrix} \begin{pmatrix} e_h(t_l) \\ \boldsymbol{\varepsilon}_l \end{pmatrix} + \begin{pmatrix} h_{n-1}^{m+1} R_{m+1,n-1}(1) \\ \mathbf{r}_{m+1,n} \end{pmatrix}. \end{aligned}$$

It is easy to check that

$$\begin{aligned} & \begin{pmatrix} 1 & \mathbf{0}_{1 \times m} \\ -h_n^{1-\alpha} \mathbf{C}_n(\alpha) \mathbf{e} & \mathbf{I}_m - h_n^{1-\alpha} \mathbf{B}_n(\alpha) \end{pmatrix}^{-1} \\ & = \begin{pmatrix} 1 & \mathbf{0}_{1 \times m} \\ (\mathbf{I}_m - h_n^{1-\alpha} \mathbf{B}_n(\alpha))^{-1} h_n^{1-\alpha} \mathbf{C}_n(\alpha) \mathbf{e} & (\mathbf{I}_m - h_n^{1-\alpha} \mathbf{B}_n(\alpha))^{-1} \end{pmatrix} \\ & = \begin{pmatrix} 1 & \mathbf{0}_{1 \times m} \\ O(h^{1-\alpha}) & \mathbf{I}_m + O(h^{1-\alpha}) \end{pmatrix}, \end{aligned}$$

so

$$\begin{aligned}
 & \begin{pmatrix} e_h(t_n) \\ \boldsymbol{\varepsilon}_n \end{pmatrix} \\
 &= \left[ \begin{pmatrix} L_0(1) & L_1(1) & \cdots & L_m(1) \\ \mathbf{0}_{m \times 1} & \mathbf{0}_{m \times 1} & \cdots & \mathbf{0}_{m \times 1} \end{pmatrix} + O(h^{1-\alpha}) \right] \begin{pmatrix} e_h(t_{n-1}) \\ \boldsymbol{\varepsilon}_{n-1} \end{pmatrix} \\
 (5.15) \quad &+ \sum_{l=1}^{n-1} h_l^{1-\alpha} \begin{pmatrix} 0 & \mathbf{0}_{1 \times m} \\ (1 + O(h^{1-\alpha})) \mathbf{C}_n^{(l)}(\alpha) \mathbf{e} & (1 + O(h^{1-\alpha})) \mathbf{B}_n^{(l)}(\alpha) \end{pmatrix} \begin{pmatrix} e_h(t_l) \\ \boldsymbol{\varepsilon}_l \end{pmatrix} \\
 &+ \begin{pmatrix} 1 & \mathbf{0}_{1 \times m} \\ O(h^{1-\alpha}) & \mathbf{I}_m + O(h^{1-\alpha}) \end{pmatrix} \begin{pmatrix} h_{n-1}^{m+1} R_{m+1, n-1}(1) \\ \mathbf{r}_{m+1, n} \end{pmatrix}.
 \end{aligned}$$

Let

$$\mathbf{M}_m := \begin{pmatrix} L_0(1) & L_1(1) & \cdots & L_m(1) \\ \mathbf{0}_{m \times 1} & \mathbf{0}_{m \times 1} & \cdots & \mathbf{0}_{m \times 1} \end{pmatrix}.$$

Then the  $m+1$  eigenvalues of  $\mathbf{M}_m$  are  $\lambda_1 = 0, \dots, \lambda_m = 0$  and

$$\lambda_{m+1} = L_0(1) = (-1)^m \prod_{i=1}^m \frac{1 - c_i}{c_i} \neq 0,$$

so  $\mathbf{M}_m$  is diagonalizable. Therefore, there exists a nonsingular matrix  $\mathbf{P}_m$ , such that

$$\mathbf{P}_m^{-1} \mathbf{M}_m \mathbf{P}_m = \begin{pmatrix} \lambda_{m+1} & \mathbf{0}_{1 \times m} \\ \mathbf{0}_{m \times 1} & \mathbf{0}_{m \times m} \end{pmatrix},$$

and the collocation solution converges to the exact solution if and only if  $|\lambda_{m+1}| = |L_0(1)| \leq 1$ .

Set

$$X_{m,n} := \left\| \mathbf{P}_m^{-1} \begin{pmatrix} e_h(t_n) \\ \boldsymbol{\varepsilon}_n \end{pmatrix} \right\|_1.$$

Then by (5.15), (5.14), (5.11), and (5.12), we obtain

$$X_{m,n} \leq |L_0(1)| X_{m,n-1} + C \sum_{l=1}^{n-1} h_l^{1-\alpha} (n-l)^{-\alpha} X_{m,l} + Ch^{\min\{r(1-\alpha), m+1\}}.$$

By induction,

$$\begin{aligned}
 X_{m,n} &\leq |L_0(1)|^{n-1} X_{m,1} + C \sum_{k=1}^{n-1} |L_0(1)|^{n-k-1} \left[ \sum_{l=1}^k h_l^{1-\alpha} (k+1-l)^{-\alpha} X_{m,l} \right] \\
 &\quad + C \frac{1 - |L_0(1)|^{n-1}}{1 - |L_0(1)|} h^{\min\{r(1-\alpha), m+1\}} \\
 &= |L_0(1)|^{n-1} X_{m,1} + C \sum_{l=1}^{n-1} \left[ \sum_{k=l}^{n-1} |L_0(1)|^{n-k-1} (k+1-l)^{-\alpha} \right] h_l^{1-\alpha} X_{m,l} \\
 &\quad + Ch^{\min\{r(1-\alpha), m+1\}} \\
 (5.16) \quad &= |L_0(1)|^{n-1} X_{m,1} + C \sum_{l=1}^{n-1} \left[ \sum_{j=1}^{n-l} |L_0(1)|^{n-l-j} j^{-\alpha} \right] h_l^{1-\alpha} X_{m,l}
 \end{aligned}$$





$$\leq Ch^{\min\{r(1-\alpha), m+1\}}.$$

Define

$$\begin{aligned} \tau_l^1(\alpha; c_i) &:= h_0^{1-\alpha} \int_0^1 \left[ \left( \frac{t_{l,i} - t_0}{h_0} - s \right)^{-\alpha} K(t_{l,i}, t_0 + sh_0) \right. \\ &\quad \left. - \left( \frac{t_{l-1,i} - t_0}{h_0} - s \right)^{-\alpha} K(t_{l-1,i}, t_0 + sh_0) \right] e_h(t_0 + sh_0) ds, \\ \tau_l^2(\alpha; c_i) &:= h_l^{m+2-\alpha} \int_0^{c_i} (c_i - s)^{-\alpha} K(t_{l,i}, t_l + sh_l) R_{m+1,l}(s) ds \\ &\quad - h_{l-1}^{m+2-\alpha} \int_0^{c_i} (c_i - s)^{-\alpha} K(t_{l-1,i}, t_{l-1} + sh_{l-1}) R_{m+1,l-1}(s) ds, \\ \tau_l^3(\alpha; c_i) &:= \sum_{k=1}^l h_k^{m+2-\alpha} \int_0^1 \left( \frac{t_{l,i} - t_k}{h_k} - s \right)^{-\alpha} K(t_{l,i}, t_k + sh_k) R_{m+1,k}(s) ds \\ &\quad - \sum_{k=1}^{l-1} h_k^{m+2-\alpha} \int_0^1 \left( \frac{t_{l-1,i} - t_k}{h_k} - s \right)^{-\alpha} K(t_{l-1,i}, t_k + sh_k) R_{m+1,k}(s) ds. \end{aligned}$$

Then

$$(5.23) \quad r_{m+1,l}(\alpha; c_i) - r_{m+1,l-1}(\alpha; c_i) = \tau_l^1(\alpha; c_i) + \tau_l^2(\alpha; c_i) + \tau_l^3(\alpha; c_i).$$

LEMMA 5.6. For  $1 \leq n \leq N$ , we have

$$\sum_{l=1}^n l^{-\alpha} \leq \frac{n^{1-\alpha}}{1-\alpha}, \quad \sum_{l=2}^n l^{-\alpha-1} \leq \frac{1}{\alpha}.$$

By (5.4), [2, Lemma 6.2.10], and Lemma 5.6, we obtain

$$\begin{aligned} \sum_{l=2}^n \left| \tau_l^1(\alpha; c_i) \right| &\leq Ch_0^{2(1-\alpha)} \sum_{l=1}^n \int_0^1 \left| \left( \frac{t_{l,i} - t_0}{h_0} - s \right)^{-\alpha} K(t_{l,i}, t_0 + sh_0) \right| ds \\ (5.24) \quad &\leq Ch_0^{2(1-\alpha)} \sum_{l=1}^n l^{-\alpha} \leq Ch_0^{2(1-\alpha)} n^{1-\alpha} \leq Ch_0^{1-\alpha} \leq Ch^{r(1-\alpha)}. \end{aligned}$$

Using (5.6) and Lemma 5.3, we obtain

$$(5.25) \quad |R_{m+1,l}(1) - R_{m+1,l-1}(1)| \leq Ch_l t_l^{-\alpha-m-1},$$

which, together with Lemma 5.3, (5.6), (5.9) of Lemma 5.1, and Lemma 5.6, yields

$$\begin{aligned} &\sum_{l=2}^n \left| \tau_l^2(\alpha; c_i) \right| \\ &\leq \sum_{l=2}^n \left( h_l^{m+2-\alpha} - h_{l-1}^{m+2-\alpha} \right) \int_0^{c_i} \left| (c_i - s)^{-\alpha} K(t_{l,i}, t_l + sh_l) R_{m+1,l}(s) \right| ds \\ &\quad + \sum_{l=2}^n h_{l-1}^{m+2-\alpha} \int_0^{c_i} \left| (c_i - s)^{-\alpha} [K(t_{l,i}, t_l + sh_l) - K(t_{l-1,i}, t_{l-1} + sh_{l-1})] R_{m+1,l}(s) \right| ds \\ &\quad + \sum_{l=2}^n h_{l-1}^{m+2-\alpha} \int_0^{c_i} \left| (c_i - s)^{-\alpha} K(t_{l-1,i}, t_{l-1} + sh_{l-1}) [R_{m+1,l}(s) - R_{m+1,l-1}(s)] \right| ds \end{aligned}$$

(5.26)

$$\begin{aligned} &\leq C \sum_{l=2}^n \frac{h_l^{m+2-\alpha}}{l} t_l^{-m-\alpha} + C \sum_{l=2}^n h_{l-1}^{m+3-\alpha} t_l^{-m-\alpha} + C \sum_{l=2}^n h_{l-1}^{m+3-\alpha} t_l^{-m-1-\alpha} \\ &\leq C \sum_{l=2}^n \frac{l^{2r(1-\alpha)-(m+1)-(2-\alpha)}}{N^{2r(1-\alpha)}} + C \sum_{l=2}^n h_l \frac{l^{2r(1-\alpha)-(m+2-\alpha)}}{N^{2r(1-\alpha)}} + C \sum_{l=2}^n \frac{l^{2r(1-\alpha)-(m+1)-(2-\alpha)}}{N^{2r(1-\alpha)}} \\ &\leq Ch^{\min\{2r(1-\alpha), m+1\}}. \end{aligned}$$

In addition,

$$\begin{aligned} &\sum_{l=2}^n \left| \tau_l^3(\alpha; c_i) \right| \\ &= \sum_{l=2}^n \left| \sum_{k=1}^l h_k^{m+2-\alpha} \int_0^1 K(t_{l,i}, t_k + sh_k) \left( \frac{t_{l,i} - t_k}{h_k} - s \right)^{-\alpha} R_{m+1,k}(s) ds \right. \\ &\quad \left. - \sum_{k=2}^l h_{k-1}^{m+2-\alpha} \int_0^1 K(t_{l-1,i}, t_{k-1} + sh_{k-1}) \left( \frac{t_{l-1,i} - t_{k-1}}{h_{k-1}} - s \right)^{-\alpha} R_{m+1,k-1}(s) ds \right| \\ &\leq \tau_n^{3,1}(\alpha; c_i) + \tau_n^{3,2}(\alpha; c_i) + \tau_n^{3,3}(\alpha; c_i) + \tau_n^{3,4}(\alpha; c_i) + \tau_n^{3,5}(\alpha; c_i), \end{aligned}$$

where

$$\begin{aligned} \tau_n^{3,1}(\alpha; c_i) &:= \sum_{l=2}^n \left| h_1^{m+2-\alpha} \int_0^1 K(t_{l,i}, t_1 + sh_1) \left( \frac{t_{l,i} - t_1}{h_1} - s \right)^{-\alpha} R_{m+1,1}(s) ds \right|, \\ \tau_n^{3,2}(\alpha; c_i) &:= \sum_{l=2}^n \sum_{k=2}^l (h_k^{m+2-\alpha} - h_{k-1}^{m+2-\alpha}) \\ &\quad \times \int_0^1 \left| K(t_{l,i}, t_k + sh_k) \left( \frac{t_{l,i} - t_k}{h_k} - s \right)^{-\alpha} R_{m+1,k}(s) \right| ds, \\ \tau_n^{3,3}(\alpha; c_i) &:= \sum_{l=2}^n \sum_{k=2}^l h_{k-1}^{m+2-\alpha} \int_0^1 \left| [K(t_{l,i}, t_k + sh_k) - K(t_{l-1,i}, t_{k-1} + sh_{k-1})] \right. \\ &\quad \left. \times \left( \frac{t_{l,i} - t_k}{h_k} - s \right)^{-\alpha} R_{m+1,k}(s) \right| ds, \\ \tau_n^{3,4}(\alpha; c_i) &:= \sum_{l=2}^n \sum_{k=2}^l h_{k-1}^{m+2-\alpha} \int_0^1 \left| K(t_{l-1,i}, t_{k-1} + sh_{k-1}) \left( \frac{t_{l-1,i} - t_{k-1}}{h_{k-1}} - s \right)^{-\alpha} \right. \\ &\quad \left. \times [R_{m+1,k}(s) - R_{m+1,k-1}(s)] \right| ds, \\ \tau_n^{3,5}(\alpha; c_i) &:= \sum_{l=2}^n \sum_{k=2}^l h_{k-1}^{m+2-\alpha} \int_0^1 \left| K(t_{l-1,i}, t_{k-1} + sh_{k-1}) \right. \\ &\quad \left. \times \left[ \left( \frac{t_{l,i} - t_k}{h_k} - s \right)^{-\alpha} - \left( \frac{t_{l-1,i} - t_{k-1}}{h_{k-1}} - s \right)^{-\alpha} \right] R_{m+1,k}(s) \right| ds. \end{aligned}$$

LEMMA 5.7. For  $2 \leq n \leq N - 1$ , we have

$$\sum_{l=1}^{n-1} \sum_{k=1}^{l-1} \frac{h_k^{1-\alpha}}{k} (l - k)^{-\alpha} \leq \frac{2^{(r-1)(1-\alpha)}(rT)^{1-\alpha}}{(1 - \alpha)^2(r - 1)}.$$



By [2, Lemma 6.2.10], (5.6), Lemmas 5.3, 5.1, 5.6, 5.7, and 5.2, and (5.25), and noticing that  $r \geq 1$ , we have

$$\begin{aligned} \tau_n^{3,1}(\alpha; c_i) &\leq C \sum_{l=2}^n h_1^{m+2-\alpha} (l-1)^{-\alpha} t_1^{-\alpha-m} \leq C \sum_{l=2}^n \frac{(l-1)^{-\alpha}}{N^{2r(1-\alpha)}} \leq Ch^{\min\{r(1-\alpha), m+1\}}, \\ \tau_n^{3,2}(\alpha; c_i) &\leq C \sum_{l=2}^n \sum_{k=2}^l \frac{h_{k-1}^{m+2-\alpha}}{k-1} (l-k)^{-\alpha} t_k^{-\alpha-m} \\ &\leq C \sum_{l=2}^n \sum_{k=2}^l \frac{h_{k-1}^{1-\alpha}}{k-1} (l-k)^{-\alpha} \frac{k^{r(1-\alpha)-(m+1)}}{N^{r(1-\alpha)}} \leq Ch^{\min\{r(1-\alpha), m+1\}}, \\ \tau_n^{3,3}(\alpha; c_i) &\leq C \sum_{l=2}^n \sum_{k=2}^l h_{k-1}^{m+2-\alpha} [h_l + h_k] (l-k)^{-\alpha} t_k^{-\alpha-m} \\ &\leq 2C \sum_{l=2}^n \sum_{k=2}^l h_l (l-k)^{-\alpha} \frac{k^{2r(1-\alpha)-(m+1)-(1-\alpha)}}{N^{2r(1-\alpha)}} \leq Ch^{\min\{r(1-\alpha), m+1\}}, \\ \tau_n^{3,4}(\alpha; c_i) &\leq C \sum_{l=2}^n \sum_{k=2}^l h_{k-1}^{m+3-\alpha} (l-k)^{-\alpha} t_{k-1}^{-\alpha-m-1} \\ &\leq C \sum_{l=2}^n \sum_{k=2}^l h_k^{1-\alpha} (l-k)^{-\alpha} \frac{k^{2r(1-\alpha)-(m+1)-1}}{N^{2r(1-\alpha)}} \leq Ch^{\min\{r(1-\alpha), m+1\}}. \end{aligned}$$

In order to estimate  $\tau_n^{3,5}(\alpha; c_i)$ , we need the following two lemmas.

LEMMA 5.8. For  $2 \leq k \leq l \leq N - 1$ ,

$$\frac{t_{l-1,i} - t_{k-1}}{h_{k-1}} - \frac{t_{l,i} - t_k}{h_k} \geq 0.$$

LEMMA 5.9. For  $k \geq 2$ ,

$$\left| h_{k-1}^{m+2-\alpha} t_k^{-\alpha-m} - h_k^{m+2-\alpha} t_{k+1}^{-\alpha-m} \right| \leq C \frac{h_k^{1-\alpha}}{k} \frac{k^{r(1-\alpha)-(m+1)}}{N^{r(1-\alpha)}}.$$

By (5.6), Lemma 5.8, [2, Lemma 6.2.10], (5.9) of Lemma 5.1, and Lemmas 5.9, 5.2, 5.6, and 5.7, we deduce

$$\begin{aligned} \tau_n^{3,5}(\alpha; c_i) &\leq C \sum_{l=2}^n \sum_{k=2}^l \int_0^1 \left( \frac{t_{l,i} - t_k}{h_k} - s \right)^{-\alpha} ds h_{k-1}^{m+2-\alpha} t_k^{-\alpha-m} \\ &\quad - C \sum_{l=2}^n \sum_{k=2}^l \int_0^1 \left( \frac{t_{l-1,i} - t_{k-1}}{h_{k-1}} - s \right)^{-\alpha} ds h_{k-1}^{m+2-\alpha} t_k^{-\alpha-m} \\ &= C \sum_{l=2}^n \sum_{k=2}^l \int_0^1 \left( \frac{t_{l,i} - t_k}{h_k} - s \right)^{-\alpha} ds h_{k-1}^{m+2-\alpha} t_k^{-\alpha-m} \\ &\quad - C \sum_{l=1}^{n-1} \sum_{k=1}^l \int_0^1 \left( \frac{t_{l,i} - t_k}{h_k} - s \right)^{-\alpha} ds h_k^{m+2-\alpha} t_{k+1}^{-\alpha-m} \\ &\leq C \sum_{k=2}^n \int_0^1 \left( \frac{t_{n,i} - t_k}{h_k} - s \right)^{-\alpha} ds h_{k-1}^{m+2-\alpha} t_k^{-\alpha-m} \end{aligned}$$



Thus,

$$(5.27) \quad |e_h(t_n)| \leq C \sum_{l=1}^n |\hat{r}_{m+1,l}(\alpha)| + C \sum_{l=1}^n |\mathbf{L}^T \mathbf{r}_{m+1,l}(\alpha)|,$$

and

$$(5.28) \quad \|\boldsymbol{\varepsilon}_n\|_1 \leq C \|\mathbf{r}_{m+1,n}(\alpha)\|_1 + Ch^{1-\alpha} \sum_{l=1}^n |\hat{r}_{m+1,l}(\alpha)| + Ch^{1-\alpha} \sum_{l=1}^n \|\mathbf{r}_{m+1,l}(\alpha)\|_1.$$

The use of (5.8) and Lemmas 5.6, 5.1, and 5.7 leads to

$$\begin{aligned} & \sum_{l=1}^n |r_{m+1,l}(\alpha; c_i)| \\ & \leq C \sum_{l=1}^n \left[ h_0^{2(1-\alpha)} l^{-\alpha} + h_l^{m+2-\alpha} t_l^{-m-\alpha} + \sum_{k=1}^{l-1} h_k^{m+2-\alpha} (l-k)^{-\alpha} t_k^{-m-\alpha} \right] \\ & \leq Ch_0^{2(1-\alpha)} n^{1-\alpha} + C \sum_{l=1}^n \frac{l^{2r(1-\alpha)-m-(2-\alpha)}}{N^{2r(1-\alpha)}} + C \sum_{l=1}^n \sum_{k=1}^{l-1} h_k^{1-\alpha} (l-k)^{-\alpha} \frac{k^{r(1-\alpha)-m-1}}{N^{r(1-\alpha)}} \\ & \leq Ch^{\min\{r(1-\alpha), m\}}, \end{aligned}$$

which, together with (5.27), (5.28), (5.5), (5.11), (5.4), and the following lemma, allows us to complete the proof.

LEMMA 5.10. *For  $2 \leq n \leq N-1$ , we have*

$$\sum_{l=2}^n h_l^{m+1} |R_{m+1,l}(1)| \leq Ch^{\min\{r(1-\alpha), m\}}.$$

**6. Numerical results.** In this section, we present a representative numerical example to illustrate the foregoing convergence results. For uniform meshes, we choose  $m=2$  and  $m=3$  with  $\alpha=0.5$ . For  $m=2$  we use the Gauss collocation parameters,  $c_1 = \frac{3-\sqrt{3}}{6}$ ,  $c_2 = \frac{3+\sqrt{3}}{6}$ ; the Radau IIA collocation parameters,  $c_1 = \frac{1}{3}$ ,  $c_2 = 1$ ; and three sets of arbitrary collocation parameters,  $c_1 = \frac{1}{4}$ ,  $c_2 = 1$ ;  $c_1 = \frac{1}{4}$ ,  $c_2 = \frac{5}{6}$ ;  $c_1 = \frac{1}{6}$ ,  $c_2 = \frac{1}{2}$ , with  $L_0(1) = 1, 0, 0, \frac{3}{5}, 5$ . For  $m=3$  we use the Gauss collocation parameters,  $c_1 = \frac{5-\sqrt{15}}{10}$ ,  $c_2 = \frac{1}{2}$ ,  $c_3 = \frac{5+\sqrt{15}}{10}$ ; the Radau IIA collocation parameters,  $c_1 = \frac{4-\sqrt{6}}{10}$ ,  $c_2 = \frac{4+\sqrt{6}}{10}$ ,  $c_3 = 1$ ; and three sets of arbitrary collocation parameters,  $c_1 = \frac{1}{3}$ ,  $c_2 = \frac{1}{2}$ ,  $c_3 = 1$ ;  $c_1 = \frac{1}{3}$ ,  $c_2 = \frac{1}{2}$ ,  $c_3 = \frac{8}{9}$ ;  $c_1 = \frac{1}{9}$ ,  $c_2 = \frac{1}{3}$ ,  $c_3 = \frac{1}{2}$ , with  $L_0(1) = -1, 0, 0, \frac{1}{4}, 16$ . For graded meshes, we choose  $m=1$  and  $m=2$  with  $\alpha=0.5$ . For  $m=1$ , we use  $c_1 = 0.1, 0.49, 0.5, 0.8, 1$ , with  $L_0(1) = -9, -\frac{51}{49}, -1, -\frac{1}{4}, 0$ . For  $m=2$ , we use the same values of  $c_1$  and  $c_2$  as those for uniform meshes.

*Example 6.1.* In (1.1) let  $K(t, s) = \frac{1}{10\Gamma(1-\alpha)}$  and  $f(t) = 1$  such that the exact solution  $u(t) = E_{1-\alpha, 1}(\frac{t^{1-\alpha}}{10})$ , where the Mittag-Leffler function  $E_{\mu, \theta}$  is defined by

$$E_{\mu, \theta}(z) := \sum_{k=0}^{\infty} \frac{z^k}{\Gamma(\mu k + \theta)} \quad \text{for } \mu, \theta, z \in \mathbb{R} \quad \text{with } \mu > 0.$$

In Tables 1–2, we use uniform meshes, while in Tables 3–4, we use graded meshes, with  $r = \frac{m+1}{1-\alpha}$ , to list the maximum of the absolute errors at the mesh points. From these tables, we observe that the numerical results agree with our theoretical analysis.

TABLE 1

Uniform mesh: Errors at the mesh points when  $m = 2$  and  $\alpha = 0.5$ .

$N$	Gauss ( $L_0(1) = 1$ )	Radau IIA ( $L_0(1) = 0$ )	$(\frac{1}{4}, 1)$ ( $L_0(1) = 0$ )	$(\frac{1}{4}, \frac{5}{6})$ ( $L_0(1) = \frac{3}{5}$ )	$(\frac{1}{6}, \frac{1}{2})$ ( $L_0(1) = 5$ )
$2^5$	5.1326e-03	3.8744e-06	1.0609e-05	1.9691e-03	7.8034e+19
$2^6$	3.6490e-03	1.9313e-06	5.2714e-06	1.4007e-03	1.1272e+42
$2^7$	2.5901e-03	9.6354e-07	2.6241e-06	9.9457e-04	3.5959e+86
$2^8$	1.8372e-03	4.8102e-07	1.3079e-06	7.0535e-04	5.7691e+175
Order	0.50	1.00	1.00	0.50	-

TABLE 2

Uniform mesh: Errors at the mesh points when  $m = 3$  and  $\alpha = 0.5$ .

$N$	Gauss ( $L_0(1) = -1$ )	Radau IIA ( $L_0(1) = 0$ )	$(\frac{1}{3}, \frac{1}{2}, 1)$ ( $L_0(1) = 0$ )	$(\frac{1}{3}, \frac{1}{2}, \frac{8}{9})$ ( $L_0(1) = \frac{1}{4}$ )	$(\frac{1}{9}, \frac{1}{3}, \frac{1}{2})$ ( $L_0(1) = 16$ )
$2^4$	5.1183e-03	1.0233e-06	9.6554e-06	4.0142e-04	7.2495e+16
$2^5$	3.6465e-03	5.0112e-07	4.7615e-06	2.8548e-04	8.6683e+35
$2^6$	2.5922e-03	2.4690e-07	2.3577e-06	2.0270e-04	1.8474e+74
$2^7$	1.8398e-03	1.2217e-07	1.1708e-06	1.4375e-04	1.2755e+151
Order	0.49	1.00	1.00	0.50	-

TABLE 3

Graded mesh: Errors at the mesh points when  $m = 1$ ,  $\alpha = 0.5$ , and  $r = 4$ .

$N$	$c_1 = 0.1$ ( $L_0(1) = -9$ )	$c_1 = 0.49$ ( $L_0(1) = -\frac{51}{49}$ )	$c_1 = 0.5$ ( $L_0(1) = -1$ )	$c_1 = 0.8$ ( $L_0(1) = -\frac{1}{4}$ )	$c_1 = 1$ ( $L_0(1) = 0$ )
$2^8$	3.2058e+237	9.3908e-04	9.2274e-07	5.4671e-07	6.8104e-08
$2^9$	-	4.7625e+00	2.3063e-07	1.3711e-07	1.7156e-08
$2^{10}$	-	5.9177e+08	5.7653e-08	3.4342e-08	4.3122e-09
$2^{11}$	-	4.7825e+25	1.4413e-08	8.5947e-09	1.0820e-09
Order	-	-	2.00	2.00	1.99

TABLE 4

Graded mesh: Errors at the mesh points when  $m = 2$ ,  $\alpha = 0.5$ , and  $r = 6$ .

$N$	Gauss ( $L_0(1) = 1$ )	Radau IIA ( $L_0(1) = 0$ )	$(\frac{1}{4}, 1)$ ( $L_0(1) = 0$ )	$(\frac{1}{4}, \frac{5}{6})$ ( $L_0(1) = \frac{3}{5}$ )	$(\frac{1}{6}, \frac{1}{2})$ ( $L_0(1) = 5$ )
$2^2$	8.6626e-03	6.1892e-05	1.0288e-04	3.9837e-03	5.1577e-01
$2^3$	2.7930e-03	9.6685e-06	1.4758e-05	7.6029e-04	3.9099e+01
$2^4$	7.6970e-04	1.2945e-06	1.9123e-06	1.0729e-04	1.8153e+06
$2^5$	1.9476e-04	1.4389e-07	2.4403e-07	1.4011e-05	3.2306e+16
Order	1.98	3.17	2.97	2.94	-

**7. Concluding remark.** In Brunner, Pedas, and Vainikko [4] convergence results were also established for collocation solutions in  $S_{m-1}^{(-1)}(I_h)$  for weakly singular VIEs with *logarithmic singularity*,

$$u(t) = f(t) + \int_0^t \log(t-s) K(t,s) u(s) ds, \quad t \in I.$$

The analogous convergence theory for collocation solutions in  $S_m^{(0)}(I_h)$  is an open problem.

**Appendix A. Lemmas of section 5: Selected proofs.**

*Proof of Lemma 5.4.* If  $n = 1$ , (5.17) is clearly true. Otherwise,

$$\begin{aligned} \sum_{j=1}^n \lambda^{n-j} \binom{n}{j}^\alpha &= \sum_{j=1}^{\lfloor \frac{n}{2} \rfloor} \lambda^{n-j} \binom{n}{j}^\alpha + \sum_{j=\lfloor \frac{n}{2} \rfloor+1}^n \lambda^{n-j} \binom{n}{j}^\alpha \\ &\leq \sum_{j=1}^{\lfloor \frac{n}{2} \rfloor} \lambda^{\frac{n}{2}} n^\alpha + \sum_{j=\lfloor \frac{n}{2} \rfloor+1}^n \lambda^{n-j} 2^\alpha \\ &\leq \frac{1}{2} \lambda^{\frac{n}{2}} n^{\alpha+1} + \frac{1}{1-\lambda} 2^\alpha. \end{aligned}$$

Since  $\lambda^{\frac{n}{2}} n^{\alpha+1}$  is the general term of the convergent series  $\sum_{n=1}^\infty \lambda^{\frac{n}{2}} n^{\alpha+1}$ , it is thus bounded. Furthermore, let  $\hat{n} = \hat{n}(\alpha; \lambda) := \left\lfloor \frac{1}{\lambda^{-\frac{1}{2(\alpha+1)} - 1}} \right\rfloor + 1$ . We distinguish between the following two cases:

- If  $n \leq \hat{n}$ , then

$$\frac{1}{2} \lambda^{\frac{n}{2}} n^{\alpha+1} \leq \frac{1}{2} \lambda^{\frac{1}{2}} \hat{n}^{\alpha+1}.$$

- Otherwise,  $\{\frac{1}{2} \lambda^{\frac{n}{2}} n^{\alpha+1}\}$  is a monotonically decreasing sequence, so for all  $n > \hat{n}$ ,

$$\frac{1}{2} \lambda^{\frac{n}{2}} n^{\alpha+1} \leq \frac{1}{2} \lambda^{\hat{n}} \hat{n}^{\alpha+1} < \frac{1}{2} \lambda^{\frac{1}{2}} \hat{n}^{\alpha+1}.$$

Let  $\mu(\alpha; \lambda) := \frac{1}{2} \lambda^{\frac{1}{2}} \hat{n}^{\alpha+1} + \frac{1}{1-\lambda} 2^\alpha (> 1)$ . Then  $\sum_{j=1}^n \lambda^{n-j} \binom{n}{j}^\alpha \leq \mu(\alpha; \lambda)$  for all  $n \geq 1$ . This completes the proof.  $\square$

*Proof of Lemma 5.5.* By (5.6), (5.25), Lemma 5.3, and (5.10) of Lemma 5.1,

$$\begin{aligned} &\sum_{l=2}^n \left| h_l^{m+1} R_{m+1,l}(1) - h_{l-1}^{m+1} R_{m+1,l-1}(1) \right| \\ &= \sum_{l=2}^n \left| (h_l^{m+1} - h_{l-1}^{m+1}) R_{m+1,l}(1) + h_{l-1}^{m+1} (R_{m+1,l}(1) - R_{m+1,l-1}(1)) \right| \\ &\leq C \sum_{l=2}^n \left[ \frac{h_{l-1}^{m+1}}{l-1} t_l^{-\alpha-m} + h_{l-1}^{m+1} h_l t_l^{-\alpha-m-1} \right] \\ &\leq C \sum_{l=2}^n \left[ \frac{l^{r(1-\alpha)-(m+2)}}{N^{r(1-\alpha)}} \right] \leq C \frac{\int_1^{n+1} s^{r(1-\alpha)-(m+2)} ds}{N^{r(1-\alpha)}} \\ &\leq C \frac{(n+1)^{r(1-\alpha)-(m+1)} - 1}{[r(1-\alpha) - (m+1)] N^{r(1-\alpha)}} \leq Ch^{\min\{r(1-\alpha), m+1\}}. \quad \square \end{aligned}$$

*Proof of Lemma 5.8.* By the definitions of the graded mesh  $I_h^r$  and the corresponding stepsizes, we obtain

$$\frac{t_{l-1,i} - t_{k-1}}{h_{k-1}} - \frac{t_{l,i} - t_k}{h_k}$$

$$\begin{aligned}
&= \frac{(l-1)^r + c_i[l^r - (l-1)^r] - (k-1)^r}{k^r - (k-1)^r} - \frac{l^r + c_i[(l+1)^r - l^r] - k^r}{(k+1)^r - k^r} \\
&= \frac{c_i l^r + (1-c_i)(l-1)^r - (k-1)^r}{k^r - (k-1)^r} - \frac{c_i(l+1)^r + (1-c_i)l^r - k^r}{(k+1)^r - k^r},
\end{aligned}$$

so we only need to prove

$$\begin{aligned}
&\left[ c_i l^r + (1-c_i)(l-1)^r - (k-1)^r \right] \left[ (k+1)^r - k^r \right] \\
&\geq \left[ c_i(l+1)^r + (1-c_i)l^r - k^r \right] \left[ k^r - (k-1)^r \right],
\end{aligned}$$

which is equivalent to

$$\begin{aligned}
&\left[ c_i \int_{k-1}^l x^{r-1} dx + (1-c_i) \int_{k-1}^{l-1} x^{r-1} dx \right] \int_k^{k+1} y^{r-1} dy \\
&\geq \left[ c_i \int_k^{l+1} x^{r-1} dx + (1-c_i) \int_k^l x^{r-1} dx \right] \int_{k-1}^k y^{r-1} dy,
\end{aligned}$$

so it is sufficient to prove

$$(A.1) \quad \int_{k-1}^l x^{r-1} dx \int_k^{k+1} y^{r-1} dy \geq \int_k^{l+1} x^{r-1} dx \int_{k-1}^k y^{r-1} dy,$$

$$(A.2) \quad \int_{k-1}^{l-1} x^{r-1} dx \int_k^{k+1} y^{r-1} dy \geq \int_k^l x^{r-1} dx \int_{k-1}^k y^{r-1} dy.$$

We will prove only (A.1) since (A.2) can be established in a similar way. Because (A.1) can be written as

$$\int_{k-1}^l x^{r-1} dx \int_{k-1}^k (y+1)^{r-1} dy \geq \int_{k-1}^l (x+1)^{r-1} dx \int_{k-1}^k y^{r-1} dy,$$

i.e.,

$$\begin{aligned}
&\int_{k-1}^k x^{r-1} dx \int_{k-1}^k (y+1)^{r-1} dy + \int_k^l x^{r-1} dx \int_{k-1}^k (y+1)^{r-1} dy \\
&\geq \int_{k-1}^k (x+1)^{r-1} dx \int_{k-1}^k y^{r-1} dy + \int_k^l (x+1)^{r-1} dx \int_{k-1}^k y^{r-1} dy,
\end{aligned}$$

i.e.,

$$\int_k^l x^{r-1} dx \int_{k-1}^k (y+1)^{r-1} dy \geq \int_k^l (x+1)^{r-1} dx \int_{k-1}^k y^{r-1} dy,$$

i.e.,

$$\int_k^l \int_{k-1}^k (xy+x)^{r-1} dx dy \geq \int_k^l \int_{k-1}^k (xy+y)^{r-1} dx dy,$$

which holds due to

$$k-1 \leq y \leq k \leq x \leq l,$$

thus (A.1) also holds. This completes the proof.  $\square$

*Proof of Lemma 5.9.* By Lemma 5.3, the mean-value theorem, and (5.10) of Lemma 5.1, for  $k \geq 2$ ,

$$\begin{aligned} & |h_{k-1}^{m+2-\alpha} t_k^{-\alpha-m} - h_k^{m+2-\alpha} t_{k+1}^{-\alpha-m}| \\ & \leq (h_k^{m+2-\alpha} - h_{k-1}^{m+2-\alpha}) t_k^{-\alpha-m} + h_k^{m+2-\alpha} (t_k^{-\alpha-m} - t_{k+1}^{-\alpha-m}) \\ & \leq C \frac{h_k^{m+2-\alpha}}{k} t_k^{-\alpha-m} + C h_k^{m+3-\alpha} (t_k + \vartheta h_k)^{-\alpha-m-1} \\ & \leq C \frac{h_k^{m+2-\alpha}}{k} t_k^{-\alpha-m} + C h_k^{m+3-\alpha} t_k^{-\alpha-m-1} \leq C \frac{h_k^{1-\alpha}}{k} \frac{k^{r(1-\alpha)-(m+1)}}{N^{r(1-\alpha)}}, \end{aligned}$$

where  $\vartheta \in (0, 1)$ . □

*Proof of Lemma 5.10.* By (5.6) and (5.10) of Lemma 5.1, we have

$$\begin{aligned} \sum_{l=2}^n h_l^{m+1} |R_{m+1,l}(1)| & \leq C \sum_{l=2}^n h_l^{m+1} t_l^{-m-\alpha} \leq C \sum_{l=2}^n \frac{l^{r(1-\alpha)-(m+1)}}{N^{r(1-\alpha)}} \\ & \leq C \frac{\int_1^{n+1} s^{r(1-\alpha)-(m+1)} ds}{N^{r(1-\alpha)}} = C \frac{(n+1)^{r(1-\alpha)-m} - 1}{[r(1-\alpha) - m] N^{r(1-\alpha)}} \\ & \leq C h^{\min\{r(1-\alpha), m\}}. \end{aligned} \quad \square$$

**Acknowledgments.** The work of the first author was carried out while she was a Visiting Scholar at Memorial University of Newfoundland (December 2017–November 2018); she gratefully acknowledges the hospitality by MUN’s Department of Mathematics and Statistics. Especially, she is also thankful to Professor Ronald Haynes for his invitation to visit MUN.

#### REFERENCES

- [1] H. BRUNNER, *The numerical solution of weakly singular Volterra integral equations by collocation on graded meshes*, Math. Comp., 45 (1985), pp. 417–437, <https://doi.org/10.1090/S0025-5718-1985-0804933-3>.
- [2] H. BRUNNER, *Collocation Methods for Volterra Integral and Related Functional Differential Equations*, Cambridge University Press, Cambridge, 2004.
- [3] H. BRUNNER, *On the divergence of collocation solutions in smooth piecewise polynomial spaces for Volterra integral equations*, BIT, 44 (2004), pp. 631–650, <https://doi.org/10.1007/s10543-004-3828-5>.
- [4] H. BRUNNER, A. PEDAS, AND G. VAINIKKO, *The piecewise collocation method for nonlinear weakly singular Volterra equations*, Math. Comp., 68 (1999), pp. 1079–1095, <https://doi.org/10.1090/S0025-5718-99-01073-X>.
- [5] Y. CAO, T. HERDMAN, AND Y. XU, *A hybrid collocation method for Volterra integral equations with weakly singular kernels*, SIAM J. Numer. Anal., 41 (2003), pp. 364–381, <https://doi.org/10.1137/S0036142901385593>.
- [6] J.-P. KAUTHEN AND H. BRUNNER, *Continuous collocation approximations to solutions of first kind Volterra equations*, Math. Comp., 66 (1997), pp. 1441–1459, <https://doi.org/10.1090/S0025-5718-97-00905-8>.
- [7] H. LIANG AND H. BRUNNER, *On the convergence of collocation solutions in continuous piecewise polynomial spaces for Volterra integral equations*, BIT, 56 (2016), pp. 1339–1367, <https://doi.org/10.1007/s10543-016-0609-x>.